



Breast Cancer Molecular Subtypes in Relation to Age, Stage and Grade among Sudanese Women Patients in Khartoum Oncology Hospital (2013 – 2017)

Mawahib Ahmed Elawad Abu Elgasim^{1*}, Heba Mustafa Abusin² and Manahil Mohamed Nouri³

¹Department of Community Medicine, Faculty of Medicine, Omdurman Islamic University, Sudan.

²Department of Oncology, Faculty of Medicine, Omdurman Islamic University, Sudan.

³Saudi Board of Preventive Medicine, Public Health Directorate- Jeddah, Saudi Arabia.

Authors' contributions

This work was carried out in collaboration among all authors. Authors MAEAE and HMA designed the study. Authors MAEAE and HMA revised the tools for data collection. Authors MAEAE and HMA maintained the quality of data collection in the field work. Author MAEAE drafted the first draft of the manuscript. Authors MAEAE revised the manuscript for grammar and spelling. All authors read and approved the final manuscript

Article Information

Editor(s):

(1) Dr. Krishnangshu Bhanja Choudhury, Assistant Professor, Department of Radiotherapy, R.G. Kar Medical College and Hospital, Kolkata, India.

Reviewers:

(1) E. Martha Ruiz-Tachiquin, Unidad de Investigación Médica en Genética Humana, México.

(2) Alessandro Borgognone, S. Eugenio Hospital, Italy.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48935>

Original Research Article

Received 04 March 2019

Accepted 19 May 2019

Published 27 May 2019

ABSTRACT

Background: There is a lack of data regarding molecular subtypes of breast cancer among Sudanese female patients. Moreover to the best of our knowledge no study had correlated the molecular subtypes with age, stage and grade till date. This study aimed to determine breast cancer molecular subtypes among Sudanese women in relation to age, clinical stage and grade and to compare the findings to other related researches.

Methods: Retrospective Study of (255) Sudanese female patients diagnosed with breast cancer from September 2013 to August 2017 in Khartoum Specialized Oncology Center, an information

sheet was used to collect data from patients medical records.

Results: The results demonstrated that, the most commonly detected molecular subtype was luminal B (34.9%), followed by triple negative and HER-2 enriched, (31.4%) and (19.2%) respectively. The least common subtype was luminal A, (14.5%). Additionally, over half of the patients (54.4%) were classified as grade 3 and (22.4%) were diagnosed as stage IIIb.

Conclusion: The main conclusion that can be drawn is that, luminal B was the most commonly diagnosed subtype and it was more common in younger age group in comparison to luminal A. The majority of the patients in the study were grade 3 and stage IIIb and were diagnosed between the age of 41 and 50. Moreover, triple negative subtype was high among Sudanese women and was associated with advance grade breast cancer.

Keywords: Subtypes; breast cancer; Sudanese; women.

1. INTRODUCTION

Breast cancer is the top cancer in females leading to 327,000 deaths annually [1,2]. In 2018, WHO reported nearly 2.1 million newly diagnosed female with breast cancer [3]. The global cancer burden of breast cancer is 11.6% [4]. According to the latest data of Globocan 2018, breast cancer ranked the most frequent cancer among women in Sudan and a total of 5,677 (36.69%) Sudanese women were newly diagnosed with breast cancer [5].

Several clinical and pathological parameters are used to classify the breast cancer subtypes, namely; lymph node (LN) status, tumor size, tumor grade, age, menopausal status, clinical and pathological stage, histological type as well as estrogen (ER), progesterone (PR) and HER-2 receptors [6]. Detecting the molecular subtype of breast cancer is a recent and advance technique for early detection and evaluation of the prognosis and management of breast cancer [7]. Thus, management options in Sudan are adapted according to breast cancer molecular subtype. The treatment guidelines in Sudan are based on the National Comprehensive Cancer Network (NCCN) and European Society for Medical Oncology(ESMO) guidelines which includes; screening, diagnosing, staging, management of local/loco- regional disease, endocrine treatment in pre- and post-menopausal patients, chemotherapy and HER2-directed therapy [8,9]. Undoubtedly molecular subtypes could provide promising prognostic and predictive information and may help identify new therapeutic targets. However, it is important to understand their limitations and to evaluate their role in improving breast cancer prognosis beyond the traditional expected outcomes in a practical and cost-effective manner [10].

Breast cancer is divided into four major molecular subtypes; the most frequent type is luminal A which is present in 50-60% of breast cancer patients and has the best prognosis rates. The second subtype, luminal B is characterized by higher histological grade and is found in 15-20% of patients [11]. The third subtype triple negative breast cancer (TNBC) which varies according to patient race and ethnicity. Furthermore, it impacts patient's survival rates and may affects management options [12]. The fourth molecular subtype, HER2-enriched (previously the HER2+/ER-subtype) is characterized by aggressive behavior [13]. There is a variation in the prevalence of molecular subtypes worldwide. Luminal A tumors were the most common tumors among some Middle East countries, namely; Saudi Arabia [14], Jordan [15] and Egypt [16] and in a number of western countries; Italy [17], Germany [18] and Atlanta (America) [19]. Regarding Asian countries, studies revealed that luminal A was high among Iranian patients [20], while Luminal B was more commonly found in Japanese [21] and Pakistani patients [22].

Studies in African countries unveiled that ; in Morocco (North Africa), Luminal B was the most common subtype [23], while triple negative had high prevalence rates in Nigeria (West Africa) [24], Uganda (sub-Saharan Africa) [25], Sudan and Eretria (North East and East Africa) [26]. Regarding race, TNBC was found to be the most prevalent subtype in African-American women [27]. Another study conducted to compare molecular subtypes between Sudanese and German women showed that triple-negative subtype was more frequent in Sudanese than German women [28].

Few studies have investigated breast cancer molecular subtypes in Sudan. Furthermore, there

is no research relating molecular subtypes to age, stage and grade of breast cancer to date. Thus, in our research we tried to classify patients according to breast cancer molecular subtypes.

The aim of this study is to determine breast cancer molecular subtype among Sudanese women in relation to age, clinical stage and grade and to compare the results to other related researches.

2. MATERIALS AND METHODS

2.1 Study Design and Setting

This is a retrospective study of histologically confirmed Sudanese women with breast at Khartoum Specialized Oncology center, in the period from September 2013 to August 2017.

Khartoum Specialized Oncology center is a specialized tertiary hospital that offers chemotherapy and radiotherapy for cancer patients, located in Khartoum State, Capital of Sudan. Entitled patients are referred from all over Sudan and hence, this study's sample is therefore representative of the Sudanese population.

2.2 Sampling

255 medical records of histologically confirmed breast cancer patients were included in the study.

2.3 Data Collection Tools

An information sheet has been used for data collection from patient's medical records.

The data retrieved included the following:-

-Patients age when diagnosed was distributed into two groups; younger age group (50 years or less) and older age group (above 50 years).

-Molecular subtypes which were identified by Immunohistochemical (IHC) markers (ER/PR/HER2). Four subtypes were defined, namely; luminal A, luminal b, triple negative and HER-2 enriched.

-Breast cancer grades (I, II and III) were detected using Nottingham Bloom-Richardson grading system [29].

-The Clinical stage of the disease was estimated from the clinical examination and was classified

according to American Joint Committee (AJC) and TNM classification [30].

2.4 Statistical Analysis

Data was entered and analyzed using the Statistical Package for the Social Sciences (version 21.0). Chi-square Test was used to evaluate the correlation between molecular subtypes and age, stage and grade were used. The results were considered significant when p (degree of significance) was less than 0.05.

3. RESULTS

A total of 255 records of female patients diagnosed with breast cancer were enrolled in the study. The mean patient's age at diagnosis was 48.8 ± 11.3 years. The majority of the patients (78.2%) were diagnosed with breast cancer before the age of 60. However, only (20.8%) of the cases were diagnosed above the age of 60 (Table 1).

The most commonly detected molecular subtype was luminal B (34.9%), followed by triple negative and HER-2 enriched, (31.4%) and (19.2%), respectively. The least common subtype was luminal A (14.5%) (Fig. 2). The vast majority of patients (22.4%) were stage IIIb, followed by stage IIa and IIb (21.1%). Furthermore, (15.7%), (8.6%) and (6.7%) of the cases were diagnosed as stage IIIa, IV and I, respectively. A small minority, (4.3%) were diagnosed as stage IIIc (Table 2). (54.4%) of the breast cancer patients were diagnosed as grade 3, while (39.2%) were classified grade 2 and only (9.4%) were diagnosed as grade 1 (Fig. 1).

Out of the (255) cases, 135 cases (52.9%) were in the younger age group (≤ 50) and 120 cases (47.1%) were in the older age group (>50 years). Most cases (34.9%) (n=89) were classified as luminal B subtype, (51.7%) of which were in the younger age group, while (48.3%) (n=43) were in the older age group. Moreover, (31.4%) (n=80) of the cases were classified as triple negative subtype, (56.3%) (n=45) were in the younger age group and (47.7%) were in the older age group. (14.5%) (n=37) of the cases were classified as luminal A subclass, (37.8%) (n=14) were in the younger age group and (62.7%) (n=23) were in the older age group. (19.2%) (n=49) were classified as HER-2 enriched subclass, (61.2%) (n=30) were in the younger age group and (38.8%) (n=19) were in the older age group.

However, the relationship between molecular subtype of breast cancer and patients age at diagnosis was not statistically significant ($p=0.162$) (Table 3).

Concerning the relationship between molecular subtypes and breast cancer stages, there was no significant association ($p=0.257$). For patients with HER-2 enriched molecular subtypes, the frequency of stage I, IIa, IIb, IIIa, IIIb, IIIc and IV breast cancer, were (2%), (16%), (24.4%), (18.3%), (26.5), (2%), and (10.2%), respectively. Regarding patients with Luminal A molecular subtypes, stage I, IIa, IIb, IIIa, IIIb, IIIc and IV breast cancer, were (16%), (32%), (18.9%), (10.8%), (16.2%) and (2.7%), respectively. For patients with Luminal B molecular subtypes, stage I, IIa, IIb, IIIa, IIIb, IIIc and IV breast cancer, were (5.6%), (22.4%), (19%), (13.4%), (22.4%), (3.4%) and (13.4%), respectively. For patients with Triple negative molecular subtypes, stage I, IIa, IIb, IIIa, IIIb, IIIc and IV breast cancer, were (6.3%), (17.5%), (22.5%), (18.8%), (22.4%), (7.5%) and (5%), respectively (Table 4).

The molecular subtypes were found to be significantly associated with breast cancer grade ($p=0.012$). Luminal B frequency of grade 1, 2 and 3 was (3.5%), (16.1%) and (15.3%), respectively, while the distribution of Luminal A was (2.4%) for grade 1, (7.8%) for grade 2 and (4.3%) for grade 3. (19.2%) of triple negative were grade 3, (9.8%) were grade 2, while only (2.4%) were grade 1. (12.5%) of Her-2 enriched were grade 3, (5.5%) were grade 2 and (1.2%) were grade 1.

4. DISCUSSION

In the current study most of the patients were fifty years and above (52.9%), while patients under fifty were (47.1%). Furthermore, the vast majority of patients were diagnosed with breast cancer between the age of 41 and 50. A similar conclusion was suggested by a study done in Nigeria [24]. According to the present study most of the cases were grade 3 (54.4%) and (39.9%) were grade 2. However, a different finding was reported by another study, which revealed that grade 2 breast cancer was more frequent among Sudanese, German and Nigerian female patients, (54.6%), (60%) and (48.57%) respectively, while the percentage of stage 2 cancer in Sudanese women was (41.8%), German women (22%) and (43.57%) for Nigerian women. The latter finding could be

partially explained by the late detection of cancer cases in Sudan.

From the results, (22.4%) of female patients were diagnosed as stage IIIb cancer, that may be attributed to lack of awareness, difficult accessibility services and absence of cancer screening programs.

With regard to molecular subtypes, luminal A is found in (50%-60%) of the patients and luminal B in (15%-20%), however, in our group the majority of the cases were classified as luminal B (34.9%) and only (14.5%) were luminal A [11]. A different conclusion was obtained from a study done in middle east countries; where percentage of patients with luminal A subtype in Saudi Arabia, Jordan and Egypt was (58.5%), (60%) and (45%), respectively [14,15,16]. Another similar study conducted in some western countries showed the following; Italy (34%), Germany (44.7%) and Atlanta (51.1%), [17,18,19]. Furthermore, the prevalence of Luminal A subtype was (63.8%) according to an Iranian study [20]. In Japan, Pakistan and Morocco, the percentage of luminal B was (71%), (69%) and (41.8%), respectively [21,22,23] which was higher than our finding. The variation in the results could be linked to the distribution of the different age groups in the studies.

The prevalence of Triple negative breast cancer subtype (TNBC) in Nigerian women was (26.53%) and (21.2%) among African-American women [24,27], nonetheless, a lower result was attained by our study (31.4%). A slightly comparable value to our finding was found in Ugandan women (34%) and Sudanese- Eritrean women (34.5%), [25,26,28].

In our examination, HER-2 enriched was found in (19.2%) of the breast cancer patients. Nevertheless, a lower finding was cited by another study; the HER-2 enriched frequency among Jordanian women was (12%) [15], Sudanese women (15.7%), German women (6.8%) [28], Sudanese Eritrean women (16%) and (9.2%) among Moroccan women [23,26].

Luminal B subtype was present in (51.7%) of patients fifty years or less. whereas, the percentage in older patients (i.e. above fifty) was (48.3%). On the other hand, Luminal A subtype prevailed in patients over fifty years old (62.2%), while (37.8%) was detected in younger ages (fifty years or less). A similar pattern of result was obtained in Jordan, where (72%) of luminal A subtype were above fifty years old [15].

Table1. Distribution of patients according to the age of breast cancer at diagnosis

| Age group | Frequency | Percent |
|-----------|-----------|---------|
| <30 | 7 | 2.7 |
| 30-40 | 54 | 21.2 |
| 41-50 | 74 | 29.0 |
| 51- 60 | 67 | 26.3 |
| >60 | 53 | 20.8 |
| Total | 255 | 100.0 |

Table 2. Distribution of patients according to the stages of breast cancer at diagnosis

| Stage | N | % |
|-------|-----|-------|
| I | 17 | 6.7 |
| Ila | 54 | 21.2 |
| Ilb | 54 | 21.2 |
| Illa | 40 | 15.7 |
| IIIb | 57 | 22.4 |
| IIIc | 11 | 4.3 |
| IV | 22 | 8.6 |
| Total | 255 | 100.0 |

Table 3. The relationship between molecular subtype of breast cancer and age at diagnosis

| Molecular subtype | Cases ≤50 years N(%) | Cases >50 years N(%) | Total N(%) | P-Value |
|-------------------|----------------------|----------------------|------------|---------|
| HER-2 enriched | 30(61.2) | 19(38.8) | 49(19.2) | 0.162 |
| Luminal A | 14(37.8) | 23(62.7) | 37(14.5) | |
| Luminal B | 46(51.7) | 43(48.3) | 89(34.9) | |
| Triple negative | 45(56.3) | 35(47.7) | 80(31.4) | |
| Total | 135(52.9) | 120(47.1) | 255(100) | |




Fig. 1. Distribution of patients according to the grade of breast cancer at diagnosis




Fig. 2. Distribution of patients according to the molecular subtypes of breast cancer at diagnosis

Table 4. The relationship between molecular subtype of breast cancer and stage at diagnosis

| Molecular subtype | | Stage | | | | | | | Total | P value |
|-------------------|-----|-------|--------|--------|--------|--------|-------|--------|--------|---------|
| | | I | II a | II b | III a | III b | III c | IV | | |
| HER-2 enriched | N | 1 | 8 | 12 | 9 | 13 | 1 | 5 | 49 | 0.257 |
| | (%) | (2) | (16) | (24.4) | (18.3) | (26.5) | (2) | (10.2) | (19.2) | |
| Luminal A | N | 6 | 12 | 7 | 4 | 6 | 1 | 1 | 37 | |
| | (%) | (16) | (32) | (18.9) | (10.8) | (16.2) | (2.7) | (2.7) | (14.5) | |
| Luminal B | N | 5 | 20 | 17 | 12 | 20 | 3 | 12 | 89 | |
| | (%) | (5.6) | (22.4) | (19) | (13.4) | (22.4) | (3.4) | (13.4) | (34.9) | |
| Triple negative | N | 5 | 14 | 18 | 15 | 18 | 6 | 4 | 80 | |
| | (%) | (6.3) | (17.5) | (22.5) | (18.8) | (22.5) | (7.5) | (5) | (31.4) | |
| Total | | 17 | 54 | 54 | 57 | 40 | 11 | 22 | 255 | |
| | | (6.7) | (21.2) | (21.2) | (22.4) | (15.7) | (4.3) | (8.6) | (100) | |

Table 5. The relationship between molecular subtypes of breast cancer and histological grades

| Molecular subtype | | N | Grade | | | Total | P-Value |
|-------------------|-----|--------|--------|--------|--------|-------|---------|
| | | | 1 | 2 | 3 | | |
| HER-2enriched | N | 3 | 14 | 32 | 49 | 0.012 | |
| | (%) | (6.1) | (28.6) | (65.3) | (19.2) | | |
| Luminal A | N | 6 | 20 | 11 | 37 | | |
| | (%) | (16.2) | (54.1) | (29.7) | (14.5) | | |
| Luminal B | N | 9 | 41 | 39 | 89 | | |
| | (%) | (10.1) | (46.1) | (43.8) | (34.9) | | |
| Triple negative | N | 6 | 25 | 49 | 80 | | |
| | (%) | (7.5) | (31.2) | (61.3) | (31.4) | | |
| Total | | N | 24 | 100 | 131 | 255 | |
| | | (%) | (9.4) | (39.2) | (51.4) | (100) | |

Triple negative and HER-2 enriched subtypes were more prominent in the younger ages (fifty years or less), (56.3%) and (61.2%), respectively. In patients over fifty years old, Triple negative subtype was detected in (43.7%) of the patients and Her-2 enriched in (38.8%). This was consistent with a study done among Sudanese and German women [28].

(7.8%) of Luminal B breast cancer subtype was stage IIIb, while only (2.4%) of luminal A was stage IIIb, which indicates that, Luminal B is more aggressive than luminal. With regard to cancer grades, the majority of luminal A breast cancer subtype (7.8%) was Grade 2. On the other hand, in Grade 3, luminal B prevailed over luminal A, (15.3%) and (4.3%) respectively. Therefore, it's suggested that, luminal B is associated with higher histological grades [11].

Another finding was that, high frequencies for grade 3 cancer was registered by Triple negative and Her-2 enriched Subtypes (19.2%) and (12.5%), respectively. This implies that Triple negative and Her-2 enriched are associated with aggressive and advanced stages of breast cancer [13].

5. CONCLUSION and RECOMMENDATION

Most of the Sudanese women were diagnosed with breast cancer between (41-50) years old. Moreover, most of them were presented with grade 3 and stage IIIb breast cancer.

Luminal B was the most prevailed molecular subtypes, followed by Triple negative.

Luminal A was more common among old age groups (over fifty). However, Her-2 enriched and TNBC subtypes were mostly Grade 3 and prevailed among younger Sudanese women.

Detecting the subtype of breast cancer is not only essential for following disease prognosis but also for the management of the breast cancer.

Breast cancer screening programs and self-examination are highly recommended for the early detection of the disease.

The effect of determining molecular subtype on survival rates is an issue for future research to explore.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Ethical approval was obtained from Institutional review board of Omdurman Islamic university- Faculty of Medicine. Data were collected after taking the necessary agreement from Khartoum State Ministry of Health and Khartoum Specialized Oncology Center.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. ZiQi Tao, Aimin Shi, Cuntao Lu, Tao Song, Zhengguo Zhang, Jing Zhao. Breast cancer: Epidemiology and etiology. *Cell Biochem Biophys*. 2015;72:333–38.
2. Hadi MA, Al Madani R, Abu Arida L, Al Ghareeb B. Breast cancer age in developing countries: The narrowing gap. *Clin Surg*. 2018;3:2074.
3. Freddie Bray, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries, *CA CANCER J CLIN*. 2018; 68:394–424.
4. WHO, Latest global cancer data. Available: <https://www.who.int/cancer/PRGIobocanFinal.pdf> [September, 2018]
5. WHO, the Global Cancer Observatory; 2019. Available: <http://gco.iarc.fr/today/data/factsheets/populations/729-sudan-fact-sheets.pdf> [March, 2019]
6. Gannon LM, Cotter MB, Quinn CM. The classification of invasive carcinoma of the breast. *Expert Rev Anticancer Ther*. 2013; 13: 941-954. (PMID: 23984896)
7. Jane Date C. Hon, Baljit Singh, Aysegul Sahin, Gang Du, Jinhua Wang, Vincent Y. Wang, Fang-Ming Deng, David Y. Zhang, Marie E. Monaco, Peng Lee Am *J Cancer Res*. 2016;6(9):1864–1872. [Published online 2016 Sep 1] [PMCID: PMC5043099]
8. National Comprehensive Cancer Network® (NCCN®), Guidelines for Breast Cancer Patients Version 4; 2018. Available: www.nccn.org/patients [February, [February, 8, 2019]
9. Senkus E, Kyriakides S, Ohno S, et al. Primary breast Cancer, *ESMO Clinical*

- Practice Guidelines for diagnosis, treatment and follow-up, *Annals of Oncology*. 2015;26(suppl_5):8-30. Available:<https://doi.org/10.1093/annonc/mdv298>
10. Emad A. Rakha, Jorge S. Reis-Filho, et al. Breast cancer prognostic classification in the molecular era: The role of histological grade *Breast Cancer Res*. 2010;12(4):207. [Published online 2010 Jul 30] DOI: 10.1186/bcr2607 [PMCID: PMC2949637]
 11. Ozlem Yersal, Sabri Barutca. World, biological subtypes of breast cancer: Prognostic and therapeutic implications. *J Clin Oncol*. 2014;5(3):412–424. [Published online 2014 Aug 10] DOI: 10.5306/wjco.v5.i3.412
 12. Ding YC, Steele L, et al. Molecular subtypes of triple-negative breast cancer in women of different race and ethnicity, *Oncotarget*. 2019;10(2):198-2. DOI: 10.18632/oncotarget.26559
 13. O'Sullivan CC, Loprinzi CL, Haddad TC. Updates in the evaluation and management of breast cancer. *Mayo Clinic Proceedings*. 2018;93(6):794-807. Available:<https://doi.org/10.1016/j.mayocp.2018.03.025>
 14. Al Tamimi DM, Shawarby MA, Ahmed A, Hassan AK, AlOdaini AA. Protein expression profile and prevalence pattern of the molecular classes of breast cancer—A Saudi population based study. *BMC Cancer*. 2010;10(1):223. Available:<https://doi.org/10.1186/1471-2407-10-223> [PubMed: 20492711 PMID: PMC2880995]
 15. Shomaf M, Masad J, Najjar S, Faydi D. Distribution of breast cancer subtypes among Jordanian women and correlation with histopathological grade: Molecular sub classification study. *JRSM Short Reports*. 2013;4(10). Available:<https://doi.org/10.1177/2042533313490516>
 16. Bodour Salhia, Coya Tapia, et al. Molecular subtype analysis determines the association of advanced breast cancer in Egypt with favorable biology, *BMC Women's Health*. 2011;11:44. Available:<https://doi.org/10.1186/1472-6874-11-44>
 17. Caldarella A, Buzzoni C, Crocetti E, Bianchi S, Vezzosi V, Apicella P, Biancalani M, Giannini A, Urso C, Zolfanelli F, Paci E. Invasive breast cancer: A significant correlation between histological types and molecular subgroups. *J Cancer Res Clin Oncol*. 2013; 139(4):617-23.
 18. Fabian Riedel, Peter Sinn. Prognosis of breast cancer molecular subtypes in routine clinical care: A large prospective cohort study, *BMC Cancer*. 2016;16:734. Available:<https://doi.org/10.1186/s12885-016-2766-3>
 19. Gaudet MM, Press MF, et al. Risk factors by molecular subtypes of breast cancer across a population-based study of women 56 years or younger, *Breast Cancer Res Treat*. 2011;130(2):587-97. DOI: 10.1007/s10549-011-1616-x [Epub 2011 Jun 11]
 20. Kadivar M, Mafi N, Joulaee A, Shamshiri A, Hosseini N. Breast cancer molecular subtypes and associations with clinicopathological characteristics in Iranian women 2002-2011. *Asian Pac J Cancer Prev*. 2012;13:1881–1886.
 21. Shibuta K, Ueo H, Furusawa H, Komaki K, Rai Y, Sagara Y, Kamada Y, Tamaki N, The relevance of intrinsic subtype to clinicopathological features and prognosis in 4,266 Japanese Women with Breast Cancer. *Breast Cancer*. 2011;18(4):292-98. Available:<https://doi.org/10.1007/s12282-010-0209-6> [PMid: 20571962]
 22. Hashmi AA, Aijaz S, Khan SM, Mahboob R, Irfan M, Zafar NI, Nisar M, Siddiqui M, Edhi MM, Faridi N, Khan A. Prognostic parameters of luminal A and luminal B intrinsic breast cancer subtypes of Pakistani patients. *World J. Sur Oncol*. 2018;16(1):1. Available:<https://doi.org/10.1186/s12957-018-0170-7>
 23. El Fatemi H, Chahbouni S, Jayi S, Mounna K, Melhouf MA, Bannani A, Mesbahi O, Amarti A. Luminal B tumors are the most frequent molecular subtype in breast cancer of North African women: An immunohistochemical profile study from Morocco, *Diagn Pathol*. 2012;7:170. DOI: 10.1186/1746-1596-7-170
 24. Kenneth A. Omoruyi, Martins A. Nnoli, Godwin A. Ebughe, Godstime Irabor, Edoise M. Isiwele. Prevalence of molecular subtypes of breast carcinoma in university of Calabar teaching hospital using immunohistochemistry as surrogates for Intrinsic DNA gene characteristics, *IOSR*

- Journal of Dental and Medical Sciences (IOSR-JDMS) e-ISSN: 2279-0853, p-ISSN: 2279-0861. 2018;17(2)Ver. 12:64-73.
25. Moses Galukande, Henry Wabinga, Florence Mirembe, Charles Karamagi, Alexzander Asea. Molecular breast cancer subtypes prevalence in an indigenous Sub Saharan African population Pan Afr Med J. 2014;17:249.
DOI: 10.11604/pamj.2014.17.249.330
[PMCID: PMC4189896]
 26. Sengal AT, Haj-Mukhtar NS, Elhaj AM, Bedri S, Kantelhardt EJ, Mohamedani AA. Immunohistochemistry defined subtypes of breast cancer in 678 Sudanese and Eritrean women; hospitals based case series, BMC Cancer. 2017;17(1):804.
DOI: 10.1186/s12885-017-38054
[PMID: 29191181]
 27. Chukwuemeka U. Ihemelandu, et al. Molecular breast cancer subtypes in premenopausal and postmenopausal African-American women: Age-specific prevalence and survival. J Surg Res. 2007; 143(1):109–118.
 28. Sengal AT, Haj Mukhtar NS, Vetter M, Elhaj AM, Bedri S, Hauptmann S, Thomssen C, Mohamedani AA, Wickenhauser C, Kantelhardt EJ. Comparison of receptor-defined breast cancer subtypes between German and Sudanese women: A facility-based cohort study. J Glob Oncol. 2018;4: 1-12.
DOI: 10.1200/JGO.2017.010082
[Epub 2017 Aug 4]
[PMID: 30241184]
 29. Elston C, Ellis IO. Pathological prognostic factors in breast cancer: The value of histological grade in breast cancer: experience from a large study with long-term follow-up. Histopathology. 2002;41: 154–61.
 30. The American Joint Committee on Cancer: The 7th edition of the AJCC cancer staging manual and the future of TNM. Edge SB, Compton CC Ann Surg Oncol. 2010; 17(6):1471-4.

© 2019 Elgasim et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48935>*