

Compositional Analysis of Sweet Italian (*Capsicum annuum*) and Habanero (*Capsicum annuum*) Peppers Consumed in Benue State, Nigeria

Ngunengen Igbakaa^a, Abraham Tartenger Girgih^a,
Dinnah Ahure^a and Pius Oteikwu Ejoha^{a*}

^a Department of Food Science and Technology, University of Agriculture, Makurdi, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100240>

Original Research Article

Received: 01/04/2023
Accepted: 03/06/2023
Published: 19/08/2023

ABSTRACT

This study investigates the physico-chemical properties and Fourier transform infrared spectral analyses of Sweet Italian (*Capsicum annum*) and Habanero (*Capsicum chinense*) pepper varieties. The two varieties of pepper were harvested fresh and dried using local drying equipment and ground into powder using a laboratory blender. The proximate composition results showed that the Sweet Italian pepper (SIP) possessed higher moisture (8.89%), crude fat (5.33%), crude protein (9.32%) and carbohydrate (51.5%) contents than the Habanero pepper (HNP). SIP showed higher content of the major (Na, K, Mg, Ca & P) and trace (Fe, Zn, Cu and Mn) minerals when compared to the HNP. The results of the infrared spectrum analysis of the two varieties of SIP and HNP showed that these pepper varieties had similar infrared spectra lines that are dominated by alcohol/phenol groups, alkanes, alkenes and amine groups, corresponding to the frequencies in the range 3435.44-675.11 cm⁻¹ for the phenols, 1037, 1379-1458 cm⁻¹ for the alkanes group, 1637.62-

*Corresponding author: E-mail: piusejoha@yahoo.com;

920.08 cm^{-1} for the alkenes and 719.47-817.57 cm^{-1} for amine groups. The result indicated that the two varieties of peppers SIP and HNP having similar spectra lines, may also possess similar structural properties.

Keywords: Proximate composition; mineral; FTIR; sweet; Italian; Habanero and pepper.

1. INTRODUCTION

Proximate composition evaluates the content of the six macronutrients in foods including moisture, protein, fats, fiber, ash and carbohydrates. The significance of determining the moisture content of food products helps a lot of people to know about the food product characteristics, including its physical appearance, texture, taste and weight, in addition to the food product shelf-life/stability, freshness and resistance to bacterial contamination [1]. Moisture content in dried foods should be $\leq 12\%$ for prolong shelf-life and storage stability [2]. Protein is very essential for body building and repair of worn out body tissues, in addition, its presence in peppered food product will help reduce the problem of protein-energy malnutrition for all ages in a population suffering this challenge [3].

The advantage of having low fat content in pepper is that its keeping quality will be enhanced as lipid peroxidation will minimally occur. In contrast, high amount of fats in pepper could provide energy for physical and physiological activities but could result in rancidity. Excess fats could also be extracted for use as essential oil for treatment of skin diseases and provide relieve from swollen body parts [4]. Crude fibers are broadly defined as non-starch polysaccharides and possess varying physicochemical properties depending on their chemical composition. Fiber has been reported to have an attenuating effect on appetite by increasing satiety thus reducing the amount of food consumed by individuals [5]. The diets rich in fiber such as fruits and vegetables have been reported to have a positive effect on health improvement since regular consumption has been linked to decreased incidences of several diseases such diabetes and cardiovascular events [6]. Fibre is an important nutrient in the body useful for colon nutrition for ease of fecal passage through gastrointestinal tract for egestion [7]. Ash is a measure of mineral content in food material [8]. Most of the energy in a diet comes from carbohydrates. Carbohydrates provide the major source of the most easily available energy in the diet. The Institute of

Medicine recommends that we should consume between 45–65% of total calories which should come from carbohydrates [9].

Mineral content of food are the essential macronutrient that plays a central role in the normal regulation of blood pressure [10] and to regulate the body homeostasis via the Na^+/K^+ pump which also promotes the proper functioning of the brain [11]. It is also involved in the regulation of the fluid balance of the body and hence, influences the cardiac output. (sodium and potassium). Calcium is one of the important macronutrients relevant in the building of strong bones and teeth [12]. Calcium is also very essential in muscle contraction, oocyte activation, blood clotting, nerve impulse, transmission and in regulation of heart beat [13]. It is also important in circulatory system, extracellular fluid, muscle and other tissues is critical for mediating vascular contraction and vasodilatation, muscle function, nerve transmission, intracellular signaling and hormonal secretion and the body to obtains calcium from the bone tissues which serves as a reservoir for source for these critical metabolic needs through the process of bone remodeling [14].

Magnesium is very important in the body because of its involvement in the respiratory process in the body [15]. Literature has provided extensive evidence of widespread magnesium deficiency the need for magnesium replenishing its content in diverse medical conditions. Magnesium is an essential element required as a cofactor for over 300 enzymatic reactions and thus important in the biochemical functioning of numerous metabolic pathways. Iron is an essential component of hemoglobin and myoglobin and thereby facilitates the transport, transitional tissue storage and cellular use of oxygen. It also has important roles in cytochromes within mitochondria, mediating the transfer of electrons in the electron transport chain [16-18]. Zinc is very important in the body and helps the immune system fight off the invading bacteria and viruses. During pregnancy, infancy and childhood, the body needs zinc for proper growth and development. Zinc also helps

wounds heal and is important for proper senses of taste and smell [12]. Copper is very important in the absorption of iron and therefore has a very important role in preventing anemia. This trace element helps in the use of iron for the transport of hemoglobin. Copper helps to form collagen and melanin and is indispensable in diets for skin and hair care [19]. Manganese also helps the body to utilize a number of vitamins such as choline, thiamine and vitamin C and E and ensures proper liver function [19].

FT-IR spectra is used recently in research for fast detection of the natural active compounds in plants and animals tissues. The utilization of Fourier transform-infrared spectroscopy (FTIR) as an essential technique has been widely employed [20] and becomes handy for research into medicinal plants, [21,22] human disease [23-28] also forensic science [29-31] and food characterization due to its nondestructive nature as well as allowance for easy identification [32,33]. The present study sought to encourage the detection of components and other relative compounds in the two pepper varieties in short time at minimal cost using FT-IR detection.

2. MATERIALS AND METHODS

2.1 Source of Materials

The two varieties of pepper, Sweet Italian (*Capsicum annum*) and Habanero (*Capsicum annum*) were harvested fresh from local farms in Makurdi metropolis in their ripened stage and dried using local drying equipment and milled into powder using a laboratory blender.

2.2 Processing of Pepper Powder

The modified method of [34] was used for pepper powder production as shown in Fig. 1. The red pepper were selected to uniform sizes, shapes, and without any defect on visual inspection and thoroughly cleaned before manually sorting. The sorted red peppers were washed in cold water to remove soil and dust particles. The thoroughly cleaned samples were manually graded on the basis of their size. Washed red peppers were sliced with knives as approximate sizes of 15 mm x 15 mm of uniform slices with thickness of 2–4 mm. After slicing, the slices were blanched with hot water at 95°C for 3-5 min. The method of blanching is similar to that of [35,36]. The blanched slices were dried using locally fabricated electric dryer, where the red pepper slices were spread on the shelves of the drying

bin and the hot air was passing through these dryer upward from the electric air collectors at the temperatures between 65-70 °C. Drying time and final moisture content for product was controlled. Also, the red pepper slices were shifted alternatively inside the electric bin in order to give the same chance for the red pepper slices to have the same drying conditions. The red pepper was ground and kept until used.




Fig. 1. Flow chart for pepper powder preparation

Source: [34]

2.3 Functional Properties of Sweet Italian and Habanero Peppers

2.3.1 Determination of bulk density (BD) of the pepper powders

The method of [37] was used to determine the bulk density of the pepper powders. A 10ml capacity graduated measuring cylinder was gently filled with the sample and the bottom of the cylinder was tapped on the laboratory bench several times until there no further diminution of the sample level after filling to the 10ml mark. Bulk density was calculated by using the formula,

$$\text{Bulk density (g/ml)} = \frac{\text{Weight of sample (g)}}{\text{Volume of sample (ml)}} \dots [1]$$

2.3.2 Determination of water absorption capacity of the pepper powders

Water absorption capacity was determined using the procedure of [38]. One gram of the sample was mixed with 10 ml distilled water for 5 min on a magnetic stirrer. The mixture was centrifuged at 3500 rpm for 30 min and the volume of the supernatant noted. WAC was calculated using as thus;

$$\text{Water absorption capacity} = \frac{\text{volume of distilled water}}{\text{weight of sample used}} \times 100$$

2.3.3 Determination of oil absorption capacity of the pepper powders

One gram of sample was weighed, 10 ml of vegetable oil of a known density (0.99 mg/ml) was added to the sample and the mixture stirred on a magnetic stirrer at 1000 rpm for 5 min. The mixture was centrifuged at 3500 rpm for 30 min and the supernatant removed and measured with 10 ml measuring cylinder [38]. The OAC was calculated

$$\text{Oil absorption capacity} = \frac{\text{volume of oil absorbed} \times \text{density}}{\text{weight of sample used}} \times 100$$

2.3.4 Determination of the foaming capacity of the pepper powders

Foaming capacity (FC) was determined by the method of [39]. One gram flour sample was added to 50 ml distilled water at ambient temperature in a graduated cylinder. The suspension were mixed and shaken for 5min to foam. The volume of foam at 30secs after whipping was expressed as foam capacity using the formula:

Where

$$\text{AW} = \text{after whipping,} \\ \text{BW} = \text{before whipping.}$$

2.3.5 Determination of swelling index of the pepper powders

The method of [40] was used to determine the swelling power of the experimental flours in duplicate. One gram of flour samples was mixed in a varl-whirl mixer with 10ml of distilled water for 30 seconds and allowed to stand for 30 minutes at ambient room temperature (29⁰C) before being centrifuged at 5,000 rpm for 30 minutes. Measuring out the volumes of the

supernatants was used to find the volumes of the remaining absorbed liquids. Multiplication of the respective absorbed volumes by the respective liquid density (mass/volume) was used to get the expression in g liquid/g sample.

2.3.6 Determination of gelation temperature of the pepper powders

The water absorption capacity was done according to the method described by [38]. A suspension of 10% of the sample in a test tube was prepared and the aqueous suspension in a boiling water bath, with continuous stirring was heated. The temperature 30secs after gelatinization was visually noticed as the gelatinization temperature was recorded.

2.3.7 Reconstitution index

Reconstitution Index was done following the method described by [38]. 5 g of pepper powder each was dissolved in 40ml of boiling water. The mixture was agitated for 90 seconds and transferred into a 50 ml of the cylinder and the volume of the sediment recorded after settling for 30 min.

2.3.8 Wettability

Wettability was determined according to the method described by [38]. 10 g of pepper was weighed into a clean, dried measuring cylinder (10 ml). The cylinder was inverted and clamped at height of 10cm from the surface of a 600 ml beaker containing 500ml distilled water. The flour in the cylinder was gradually spread on the surface of water at a moderate speed. The time taken for the sample to completely wet was noted as wettability.

2.4 Determination of Proximate Composition of Pepper Powders

2.4.1 Determination of moisture content of pepper powders

The moisture content was determined by hot air oven method as described by [37]. Empty crucible was weighed and 2 g of the sample was transferred into the crucible. This was taken into the hot air oven and dried for 24 hours at 100⁰C. The loss in weight was regarded as moisture content and expressed as:

$$\% \text{ Moisture} = \frac{W_2 - W_1}{W} \times 100 \dots\dots\dots [2]$$

Where:

W₂=Weight of the crucible and dry sample;
 W₁=Weight of empty crucible
 W=Weight of the sample

2.4.2 Determination of crude protein of pepper powders

The micro Kjeldahl method described by [37] was used to determine crude protein. Two grams of the sample was put into the digestion flask. A gram of copper sulphate and sodium sulphate (catalyst) in the ratio 5:1 respectively and 25 ml concentrated sulphuric acid was added to the digestion flask. The flask was placed into the digestion block in the fume cupboard and was heated until frothing ceased giving clear and light blue green colouration. The mixture was then allowed to cool and diluted with distilled water until it reaches 250ml of volumetric flask. Distillation apparatus was connected and 10ml of the mixture was poured into the receiver of the distillation apparatus, also 10ml of 40% sodium hydroxide was added. The released ammonia by boric acid was then treated with 0.2 M of hydrochloric acid until the green colour changed to purple. Percentage of nitrogen in the sample was calculated using the formula:

Crude protein = % Total Nitrogen = (Titre blank) x Normality x N₂.

Nitrogen factor = 6.25
 Crude protein= % total Nitrogen X 6.25

2.4.3 Determination of crude fat of pepper powders

The Soxhlet extraction method described by [37] was used in determining the crude fat content of the samples. Two grams of the sample was weighed and the weight of the flat bottom flask taken with the extractor. Mounted on it, the thimble was held half way into the extractor and the weight of sample. Extraction was carried out using boiling point 60°C. The thimble was plugged with cotton wool. At completion of extraction which lasted for 8 hours, the solvent was removed by evaporation on a water bath and remaining part in the flask was dried at 80°C for 30 minutes in the air oven to dry the fat then cooled in a desiccator. The flask was reweighed and percentage fat was calculated as thus:

$$\% \text{ fat} = \frac{\text{Weight of extracted fat}}{\text{Weight of sample}} \times 100 \dots \dots \dots [3]$$

2.4.4 Determination of ash content of pepper powders

The [37] method for determining ash content was used. Two (2) grams of sample was weighed into an ashing dish which had been pre-heated, cooled in a desiccator and weighed soon after reaching room temperature. The crucible and content was then heated in a muffle furnace at 550°C for 6-7 h. The dish was cooled in a desiccator and weighed soon after reaching room temperature. The total ash was calculated as percentage of the original sample weight.

$$\% \text{ Ash} = \frac{(W_3 - W_1)}{(W_2 - W_1)} \times 100 \dots \dots \dots (4)$$

Where:

W₁ = Weight of empty crucible,
 W₂ = Weight of crucible + sample before ashing,
 W₃ = Weight of crucible + content after ashing.

2.4.5 Determination of crude fiber of pepper powders

The method described by [37] was used for fibre determination. Two (2) grams of the sample was extracted using Diethyl ether. This was digested and filtered through the California Buchner system. The resulting residue was dried at 130 ± 2°C for 2 h, cooled in a desiccator and weighed. The residue was then transferred to a muffle furnace (Shanghai box type resistance furnace, No.:SX2-4-10N) and ignited at 550°C for 30 min, cooled and weighed. The percentage crude fibre content was calculated as:

$$\% \text{ Crude fibre} = \frac{\text{Loss in weight after incineration}}{\text{Weight of original food}} \times 100 \dots \dots \dots (5)$$

2.4.6 Determination of carbohydrate content of pepper powders

Carbohydrate content was determined by difference according to [41] as follows:

$$\% \text{ Carbohydrate} = 100 - (\% \text{moisture} + \% \text{Protein} + \% \text{Fat} + \% \text{Ash} + \% \text{Fibre} \dots [6])$$

2.5 Determination of Mineral Content of Pepper Powders

Mineral content of pepper was determined using the method of [37]. Five hundred milligram (500

mg) of sample was weighed in a digesting flask and 10 ml of each of HCl and HNO₃ was added. The mixture was digested for 10 minutes on a Bunsen burner and allowed to cool. The mixture was then filtered using filter paper and the filtrate was made up-to mL with distilled water and injected into the atomic absorption spectroscopy for quantification of the mineral elements except for sodium and potassium which were determined using flame photometer.

2.6 Fourier Transform Infrared Spectra Analysis of SIP and HNP Pepper Powders

The Fourier transform infrared spectra analysis of capsicum pepper samples was determine using validation of analytical methods of methamphetamine described by [42]. A total of 1 g; 0.9 g; 0.8 g; 0.7 g; 0.6 g; 0.5 g; 0.4 g; 0.3 g; 0.2 g; 0.1 g of standard methamphetamine pure powders and pepper powders (CC and CA) were put in a each cup that has been labeled. Each coupled with alum-weight to 1 g and shaken until homogeneous. It is a standard methamphetamine with a concentration of 100%, 90%, 80%, 70%, 60%, 50%, 40%,30%, 20%, and 10% were used for the preparation of calibration curve and determination of methamphetamine. The values for the precession of the system (RSD% ≤ 2.0, n = 7) show that the system is precise and method validation. The linear range of the method was studied by analyzing in duplicate ten concentrations of each compound ranging from 10.0 to 100.0%. The obtained linear ranges for each of the methamphetamine with corresponding correlation coefficients (R²) are given in paper. Limit of detection (LOD) and limit of quantification (LOQ) by an empirical method that consisted of analyzing a series of standard solutions containing decreasing amounts of methamphetamine were determined.

The limit of detection was calculated by $LOD = (3S_y/x)/a$.

Where S_y/x is the standard deviation of the response of the blank and a is the slope of the calibration curve.

The limit of quantification was calculated by $LOQ = (10S_y/x)/a$. The limits were validated by analyzing standards prepared at the concentrations of the LOQs for each standard and their precision and accuracy were assessed.

2.7 Statistical Analyses of the Samples

Data was subjected to Analysis of Variance (ANOVA) followed by T-Test and means were separated by Duncan multiple range test; significant levels were obtain at 95% ($P>0.05$). SPSS version 17 software was used.

3. RESULTS AND DISCUSSION

3.1 Proximate Composition of Pepper Powders

Table 1 shows the proximate composition of Sweet Italian (SIP) and Habanero (HNP) peppers. The moisture content of the samples were 8.89 and 8.54% for the SIP and HNP, respectively. The moisture content of SIP was significantly ($p<0.05$) higher than that of HNP pepper variety. Since the moisture content of the pepper varieties is less than 10%, it suggests that the samples may have longer shelf-life and inhibit microbial growth [43,2]. Similarly, the crude protein, crude fat and carbohydrate contents of SIP sample (9.32, 5.33 and 51.5%) were significantly ($p<0.05$) higher than that of the HNP sample (8.77, 4.87 and 50.9, respectively). In contrast, the HNP sample had significantly ($p<0.05$) higher crude fibre and ash contents (21.8 and 5.21%) than was exhibited by the SIP sample with lower values of 20.3 and 4.63%, respectively. Protein presence in peppered food product will help reduce the problem of protein-energy malnutrition for all ages in a population suffering this challenge [3]. The fat values reported in the present study were lower than those obtained for dried pepper (12.5-15.8%) harvested at different times of the year by [44]. The advantage of having low fat content in pepper is that its keeping quality will be enhanced as lipid peroxidation will minimally occur. The fibre content of the pepper varieties in this study were higher than 2.10 and 3.25% reported for garlic and ginger powders [19], suggesting that these pepper varieties had better fibre contents compared to other spices as reported by [45]. Including these pepper samples in product development could confer on such functional food products physiological and health benefits that will ameliorate morbidities and manage chronic disease conditions. The Ash results suggest that HNP and SIP are better mineral source [8]. The carbohydrate content of SIP and HNP samples obtained in this study are within recommended (45–65%) threshold levels [9].

3.2 Functional Properties of Pepper Powders

Table 2 presents the functional properties of Sweet Italian (*Capsicum annuum*) and Habanero (*Capsicum chinense*) powders. The bulk density results for the samples were 0.49 and 0.54 g/ml for SIP and HNP respectively. This showed that the pepper samples in this study are low in calories [46]. Results in this study are in line with [47] who reported a value of 0.71 for green pepper and 0.54 for red pepper for SIP and HNP respectively. The water absorption capacity for the sample were 1.36 and 1.97 g/ml for SIP and HNP respectively with HNP significantly ($p>0.05$) higher as compared to SIP. This results is in line with 1.0g/ml for Negro pepper reported by [48]. Results of oil absorption capacity for peppers in this study were 3.18 and 2.17 g/ml for SIP and HNP respectively with SIP significantly ($p>0.05$) higher than HNP. The results of the wettability showed sample SIP had higher wettability of 71.00 as compared to HNP with wettability of 57.0. The results of foaming capacity in this study were 9.10 and 6.30 for SIP and HNP respectively with SIP been significantly ($p>0.05$) higher as compared to HNP. Swelling index results revealed that SIP had higher swelling index of 2.89 as compared to HNP 2.86 but were not significantly ($p>0.05$) difference. The results of reconstitution index showed 5.87 and 6.53% for SIP and HNP respectively with SIP having higher vales than HNP. There were significant

($p>0.05$) differences observed in the functional properties of the two varieties of peppers except in the swelling index.

3.3 Mineral Content of Pepper Powders

Table 3 shows the results of the mineral content of Sweet Italian and Habanero peppers. For both the macro and trace elements content of the studied samples, the SIP sample showed significantly ($p<0.05$) higher values than HNP sample except for magnesium in which no statistical significant difference between the SIP and HNP samples was observed (8.82 & 8.52 mg/100 g). The mineral results showed that the content of the following macro-elements Na (15.3 mg/100 g), K (273 mg/100 g), Ca (38.6 mg/100 g) and P (0.05 mg/100 g) were significantly ($p<0.05$) in SIP than in HNP which recorded 12.4 mg/100 g (Na), 122 mg/100 g (K), 25.1 mg/100 g (Ca) and 0.03 mg/100 g for Phosphorus. The trace elements (Fe, Zn, Cu and Mn) measured in this study showed that SIP variety contained denser quantities than their HNP counterparts. HNP variety contained 5.76 mg/100 g of Fe, 3.21 mg/100 g of Zn, 0.31 mg/100 g of Cu and 0.83 mg/100 g of Mn. Correspondingly, SIP contained higher values of same trace elements as follows: 8.63, 4.39, 0.63 and 1.27, respectively. *The results microelement in SIP and HNP varieties has revealed the potential diversity of elements in the two samples which could impact several physiological benefits [12, 14].*

Table 1. Proximate composition of Sweet Italian and habanero peppers (dry wt basis)

Parameter	Sweet Italian pepper	Habanero pepper
Moisture content (%)	8.89 ^a ±0.85	8.54 ^b ±0.23
Crude protein content (%)	9.32 ^a ±0.93	8.77 ^b ±0.65
Crude fat content (%)	5.33 ^a ±0.84	4.87 ^b ±0.41
Crude fibre content (%)	20.3 ^b ±1.04	21.8 ^a ±0.74
Ash content (%)	4.63 ^b ±0.09	5.21 ^a ±0.34
Carbohydrate (%)	51.5 ^a ±2.0	50.9 ^b ±1.43

Values are mean±standard deviation of triplicate determinations. Mean values with same superscript letter(s) along each row are not significantly ($p>0.05$) different

Table 2. Functional properties of Capsicum varieties

Parameters	Sweet Italian Pepper	Habanero Pepper
Water Absorption Capacity(ml/g)	1.36 ^b ±0.03	1.97 ^a ±0.32
Oil Absorption Capacity(ml/g)	3.18 ^a ±0.64	2.17 ^b ±0.58
Bulk Density(g/ml)	0.49 ^b ±0.03	0.54 ^a ±0.06
Foaming Capacity	9.10 ^a ±0.81	6.30 ^b ±0.26
Swelling Index	2.89 ^a ±0.02	2.86 ^a ±0.01
Reconstitution Index	5.87 ^b ±0.12	6.53 ^a ±0.12
Wettability	71.00 ^a ±1.00	57.00 ^b ±1.00

Values are mean±standard deviation of triplicate determinations. Mean values with same superscript letter(s) along each row are not significantly ($p>0.05$) different

3.4 Fourier Transform Infrared Spectra Analysis of SIP and HNP Varieties

The results of the infrared spectrum analysis of the two pepper varieties are shown in Table 4, Fig. 2. The results showed that the pepper varieties (SIP and HNP) had similar infrared spectra that are dominated by the presence of alcohol/phenol, alkanes, alkenes and amine

groups, corresponding to the frequencies in the range 3435.44-675.11 cm^{-1} for the phenols, 1037, 1379-1458 cm^{-1} for the alkanes, 1637.62-920.08 cm^{-1} for the alkenes and 719.47-817.57 cm^{-1} for amine groups. The result indicated that the two varieties of peppers (SIP and HNP) had similar spectra lines, suggesting that they may possess similar structural, functional and bioactive properties [22,29,31].




Fig. 2(A). Infrared spectrum analysis of Sweet Italian pepper




Fig. 2(B). Infrared spectrum analysis of Habanero peppers

Table 3. Mineral composition (mg/100 g) of Sweet Italian and Habanero pepper powders

Minerals	Sweet Italian peppers	Habanero peppers
Sodium	15.3 ^a ±0.05	12.4 ^b ±0.04
Potassium	273 ^a ±0.09	122 ^b ±0.03
Calcium	38.6 ^a ±0.02	25.1 ^b ±0.05
Magnesium	8.82 ^a ±0.07	8.52 ^a ±0.08
Phosphorus	0.05 ^a ±0.07	0.03 ^b ±0.01
Iron	8.63 ^a ±0.05	5.76 ^b ±0.11
Zinc	4.39 ^a ±0.07	3.21 ^b ±0.04
Copper	0.63 ^a ±0.04	0.31 ^b ±0.07
Manganese	1.27 ^a ±0.08	0.83 ^b ±0.03

Values are mean±standard deviation of triplicate determinations. Mean values with same superscript letter(s) along each row are not significantly ($p>0.05$) different

Table 4. Infrared spectrum analysis of SIP and HNP pepper powders

Frequencies	Functional group	Assignment	Assignment
3435.44	Alcohols and Phenols	O-H (H-bonded)	O-H (H-bonded)
2928.04	Aliphatic/Methylene	C-H (asymmetry)	C-H (asymmetry)
2854.74	Aliphatic/methylene	C-H (symmetry stretch)	C-H (symmetry stretch)
1637.62	Aliphatic/Alkenyl	C=C (Stretch)	C=C (Stretch)
1458.23	Aliphatic/methyl	C-H (asymmetry)	C-H (asymmetry)
1379.25	Aliphatic/methyl	C-H (symmetry bend)	C-H (symmetry bend)
1319.35	Aliphatic/Methyne	C-H (bend)	C-H (bend)
1242.20	Aliphatic/vibrations	C-C (vibrations)	C-C (vibrations)
1157.33	Aliphatic/vibrations	C-C (vibrations)	C-C (vibrations)
1101.39	Aliphatic	Cyclohexane ring vibrations	Cyclohexane ring vibrations
1062.81	Alcohol	C-O (Broad)	C-O (Broad)
1037.64	Aliphatic	Cyclohexane ring vibrations	Cyclohexane ring vibrations
920.08	Olefinic (Alkene)	Vinyl CH out-of-plane bend	Vinyl CH out-of-plane bend
817.57	Amines	NH ₂ and N-H wagging (shifts on H-bonding)	NH ₂ and N-H wagging (shifts on H-bonding)
779.27	Amines	NH ₂ and N-H wagging (shifts on H-bonding)	NH ₂ and N-H wagging (shifts on H-bonding)
719.47	Amines	NH ₂ and N-H wagging (shifts on H-bonding)	NH ₂ and N-H wagging (shifts on H-bonding)
675.11	Alcohols and phenols	O-H bend (out-of-plane)	O-H bend (out-of-plane)

4. CONCLUSION

Including these pepper samples in product development could confer on such functional food products physiological and health benefits that will ameliorate morbidities and manage chronic disease conditions. The SIP and HNP pepper samples could potentially contribute to energy provision in diet when used in product development or added to foods as spice. The two varieties of peppers, SIP and HNP having similar spectra lines may also possess similar structural properties. Presence of amine group in the samples indicated that they may have some quantities of Nitrogen. However, the samples also possessed broad C-O groups, indicating that the samples are richer in phenols/alcohol groups compared to other groups.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Marques ÍE, Lucion FB, Bizzi CA, Cichoski AJ, Wagner R, de Menezes CR. Are infrared and microwave drying suitable alternatives for moisture determination of meat products? Journal of Food Quality. 2016;39(4):391-397.
2. Zambrano MV, Dutta B, Mercer DG, MacLean HL, Touchie M. Assessment of moisture content measurement methods of dried food products in small-scale operations in developing countries: A

- review. Trends in Food Science & Technology. 2019;88:484–49.
3. Ubesie A, Ibeziakor N. High burden of protein–energy malnutrition in Nigeria: Beyond the health care setting. Annals of Medical and Health Sciences Research. 2012;2(1):66-69.
 4. Okwu D, Nnamdi FU. Evaluation of the chemical composition of *Dacryodes edulis* and *Raphia hookeri* Mann and Wendl exudates used in herbal medicine in south eastern Nigeria. African Journal of Traditional, Complementary and Alternative Medicines. 2008;5(2):194-200.
 5. Hervik AK, Svihus B. The role of fiber in energy balance. Journal of Nutrition and Metabolism. 2019;498365.
 6. Dhingra D, Michael M, Rajput H, Patil R. Dietary fibre in foods: A review. Journal of Food Science and Technology. 2012; 49(3):255-266.
 7. Oloyede O. Chemical profile of unripe pulp of *Carica papaya*. Pakistan journal of nutrition. 2005;4(6):379-381.
 8. Okwu DE, Okoro E. Phytochemical composition of *Brachystegia eurycoma* and *Mucuna flagellipes* seeds. Medicinal and Aromatic Plant Science and Biotechnology. 2006;26:1-4.
 9. Manore MM. Exercise and the Institute of Medicine recommendations for nutrition. Current Sports Medicine Reports. 2005; 4(4):193-198.
 10. Huang F, Yu P, Yuan Y, Li Q, Lin F, Gao Z. The relationship between sodium excretion and blood pressure, urine albumin, central retinal arteriolar equivalent. BMC Cardiovascular Disorders. 2016;16(1):194.
 11. Forrest MD. The sodium-potassium pump is an information processing element in brain computation. Frontiers in Physiology. 2014;5:472.
 12. Esayas K, Shimelis A, Ashebir F, Negussie R, Tilahun B, Gulelat D. Proximate composition, mineral content and antinutritional factors of some capsicum (*Capsicum annum*) varieties grown in Ethiopia. Bulletin of the Chemical Society of Ethiopia. 2011;25(3):53-59.
 13. Pravina P, Sayaji D, Avinash M. Calcium and its role in human body. International Journal of Research in Pharmaceutical and Biomedical Sciences. 2013;4(2):659-668.
 14. Del Valle HB, Yaktine AL, Taylor CL, Ross AC. Dietary reference intakes for calcium and vitamin D: 2011; National Academies Press.
 15. Jannen-Dechent W, Ketteler M. Magnesium basics. Clinical Kidney Journal. 2012;5(1):13-14.
 16. Abbaspour N, Hurrell R, Kelishadi R. Review on iron and its importance for human health. Journal of research in medical sciences: the official Journal of Isfahan University of Medical Sciences. 2014;19(2):164.
 17. Olapade A, Akinyanju F. Chemical and functional properties and performance of blends of water yam (*Dioscorea Alata*) and Soybean (*Glycine Max*) flours for water yam ball (Ojojo) preparation. American Journal of Chemistry. 2014;4(3):89-96.
 18. Gozzelino R, Arosio P. The importance of iron in pathophysiologic conditions. Frontiers in pharmacology. 2015;6:26.
 19. Otunola GA, Oloyede OB, Oladiji AT, Afolayan AJ. Comparative analysis of the chemical composition of three spices–*Allium sativum* L. *Zingiber officinale* Rosc. and *Capsicum frutescens* L. commonly consumed in Nigeria. African Journal of Biotechnology. 2010;9(41):6927-6931.
 20. El Kaaby EA, Al Hattab ZN, Al-Anny JA. FT-IR Identification of Capsaicin from callus and seedling of chilli pepper plants *Capsicum annum* L. *in vitro*. Int. J. of Multidisciplinary and Current research. 2016; 4.
 21. Ashokkumar R, Ramaswamy M. Phytochemical screening by FTIR spectroscopic analysis of leaf extracts of selected Indian medicinal plants. International journal of Current Microbiology and applied Sciences. 2014; 3(1):395-406.
 22. Al-Shareefi E, Sahib AHA, Hameed IH. Phytochemical Screening by FTIR Spectroscopic Analysis and Anti-Fungal Activity of Fruit Extract of Selected Medicinal Plant of *Ruta graveolens*. Indian Journal of Public Health Research & Development. 2019;10(1):994-999.
 23. Sankari G, Krishnamoorthy E, Jayakumar S, Gunasekaran S, Priya VV, Subramaniam S. Analysis of serum immunoglobulins using Fourier transform infrared spectral measurements. Biology and Medicine. 2010;2(3):42-48.
 24. Baker MJ, Trevisan J, Bassan P, Bhargava R, Butler HJ, Dorling KM. Using Fourier transform IR spectroscopy to analyze

- biological materials. Nature protocols. 2014;9(8):1771.
25. Rabe JH, Sammour DA, Schulz S, Munteanu B, Ott M, Ochs K. Fourier transform infrared microscopy enables guidance of automated mass spectrometry imaging to predefined tissue morphologies. Scientific Reports. 2018; 8(1):1-11.
 26. Maitra I, Morais CL, Lima KM, Ashton KM, Date RS, Martin FL. Attenuated total reflection Fourier-transform infrared spectral discrimination in human bodily fluids of oesophageal transformation to adenocarcinoma. Analyst. 2019;144(24): 7447-7456.
 27. Rodrigues RP, Aguiar EM, Cardoso-Sousa L, Caixeta DC, Guedes CC, Siqueira WL. Differential Molecular Signature of Human Saliva Using ATR-FTIR Spectroscopy for Chronic Kidney Disease Diagnosis. Brazilian Dental Journal. 2019;30(5):437-445.
 28. Ferreira IC, Aguiar EM, Silva AT, Santos LL, Cardoso-Sousa L, Araújo TG. Attenuated Total Reflection-Fourier Transform Infrared (ATR-FTIR) Spectroscopy Analysis of Saliva for Breast Cancer Diagnosis. Journal of oncology; 2020.
 29. Zhang J, Li B, Wang Q, Wei X, Feng W, Chen Y. Application of Fourier transform infrared spectroscopy with chemometrics on postmortem interval estimation based on pericardial fluids. Scientific Reports. 2017;7(1):1-8.
 30. Takamura A, Watanabe K, Akutsu T, Ozawa T. Soft and robust identification of body fluid using Fourier transform infrared spectroscopy and chemometric strategies for forensic analysis. Scientific Reports. 2018;8(1):1-10.
 31. Zha S, Wei X, Fang R, Wang Q, Lin H, Zhang K. Estimation of the age of human semen stains by attenuated total reflection Fourier transform infrared spectroscopy: A preliminary study. Forensic Sciences Research. 2019; 1-7.
 32. Amir RM, Anjum FM, Khan MI, Khan MR, Pasha I, Nadeem M. Application of Fourier transform infrared (FTIR) spectroscopy for the identification of wheat varieties. Journal of Food Science and Technology. 2013; 50(5):1018-1023.
 33. Kowalczyk D, Pitucha M. Application of FTIR Method for the Assessment of Immobilization of Active Substances in the Matrix of Biomedical Materials. Materials. 2019;12(18):2972.
 34. Raja PB, Sethuraman MG. Inhibitive effect of black pepper extract on the sulphuric acid corrosion of mild steel. Materials Letters. 2008;62(17-18):2977-2979.
 35. Ahmed J, Shivhare U. Thermal kinetics of color change, rheology, and storage characteristics of garlic puree/paste. Journal of Food Science. 2001;66(5):754-757.
 36. Vengaiah P, Pandey J. Dehydration kinetics of sweet pepper (*Capsicum annum* L.). Journal of Food Engineering. 2007;81(2):282-286.
 37. AOAC. Official Methods of Analysis of AOAC International. 19th edition. AOAC International, Gaithersburg, Maryland, USA; 2012.
 38. Onwuka GI. Food analysis and instrumentation: theory and practice: Naphthali Prints; 2005.
 39. Narayana K, Narasinga Rao M. Functional properties of raw and heat processed winged bean (*Psophocarpus tetragonolobus*) flour. Journal of Food Science. 1982;47(5):1534-1538.
 40. AOAC. Association of Official Analytical Chemists. Official methods of analysis (16th ed.). Washington, DC: Author.
 41. Ihekoronye AI, Ngoddy PO. Integrated food science and technology for the tropics: Macmillan; 1985.
 42. UNODC. United Nations Office on Drugs and Crime. Recommended Methods for the Identification and Analysis of Amphetamine, Methamphetamine and Their Ring-Substitutes Analogues in Seized Materials. United Nations, New York; 2006.
 43. Nadeem M, Rehman S, Anjum F, Bhatti I. Textural profile analysis and phenolic content of some date palm varieties. J. Agric. Res. 2011;49(4):525-539.
 44. Bhandari SR, Bashyal U, Lee YS. Variations in proximate nutrients, phytochemicals, and antioxidant activity of field-cultivated red pepper fruits at different harvest times. Horticulture, Environment, and Biotechnology. 2016;57(5):493-503.
 45. Otunola GA, Afolayan AJ, Ajayi EO, Odeyemi SW. Characterization, antibacterial and antioxidant properties of silver nanoparticles synthesized from aqueous extracts of *Allium sativum*, *Zingiber officinale*, and *Capsicum*

- frutescens*. Pharmacognosy Magazine. 2017;13(2):201.
46. Nuru M, Muradashvili N, Kalani A, Lominadze D, Tyagi N. High methionine, low folate and low vitamin B6/B12 (HM-LF-LV) diet causes neurodegeneration and subsequent short-term memory loss. Metabolic brain disease. 2018;33(6):1923-1934.
47. Dawit L. Product Development and Evaluation of Fermented hot Pepper (*Capsicum annuum* L.), Datta. Addis Ababa University; 2018.
48. Oso BJ, Oladiji AT. Total Phenolic Contents and Antioxidant Variations in Raw and Cooked Dried Fruit of *Xylopia aethiopica*. International Annals of Science. 2019;6(1):13-17.

© 2023 Igbakaa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/100240>