

# Esophageal Papilloma: Case Series of 4 Patients

Seydou Roubanatou <sup>a\*</sup>, Ghenim Houda <sup>a</sup>, Dauvois Barbara <sup>a</sup>  
and Lagasse Jean Paul <sup>a</sup>

<sup>a</sup> Hepato-Gastro-Enterology Department, UHC ORLEANS, Orleans, France.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108237>

**Case Study**

**Received: 10/09/2023**  
**Accepted: 14/11/2023**  
**Published: 27/11/2023**

## **ABSTRACT**

**Background:** Papillomas are rare benign lesions that can be found in the oesophagus. Being rarely seen like a single small whitish, wart-like lesion grown from the mucosa. Histopathological examination is the gold standard for a definitive diagnosis. Excision of papillomas is recommended for definitive treatment in the fact of the potential for malignant transformation.

**Objectives:** Through these differences cases, we would like to show the clinical and endoscopic difference that papillomas in the oesophagus can present.

**Methods:** We report clinicopathologic features of four esophageal papilloma cases detected in the endoscopy unit of our department. These patients had undergone oesophageal duodenal fibroscopy during 2022. The inclusion criteria were endoscopic appearance and histology.

**Results:** We present four cases of this rare endoscopic finding. While three of our patients were asymptomatic, one had an acute epigastric pain, all the lesions were excised.

**Conclusion:** Endoscopists should be able to make the differential diagnosis of papilloma detected in the esophagus and have knowledge about these lesions as they may carry malignant potential.

*Keywords: Oesophageal papilloma; histology; upper endoscopy; endoscopic resection.*

\*Corresponding author: Email: [seydouroubanatou20@gmail.com](mailto:seydouroubanatou20@gmail.com);

## 1. INTRODUCTION

Papilloma of the oesophagus was first described by Adler et al. in 1959 [1]. It is a rare benign tumour of the oesophagus, mostly discovered incidentally at endoscopy, but can also be symptomatic. Its natural history remains poorly understood, and the risk of malignancy exists [2]. There is no consensus on treatment or monitoring methods. We report four cases in our department.

## 2. CASE PRESENTATION

### Case 1:

A 54 year old male patient known to be diabetic, on insulin, with a history of decompensated viral B and delta cirrhosis (MELD score of 17 and CHILD- PUGH C 11) admitted to hospital for acute epigastric pain which did not respond to symptomatic treatment. The upper endoscopy revealed a duodenal ulcer with grade II oesophageal varices and hypertensive gastropathy in addition to two small lesions of the middle third of the oesophagus, one was pedunculated which was totally resected by cold forceps. The histological analysis showed papilloma.

### Case 2:

A 66 year old male patient has been treated for prostatic adenocarcinoma, operated for subependymoma of the 4th ventricle complicated post operatively by pharyngo-laryngeal paralysis leading to a tracheotomy, and subsequently swallowing problems needing a definitive gastrostomy. During endoscopic follow-up 10 years after placement of the gastrostomy tube, several centimetric lesions in the middle oesophagus were found. The histo-pathology confirmed the papillomatous nature of the lesions. He subsequently underwent several sessions of argon plasma electrocoagulation of these lesions, which resulted in their almost complete disappearance.

### Case 3:

A 65-year-old female smoker patient known to be hypertensive with a family history of adenomatous polyps underwent an upper endoscopy which revealed an 8 mm pedunculated papilloma in the upper third of the oesophagus. It was completely resected by cold forceps.

### Case 4:

A 64 year old male patient, treated for alcoholic compensated cirrhosis. An upper endoscopy was performed searching for esophageal varices. Surprisingly, 3 small lesions measuring 2-3 mm in the lower third of the oesophagus were found. A biopsy was performed to confirm the diagnosis by cold forceps.

## 3. DISCUSSION

Oesophageal papilloma is a rare epithelial tumour with a prevalence in patients undergoing upper GI endoscopy of 0.01%-0.45%. In the general population, their prevalence is estimated at 0.006%-0.04% based on autopsy series [3-6]. The age of onset of papilloma is often around 50 years [7]; in our series, the average age at diagnosis of papilloma was 59 years. Several publications, such as TSAI et al. [7] and Szanto et al. [4], show that oesophageal papillomas are more frequent in women, whereas other publications, such as Takeshita et al. [8] and Huart et al. [2], report a higher number in men, as the case in our series.

There is not much known about the pathophysiology. Two main causes have been discussed and appear to be synergistic [2]: chronic irritation of the oesophageal mucosa due to chemical or mechanical factors, and HPV infection. The mechanism of chronic inflammation can be found in gastro-oesophageal reflux, Barrett's esophagus, peptic oesophagitis, and mechanical esophageal trauma such as variceal sclerosis or self-expanding metal prostheses [2]. This coincides with our case reports, most of which present these favourable factors described above. Human papillomavirus (HPV) has also been implicated in the formation of oesophageal papilloma. HPVs are known to cause benign lesions, but can lead to the development of cancer. Infection with human papilloma virus (HPV) has been suggested as a risk factor for oesophageal carcinoma. Some authors suggest a role for HPV in oesophageal carcinogenesis in areas of high incidence of oesophageal squamous cell carcinoma, a role not found in areas of low incidence. The HPV subtypes involved appear to be high-risk HPVs such as HPV-16 and 18 [2]. Some studies, shown that the esophageal papilloma, has no relation with HPV. Its role is still, very much debated [3]. The prevalence of HPV detection in oesophageal papilloma varies between 0% and 87.5% in the

various series published [8]. Takeshita et al found a prevalence of 10.5% of HPV positive in his series, while Bohn et al reported 85.7% with a positive HPV PCR [9]. However, in other series HPV was not identified, which may be due to the involvement of as yet unknown subtypes of HPV or the sensitivity of microbiological tests [2]. HPV was not identified in the histological specimens of our patients and we did not request for PCR.

The endoscopic appearance is characteristic but not pathognomonic. It may be confused with glycogenic acanthosis or verrucous carcinoma, a particular type of squamous cell carcinoma. It may present as a sessile or pedunculated polypoid formation, demarcated from adjacent tissue [2].

Most commonly, the size of the papilloma varies between 3 and 12 mm, with an average size of less than 5 mm [10]. Papillomas may be single or multiple, sometimes forming large papillomatoses, and giant forms up to 5 cm have been described [2]. In our series, we also found lesions that were millimetre-sized in the majority of cases, sometimes single or multiple.

Previous studies report variable data regarding the location of oesophageal papilloma. For Bohn et al, the lesions were mainly located in the upper oesophagus. However, other studies suggest that the middle and distal oesophagus may also be involved. This could be explained by damage to the mucosa caused by acid reflux [10].

The diagnosis of papilloma is confirmed histologically. It is a benign tumour in which the general architecture of the squamous epithelium is preserved and the basement membrane is intact [2]. In our series, histology was crucial in confirming the diagnosis. Treatment is not yet fully codified, but is essentially endoscopic. Some carry out endoscopic resection using cold forceps [11], a method that has been validated for the majority of our patients, or using hot snare polypectomy or mucosectomy [3,12].

No surveillance has been established. However, some authors suggest that the risk of malignant transformation is very low in case of solitary papilloma, but higher in the case of oesophageal papillomatosis or giant papilloma [2].


#### 4. CONCLUSION

Papilloma of the oesophagus is a rare benign tumour, for which certain factors play a role in its

appearance. It has the potential for malignant transformation and the risk increases with the number and size of lesions, hence the importance of identifying them during endoscopy and treating them using the most conservative method possible, often endoscopic resection.



**Fig. 1. Upper endoscopy showing an esophageal polyp**



**Fig. 2. Several large esophageal lesions**

#### CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

#### ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Adler RH, Carberry DM, Ross CA. Papilloma of the esophagus. Association with hiatus hernia. J Thorac Surg. 1959;37:625–35.
2. D'Huart MC, Kull E, Peyrin-Biroulet L, et al. Endoscopic management of oesophageal papillomas. Acta Endosc. 2014;44:43-50.
3. Moasca S, Manes G, Monaco R, Bellomo PF, Botttini V, Balzano. A squamous papilloma of the oesophagus : long-term follow up. J Gastroenterol Hepatol. 2001;16:857-61.
4. Szanto I, Szentirmay Z, Banai J, et al. Squamous papilloma of the oesophagus. Clinical and pathological observations based on 172 patients.Orv Hetil. 2005;146:547-552
5. Plachta A. Benign tumors of the esophagus. Review of literature and report of 99 cases. Am J Gastroenterol. 1962;38:639-652.
6. Sablich R, Benedetti G, Bignucolo S, Serraino D. Squamous cell papilloma of the esophagus. Report on 35 endoscopic cases. Endoscopy. 1988;20:5-7.
7. Tsai SJ, Lin CC, Chang CW, et al. Benign oesophageal lesions: Endoscopic and pathologic features. World J Gastroenterol Hepatol. 2015;21:1091-1098.
8. Takeshita K, Murata S, Mitsufuji S, et al. Clinicopathological characteristics oesophageal squamous papillomas in japanese patients-with comparaisons of findings from western countries. Acta Histochem. 2006 ;10:228-230
9. Bohn OL, Navarro L, Saldivar J, Sanchez-Sosa S. Identification of human papillomavirus in oesophageal squamous papillomas. World J Gastroenterol. 2008; 14:7107-11.
10. Ekin N, Bestas R, Cetin A. Clinicopathological characteristics of patients with oesophageal squamous papilloma in Turkey and comparaisn with the littérature data: The largest case series ever reported from Turkey. Int J C clin Pract. 2021;75:e14420
11. Mavilla MG, Wu GY. Esophageal squamous papilloma: A case series and littérature review. J Dig Dis. 2018;19 (4):254-6
12. Sablich R, Benedetti G, Bignucolo S, Serraino D. Sqamous cell papilloma of the oesophagus: report of 35 endoscopic.Case endoscopy. 1988;20:5-7.

© 2023 Roubanatou et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/108237>