



# Small Bowel Obstruction Due to Appendix Adhesion to an Aortic Graft

Oluwatobi Onafowokan <sup>a++</sup> and Hugo J.R. Bonatti <sup>a+++\*</sup>

<sup>a</sup> Meritus Health, 11110 Medical Campus Road, Suite 147, Hagerstown, MD 21742, US State, USA.

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113960>

**Case Study**

**Received: 05/01/2024**

**Accepted: 09/03/2024**

**Published: 18/03/2024**

## **ABSTRACT**

**Background:** Small bowel obstruction (SBO) is a common cause of surgical consultation. Adhesion bands, internal hernias, tumors and many other causes need to be considered. If non-operative management fails, exploratory laparoscopy or laparotomy is indicated.

**Case Report:** A 63 year old male with multiple comorbidities had undergone open aortobifemoral bypass grafting for occlusive disease. The graft failed as the patient was non-compliant with medical management and he underwent right below knee amputation. During the following 3 months, he had multiple emergency room visits for recurrent abdominal pain and vomiting. He ultimately was admitted with complete SBO and CT-scan showed a transition point in the mid-ileum. A mass in the tail of the pancreas was also found. He initially responded to non-operative management, however, upon oral challenge SBO relapsed and decision for surgery was made. On laparoscopy, the transition point was identified and strangulation was found to be caused by the appendix, which crossed the midline. The appendix base at the cecum was identified but the tip could not be found. Therefore, the appendix base was stapled off and the vascular pedicle was secured. The small bowel could be freed and the appendix was followed and cut out of dense adhesions to the aortic graft; pathology indicated chronic appendicitis. The patient recovered without complications from this surgery.

<sup>++</sup> Meritus Surgical Specialists;

<sup>\*</sup>Corresponding author: E-mail: [hugo.bonatti@dr.com](mailto:hugo.bonatti@dr.com);

**Conclusions:** This is a very unusual case of SBO caused by an appendix, which became adhered to an aortic graft. It remains unclear if this pathology was caused by post-operative inflammation or if the patient had appendicitis. Laparoscopic exploration is useful in the management of SBO that does not resolve with non-operative management.

*Keywords: Appendicitis; aortobifemoral bypass; small bowel obstruction; laparoscopy.*

## 1. INTRODUCTION

Small bowel obstruction (SBO) is one of the most common disorders requiring surgical consultation [1]. SBO may be caused by a variety of conditions including adhesions after surgery, obstruction due to inflammatory, and infectious or neoplastic disorders amongst many others [2]. Symptoms of obstruction can also be due to a paralytic ileus which again may be caused by a variety of conditions [2]. Non operative management is initiated in most patients using decompression with a nasogastric tube with subsequent work up for the underlying pathology. If patient do not respond to nonoperative management surgery is indicated with laparoscopy becoming the preferred approach by many surgeons [3-5].

Traditionally, aortic aneurysm and aortoiliac occlusive disease has been treated by open replacement of the distal aorta but endovascular treatment has become an excellent alternative [6,7]. Multiple complications may occur following open surgery including intraabdominal adhesions that may lead to SBO requiring surgical intervention particularly when using the transperitoneal approach [8,9]. SBO may be associated with acute and chronic appendicitis [10]. In the case of acute appendicitis paralytic ileus due to the inflammation or local obstruction by the inflammatory process may be encountered and in the case of chronic appendicitis the appendix may get stuck to a bowel loop or to the abdominal wall thus creating a pocket with possible herniation of intestine [11].

We present a case of small bowel obstruction due to adhesion of the appendix to an aortobifemoral bypass graft.

## 2. CASE PRESENTATION

A 63 year old male with multiple comorbidities including coronary artery disease, hypertension


and chronic obstructive pulmonary disease had previously undergone open aortobifemoral bypass grafting for aortoiliac occlusive disease. The graft failed as the patient was non-compliant with medical management, and he subsequently underwent right below-knee amputation. During the following three months, he had multiple visits to the emergency room and multiple admissions for recurrent abdominal pain and vomiting. He ultimately was re-admitted and CT scan confirming small bowel obstruction with a transition point in the mid-ileum (Fig. 1). A mass in the pancreas tail was also incidentally found. He initially responded to non-operative management with bowel rest and decompression using a nasogastric tube (NGT); his pain improved, NGT output declined, and he passed gas. However, upon oral challenge he again worsened, abdominal x-ray indicated ongoing small bowel obstruction (Fig. 2), and the decision was made to perform an exploratory laparoscopy.

Trocars were placed in the left upper and lower quadrants, and suprapubically in his previous midline incision. Omental adhesions were taken down and on exploration, the transition point as previously seen on the CT image was found when running the small bowel. The intestinal strangulation was caused by the appendix, which crossed the midline. The appendix base at the cecum was visualized but the tip could not be mobilized as it was stuck in the left abdomen. Therefore, the appendix base and the vascular pedicle were stapled off. The small bowel could be freed and recovered during the procedure and did not require resection. The appendix was followed and was found to be densely adhered to the anterior surface of the aortic graft. The appendix tip was cut out of the adhesions and removed in a retrieval bag, with subsequent pathology indicating chronic appendicitis.

The patient recovered without complications from this surgery but died within 6 months from metastatic pancreatic cancer.



**Fig. 1. CT-scan with transition point where appendix crossed the small bowel loop (arrow)**



**Fig. 2. Abdominal X-ray: Ongoing small bowel obstruction with dilated loops up to >4cm, some oral contrast had passed into the ascending colon**

### 3. DISCUSSION

Surgical approach to aortobifemoral grafting may be transperitoneal or retroperitoneal using a rather large incision but recently also

laparoscopic techniques have been reported [12]. Neither approach has been shown to be definitively superior to the other [13], however, some studies have indicated use of the retroperitoneal approach resulting in decreased

blood loss and shorter hospital and ICU stay, and possible lower risk for ileus when compared to the transperitoneal approach [14]. The transperitoneal approach more commonly results in gastrointestinal complications, due to the peritoneal defect creating significant potential for omental and/or intraperitoneal organ interaction with the graft. Aortobifemoral bypass has remained a procedure with a high risk for a multitude of complications [12,15] including bowel obstruction [16].

Only few appendix-related complications associated with aortic grafts have been published. Graft complications involving acutely inflamed appendices are very rare, making up 0.6-5% of all secondary aorto-enteric prosthetic fistulas. The few known cases demonstrated a fistula between the graft and the appendix, which may be acutely or chronically inflamed and being in a retro-ileal position allowing the appendix to interact with the aortic prosthesis [17,18]. Such a fistula may occur even with retroperitoneal closure. The pathology is more likely due to mechanical factors such as graft dehiscence and/or anastomotic aneurysm than due to paraprostatic infection following acute appendicitis [19]. The majority of patients with aorto-appendiceal fistula present with hematochezia [18]. Management includes replacement of the graft with simultaneous appendectomy.

Our patient presented with a SBO and only during laparoscopy the unusual pathology could be identified and treatment consisted of a simple appendectomy. As the graft did not show any signs of infection the aortobifemoral graft was left in situ, with the patient being carefully monitored post-operatively for any signs of infection. Final pathology indicated inflammation of the appendix but it cannot be determined if chronic appendicitis was the cause of the condition or if the appendix tip developed adhesions to the graft with a subsequent inflammatory reaction.

As the SBO did not resolve within few days with bowel rest and decompression using a nasogastric tube, the patient underwent laparoscopic exploration revealing this unusual case. The appendectomy released the strangulated small bowel segment and the patient made a quick recovery.

#### 4. CONCLUSION

Exploratory laparoscopy should be considered in the management of small bowel obstruction that

does not resolve with non-operative management [20]. Rare causes of SBO as in our case should always be considered.

#### CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

#### ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

#### DISCLAIMER

Parts of the data were presented as a poster at the Austrian Surgical Association Meeting in Vienna, Austria.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Kuehn F, Weinrich M, Ehmann S, Kloker K, Pergolini I, Klar E. Defining the need for surgery in small-bowel obstruction. *J Gastrointest Surg.* 2017 Jul;21(7):1136-41.
2. Miller G, Boman J, Shrier I, Gordon PH. Etiology of small bowel obstruction. *Am J Surg.* 2000 Jul;180(1):33-6.
3. Morelli M, Strambi S, Cremonini C, Musetti S, Tonerini M, Coccolini F, et al. Adhesive small bowel obstruction: Predictive factors of laparoscopic failure. *Updates Surg.* 2023 Dec 27.
4. Yamamoto Y, Kitazawa M, Otsubo T, Miyagawa Y, Tokumaru S, Nakamura S, et al. Comparison of clinical outcomes and safety between open and laparoscopic surgery for adhesive small bowel obstruction: A propensity-matched analysis of a national inpatient database. *J Laparoendosc Adv Surg Tech A.* 2022 Oct;32(10):1064-70.
5. Fu WJ, Xiao X, Gao YH, Hu S, Yang Q. Analysis of risk factors for recurrence and prognosis of adhesive small bowel obstruction. *Asian J Surg.* 2023 Sep;46(9):3491-5.
6. Paravastu SC, Jayarajasingam R, Cottam R, Palfreyman SJ, Michaels JA, Thomas

- SM. Endovascular repair of abdominal aortic aneurysm. *Cochrane Database Syst Rev.* 2014 Jan 23(1):CD004178.
7. Dua A, Kuy S, Lee CJ, Upchurch GR, Jr., Desai SS. Epidemiology of aortic aneurysm repair in the United States from 2000 to 2010. *J Vasc Surg.* 2014 Jun;59(6):1512-7.
  8. Mann AJ, Laconi N, Smith RS. Closed-loop bowel obstruction years after an open abdominal aortic aneurysm repair. *Cureus.* 2021 Oct;13(10):e18586.
  9. Glover K, Lyden S, Bena JF, Smolock C, Parodi F. Mid-Term Outcomes of Retroperitoneal and Transperitoneal Exposures in Open Aortic Aneurysm Repair. *Ann Vasc Surg.* 2020 Jul;66:35-43 e1.
  10. Deshmukh SN, Maske AN, Bote SM, Parashi HS. Small bowel obstruction caused by appendiceal tourniquet. *Am J Surg.* 2011 Feb;201(2):e21-2.
  11. Maly O, Paral J. Appendicitis as a rare cause of mechanical small-bowel obstruction: A literature review of case reports. *Int J Surg Case Rep.* 2016;29:180-4.
  12. Bruls S, Quaniers J, Tromme P, Lavigne JP, Van Damme H, Defraigne JO. Comparison of laparoscopic and open aortobifemoral bypass in the treatment of aortoiliac disease. Results of a contemporary series (2003-2009). *Acta Chir Belg.* 2012 Jan;112(1):51-8.
  13. Ma B, Wang YN, Chen KY, Zhang Y, Pan H, Yang K. Transperitoneal versus retroperitoneal approach for elective open abdominal aortic aneurysm repair. *Cochrane Database Syst Rev.* 2016 Feb 5;2:CD010373.
  14. Arya N, Sharif MA, Lau LL, Lee B, Hannon RJ, Young IS, et al. Retroperitoneal Repair of Abdominal Aortic Aneurysm Reduces Bowel Dysfunction. *Vascular and Endovascular Surgery.* 2009;43(3):262-70.
  15. Bredahl K, Jensen LP, Schroeder TV, Sillesen H, Nielsen H, Eiberg JP. Mortality and complications after aortic bifurcated bypass procedures for chronic aortoiliac occlusive disease. *Journal of Vascular Surgery.* 2015;62(1):75-82.
  16. Citgez B, Yetkin G, Uludag M, Akgun I, Ekici U, Kartal A. A rare complication of aortobifemoral bypass operation: internal herniation. *Ulus Travma Acil Cerrahi Derg.* 2013 Mar;19(2):164-6.
  17. Van der Vliet QM, Berger P, De Borst GJ, Moll FL. Graft-appendiceal fistula treated with a rifampicin-silver in-situ graft. *Surg Infect (Larchmt).* 2014 Aug;15(4):450-3.
  18. Tse DM, Thompson AR, Perkins J, Bratby MJ, Anthony S, Uberoi R. Endovascular repair of a secondary aorto-appendiceal fistula. *Cardiovascular and Interventional Radiology.* 2011 Oct;34(5):1090-3.
  19. Lauwers P, De Greef K, Van den Brande F, Hendriks J, De Maeseneer M, Van Schil P. Aortic graft infection from appendicitis. A case report. *Acta Chir Belg.* 2004 Aug;104(4):454-6.
  20. Behman R, Nathens AB, Karanicolas PJ. Laparoscopic Surgery for Small Bowel Obstruction: Is It Safe? *Adv Surg.* 2018 Sep;52(1):15-27.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/113960>