



# Molecular Characterization of Extended-spectrum $\beta$ -Lactamase Producing *Escherichia coli* from Suspected Cases of Diarrhea in Nasarawa South Senatorial Zone, Nigeria

Zakou A.T <sup>a,b\*</sup>, Nkene I.H <sup>a</sup>, Bolarinwa O.F <sup>a</sup>, Tama S.C <sup>a</sup>, Ngwai Y.B <sup>a</sup>, Ishaleku D <sup>a</sup> and Abimiku R.H <sup>c</sup>

<sup>a</sup> Department of Microbiology, Nasarawa State University, P.M.B. 1022, Keffi, Nigeria.

<sup>b</sup> Dalhatu Araf Specialist Hospital, Lafia, Nasarawa State, Nigeria.

<sup>c</sup> Institute of Human Virology, Nigeria, Abuja, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. Author NYB designed the study, while Author ZAT performed the statistical analysis, and wrote the protocol. Author ID wrote the first draft of the manuscript and Authors NIH and ARH managed the analyses of the study. Authors TSC and BOF managed the literature searches. All authors read and approved the final manuscript.

## Article Information

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123624>

**Original Research Article**

**Received: 11/07/2024**

**Accepted: 13/09/2024**

**Published: 17/09/2024**

\*Corresponding author: E-mail: zakouahmadu2013@gmail.com;

## ABSTRACT

**Aims:** To investigate the prevalence and molecular diversity of ESBL producing *E. coli* isolates from children with suspected cases of diarrhea, in Nasarawa South, senatorial zone, Nigeria

**Study Design:** Investigative study.

**Place and Duration of Study:** Nasarawa South Senatorial District, Nigeria, in 2023.

**Methodology:** A total of eight (8) phenotypic ESBL producing isolates that form suspected cases of diarrhea in Nasarawa south, Nigeria, from previous study were used for polymerase chain reaction detection of ESBL genes and molecular diversity of ESBL isolates were determined using restriction fragment length polymorphism.

**Results:** The prevalence of *bla*<sub>CTX-M</sub> gene (4, 50%) was found to more prevalent among ESBL producing isolates than *bla*<sub>SHV</sub> and *bla*<sub>TEM</sub> (3, 37.5%). The strains of ESBL isolates that form the suspected cases of diarrhea among children in Nasarawa South were A and B and the proportion was 50%. The *bla*<sub>CTX-M</sub> gene was frequently detected in the ESBL producers and strain A and B were the only strain of ESBL producing isolates that form suspected cases of diarrhoea in the study location.

**Conclusion:** *bla*<sub>CTX-M</sub> gene was more predominant in ESBL producing isolates in the study location and the strains A and B ESBL producing isolates were prevalent among children with suspected cases of diarrheic in the study location.

**Keywords:** *Escherichia coli*; diarrhea; antimicrobial; Nasarawa; extended-spectrum beta-lactamase; Nigeria.

## 1. INTRODUCTION

Diarrhoea disease cause by *Escherichia coli* (*E. coli*) is serious global health problem in children and adults worldwide [1] with highest mortality rate in Sub-Sahara Africa [2,3]. This disease account 1.7 billion annual global cases with estimated 1.3 million deaths of which 25% of this were mainly occurring in Low- and Middle-Income Countries (LMICs) [4,2].

“Worldwide, reports have shown that *E. coli* causing diarrhoea are classified into six different pathotypes namely; enteroaggregative *Escherichia coli* (EAEC), Enteroinvasive *Escherichia coli* (EIEC) Enterohemorrhagic *Escherichia coli* (EHEC)/ Shiga-toxin producing *Escherichia coli* (STEC), enteropathogenic *Escherichia coli* (EPEC), enterotoxigenic *Escherichia coli* (ETEC) and diffusely adherent *Escherichia coli* (DAEC) [5,6,1]. Among the diarrheagenic *E. coli* pathotypes, EAEC along with the well-established ETEC and EPEC cause a substantial health burden of infant diarrheal cases and a variety of animal’s species” [7].

“The antimicrobials such as beta-lactams and fluoroquinolones are commonly as therapeutic options of infections caused by Gram-negative Enterobacteriaceae in both human and veterinary medicine” [8].

“Antimicrobial resistance in enteric bacteria is a serious global problem that have associated with

increased prolonged hospitalization, increased in morbidity and mortality; and high cost of treatment” [9,3].

The emergence and spread of ESBL strains of diarrheagenic *E. coli* is a serious challenge, and has led to increase severity of infection, cost of treatment, and duration of diarrhoea episodes [10,11,12]. Therefore, the focus of this study is on the prevalence and molecular diversity of ESBL producing *E. coli* isolated from stool of suspected cases of diarrhoea among children of <5 years in Nasarawa South Senatorial zone, Nasarawa State, Nigeria.

## 2. MATERIALS AND METHODS

### 2.1 Materials

#### 2.1.1 Bacteria isolates

A total of eight (8) ESBL producing *E. coli* from stool of children with suspected cases of diarrheic from our previous study [13] maintained at -80 °C were used for this study.

#### 2.2 DNA extraction

The DNA of phenotypic ESBL producing isolates was extracted using boiling method with minor modification [14]. One thousand microlitre (1000 µL) of an overnight broth culture of the

**Table 1. Primers and amplicon sizes for Extended spectrum  $\beta$ -lactamase genes**

Target genes	Gene sequence	Amplicon size(bp)	References
<i>bla<sub>TEM</sub></i>	F 5'-TCGGGGAAATGTGCGCG-3' R 3'-TGCTTAATCAGTGAGGCACC-5'	972	Abrar et al., 2019 [15]
<i>bla<sub>SHV</sub></i>	F 5'-GGGTTATTCTTATTTGTCGC-3' R 3'-TTAGCGTTGCCAGTGCTC-5'	615	Abrar et al., 2019 [15]
<i>bla<sub>CTX-M</sub></i>	F 5'-ACGCTGTTGTTAGGAAGTG-3' R 3'-TTGAGGCTGGGTGAAGT-5'	757	Abrar et al., 2019 [15]

isolates were centrifuge at 14000 rpm for 3 min. The supernatant was discarded, and the harvested cell pellet was resuspended in 1000  $\mu$ L sterile distilled water and centrifuged at 14000 rpm for 10 min. The supernatant was discarded carefully. The pellet was resuspended in 100  $\mu$ L of sterile distilled water by vortexing. The tube was centrifuged again at 14000 for 10 min, and the supernatant was discarded carefully. The cells were re-suspended in 500 $\mu$ L of normal saline and heated at 95 °C for 20 min. The heated bacterial suspension was cooled on ice for 10 mins and spun for 3 min at 14000 rpm. The supernatant containing the DNA 52 was transferred to a 1.5ml microcentrifuge tube and stored at -20 °C for other downstream reactions. The purity and concentration of extracted DNA was quantified using the Nanodrop 1000 spectrophotometer.

### 2.3 Amplification and Separation of Extended-Spectrum Beta-Lactamase Genes

The single plex Polymerase Chain Reaction (PCR: ABI 9700 Applied Biosystems thermal cycler) were performed to amplify the presence of ESBL genes such as *bla<sub>CTX-M</sub>*, *bla<sub>SHV</sub>* and *bla<sub>TEM</sub>* in the phenotypic ESBL producing isolates using the following forward (F) and reverse (R) primers as shown in Table 1 under the following condition with minor modification [14]; initial denaturation step at 95 °C for 3 minutes, followed by 35 cycles of denaturation at 94 °C for 3 sec, annealing at 58 °C for 30 sec, initial extension at 55 °C for 30 sec, final extension at 68 °C for 7 minutes and hold at 4 °C indefinitely for cooling. The amplified ESBL genes were separated on 1% agarose gel at 120 V for 15 minutes and visualized on a UV trans illuminator.

### 2.4 16s rRNA Amplification

The amplification of 16s rRNA of the ESBL producing isolates was carried out using the F:

AGAGTTTGATCMTGGCTCAG and R: CCGTTA CCTTGTTACGACTT and the PCR conditions were as follows: Initial denaturation, 96°C for 5 min; denaturation, 95 C for 30 sec; annealing, 52°C for 30 sec; extension, 72°C for 30 sec for 25 cycles and final extension, 72°C for 5 minutes [14]. The amplified 16s rRNA of the phenotypic ESBL isolates were separated on 1% agarose gel at 120 V for 15 minutes and visualized on a UV trans illuminator.

### 2.5 Restriction Fragment Length Polymorphism

The Restriction Fragment Length Polymorphism (RFLP) assay of the ESBL producing isolates by digesting the amplified 16s rRNA of the ESBL isolates with restriction enzymes ECOR1 following a method described by Nkene et al. [14]. The restriction fragments were separated on 1% agarose gel and visualized on a UV trans illuminator.

## 3. RESULTS AND DISCUSSION

### 3.1 Prevalence of Extended Spectrum $\beta$ -Lactamase Genes


The prevalence of ESBL genes in phenotypic ESBL producing isolates was determined using descriptive statistics and the overall prevalence. *bla<sub>CTX-M</sub>*, *bla<sub>SHV</sub>* and *bla<sub>TEM</sub>* were 4(50.0%) and 3(37.5%) respectively as shown in Fig. 1. Similarly, isolates with co-carriage of *bla<sub>SHV</sub>/bla<sub>TEM</sub>* (1, 12.5%) were found to be more prevalent, but none of the isolates were co-carriage of *bla<sub>TEM</sub>/bla<sub>CTX-M</sub>*, *bla<sub>SHV</sub>/bla<sub>CTX-M</sub>* and *bla<sub>CTX-M</sub>/bla<sub>SHV</sub>/bla<sub>TEM</sub>*.

### 3.2 Molecular Diversity of Extended-Spectrum $\beta$ -Lactamase Genes producing *Escherichia coli*


To determine the diversity of ESBL producers from children with suspected carrier of diarrheic in Nasarawa South, Nigeria RFLP approach was

used to digest the amplified 16rRNA sequence with ECOR1 and the different strains of isolates based on the size of fragments generated were assigned as A (200bp, 300bp, 500bp, 700bp) and B (100bp, 200bp,300bp, 600bp) of which the

proportion of both A and B were 4(50.0%) as shown in Fig. 2 and present both in General Hospital Doma (GHD) and Dalhatu Araf Specialist Hospital Lafia (DASHL) as shown in Plate 4.



**Fig. 1. Prevalence of Extended spectrum  $\beta$ -lactamase genes in *Escherichia coli* from children with suspected cases of diarrheic in Nasarawa South, Senatorial Zone, Nigeria**  
 ESBL= Extended spectrum  $\beta$ -lactamase




**Fig. 2. Prevalence of strains of Extended spectrum  $\beta$ -lactamase producing *Escherichia coli* from children with suspected cases of diarrheic in Nasarawa South, Senatorial Zone, Nigeria**




**Plate 1. Agarose gel electrophoresis of the amplified ESBL gene bacteria isolates. Lane M represents 1500bp DNA molecular ladder. Lane E1, E2, E4 and E8 represent the expression of the *bla<sub>SHV</sub>* (615bp) gene for *Escherichia coli*. While other Lanes did not show the expression of any ESBL genes**

Key: E1-E4=DASHL; -E1&E2=GHD, E8=GHO; DASHL= Dalhatu Araf Specialist Hospital Lafia, GHD=General Hospital Doma, GHO=General Hospital Obi




**Plate 2. Agarose gel electrophoresis of the amplified ESBL gene in *Escherichia coli* isolates. Lane M represents 1500bp DNA molecular ladder. Lane E1, E5 and E7 represent the expression of the bla<sub>TEM</sub> (972bp) gene for *Escherichia coli*. While other Lanes did not show the expression of any ESBL genes**

Key: E1-E4=DASHL; E5-E7=GHD, E8=GHO; DASHL= Dalhatu Araf Specialist Hospital Lafia, GHD=General Hospital Doma, GHO=General Hospital Obi



**Plate 3. Agarose gel electrophoresis of the amplified ESBL gene in *Escherichia coli* isolates. Lane M represents 1500bp DNA molecular ladder. Lane E3, E4, E6 and E8 represent the expression of the bla<sub>CTX-M-4</sub> (501bp) gene for *Escherichia coli*. While other Lanes did not show the expression of any ESBL genes**

Key: E1-E4=DASHL; E5-E7=GHD, E8=GHO; DASHL= Dalhatu Araf Specialist Hospital Lafia, GHD=General Hospital Doma, GHO=General Hospital Obi



**Plate 4. Restriction Fragment Length Polymorphism (RFLP) analysis of *Escherichia coli* isolates cluster. Lane M represents 1500bp DNA molecular ladder. Each lane corresponds to an isolate' DNA. Identical and unidentical banding patterns were evident in all samples from Lane E1 to Lane E6**

Key: E1-E4=DASHL; E5-E7=GHD, E8=GHO; DASHL= Dalhatu Araf Specialist Hospital Lafia, GHD=General Hospital Doma, GHO=General Hospital Obi

### 3.3 Discussion

This study investigates the prevalence and molecular diversity of ESBL producing *E. coli* isolates from children with suspected cases of diarrheic. The results of our study on detection of ESBL genes in ESBL producing *E. coli* with suspected cases of diarrheic shows that bla<sub>CTX-M</sub>,

bla<sub>SHV</sub> and bla<sub>TEM</sub> were detected and this suggest that the genes may be responsible for extended-spectrum cephalosporin resistance in *E. coli* with suspected cases of diarrheic in the study location.

The result of our study also shows that bla<sub>CTX-M</sub> gene was found to be more prevalent than other

ESBL gene and this however contradict the previous study conducted by Abimiku et al. [7] who reported high prevalence of *bla<sub>SHV</sub>* in *E. coli* with suspected cases of diarrheic. The results of our study also agree with previous study that reported *bla<sub>CTX-M</sub>* as most prevalent ESBL gene in intestinal *E. coli* causing diarrheic, blood stream pathogenic *E. coli* [16,17,18,19].

The results of our study on the RFLP of the ESBL producing isolates shows that the isolates were distributed into two different strains namely A and B with equal proportion and these findings contradict with the study conducted by Abimiku et al. [7] and Castanheira et al [20] and 2021 respectively), who reported the distribution of the ESBL strain A-J. The presence of the above strains suggest that they are likely response for suspected cases of diarrheic in the study location [21].

#### 4. CONCLUSION

The results of this study shows that *bla<sub>CTX-M</sub>* gene was more predominant in ESBL producing isolates in the study location and the strains A and B ESBL producing isolates were prevalent among children with suspected cases of diarrheic in the study location.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Dela H, Egyir B, Majekodunmi AO, Behene E, Yeboah C, Ackah D, Bongo RN, Bonfoh B, Zinsstag J, Bimi L, Addo KK. Diarrhoeagenic *E. coli* occurrence and antimicrobial resistance of Extended Spectrum Beta-Lactamases isolated from diarrhoea patients attending health facilities in Accra, Ghana.2022. PLoS One. 17(5):e0268991.
2. Ogunbiyi TS, Fayemi OE, Akanni GB, Ayolabi CI, Hald T. Molecular Characterization of Hetero-Pathogenic and Diarrheagenic *Escherichia coli* Pathotypes in Diarrheic Children under Five Years and Exposure Environment in Ogun State, South-West Nigeria. 2023. Pathogens. 2023;12(11):1358.
3. Zelelie TZ, Eguale T, Yitayew B, Abeje D, Alemu A, Seman A, Jass J, Mihret A, Abebe T. Molecular epidemiology and antimicrobial susceptibility of diarrheagenic *Escherichia coli* isolated from children under age five with and without diarrhea in Central Ethiopia.2023. PLoS One. 2023;18(7):e0288517.
4. Mokomane M, Kasvosve I, Melo ED, Pernica JM, Goldfarb DM. The global problem of childhood diarrhoeal diseases: emerging strategies in prevention and management. Therapeutic advances in infectious disease. 2018 Jan;5(1):29-43.
5. Sani A, Onaolapo JA, Ibrahim YK, Idris HW, Igwe JC, Nworie A. Prevalence of *Escherichia coli* pathotypes among children with diarrhoea in Zaria, Nigeria. British Journal of Medicine and Medical Research. 2015 Jan 29;7(1):17-24.
6. Abimiku RH, Ngwai YB, Nkene IH, Tاتفeng YM. Molecular detection of diarrheagenic pathotypes of *Escherichia coli* from diarrheic patients in Keffi, Nigeria. Journal of Microbiology and Biomedical Research. 2016 Dec;2(3):1-6.
7. Abimiku RH, Ngwai YB, Nkene IH, Bassey BE, Tsaku PA, Ibrahim T, Tama SC, Ishaleku D, Pennap GR. Molecular Diversity and extended spectrum beta-lactamase resistance of diarrheagenic *Escherichia coli* from patients attending selected health care facilities in Nasarawa state, Nigeria. International Journal of Pathogen Research. 2019 Sep 2;3(1):1-8.
8. Egwu E, Ibiyam FA, Moses IB, Iroha CS, Orji I, Okafor-Alu FN, Eze CO, Iroha IR. Antimicrobial susceptibility and molecular characteristics of beta-lactam-and fluoroquinolone-resistant *E. coli* from human clinical samples in Nigeria. Scientific African. 2023 Sep 1;21:e01863.
9. Chellapandi K, Dutta TK, Sharma I, De Mandal S, Kumar NS, Ralte L. Prevalence of multi drug resistant enteropathogenic and enteroinvasive *Escherichia coli* isolated from children with and without diarrhea in Northeast Indian population.

- Annals of clinical microbiology and antimicrobials. 2017 Dec;16:1-9.
10. Ugwu MC, Omanukwue C, Chimezie C, Okezie U, Ejikeugwu CP, Nnnabuife-Iloh E, Esimone CO. Poultry farm and poultry products as sources of multiple antimicrobial-resistant Salmonella and S. aureus. J Trop Dis. 2019;7(3):1-23.
  11. Heredia N, García S. Animals as sources of food-borne pathogens: A review. Animal nutrition. 2018 Sep 1;4(3):250-5.
  12. Odetoyin B, Ogundipe O, Onanuga A. Prevalence, diversity of diarrhoeagenic Escherichia coli and associated risk factors in well water in Ile-Ife, Southwestern Nigeria. One Health Outlook. 2022 Feb 8;4(1):3.
  13. Zakou AT, Ngwai YB, Nkene IH, Ishaleku D, Abimiku RH, Ekeleme IK. Antimicrobial Resistance and Phenotypic Detection of Extended Spectrum Beta-Lactamase in Escherichia coli from Children with Cases of Diarrhea in Nasarawa-South, Nasarawa State, Nigeria. Asian Journal of Biochemistry, Genetics and Molecular Biology. 2024 Jul 25;16(8):24-32.
  14. Abrar S, Ain NU, Liaqat H, Hussain S, Rasheed F, Riaz S. Distribution of bla CTX- M, bla TEM, bla SHV and bla OXA genes in Extended-spectrum- $\beta$ -lactamase-producing Clinical isolates: A three-year multi-center study from Lahore, Pakistan. Antimicrobial Resistance & Infection Control. 2019 Dec;8:1-0.
  15. Nkene IH, Ngwai YB, Bassey EB, Abimiku H, Ibrahim T, Yahaya I. Molecular Characterization of Escherichia coli with Ciprofloxacin and Extended-Spectrum Cephalosporins Co-resistance from Patients Attending Tertiary Hospitals in Nasarawa State, Nigeria. GSJ. 2020 May;8(5).
  16. Yu K, Huang Z, Xiao Y, Bai X, Gao H, Wang D. Epidemiology and molecular characterization of CTX-M-type ESBLs producing Escherichia coli isolated from clinical settings. Journal of Global Antimicrobial Resistance. 2024 Mar 1;36:181-7.
  17. Kuenzli E. Antibiotic resistance and international travel: Causes and consequences. Travel medicine and infectious disease. 2016;14(6):595-8.
  18. DeFrancesco AS, Tanih NF, Samie A, Guerrant RL, Bessong PO. Antibiotic resistance patterns and beta-lactamase identification in Escherichia coli isolated from young children in rural Limpopo Province, South Africa: The MAL-ED cohort. South African Medical Journal. 2017 Mar 1;107(3):205-14.
  19. Hamad WF. Genotypic and phenotypic study of E. coli isolated from children suffering from severe diarrhea with some antibiotic resistant gene. World Journal of Advanced Research and Reviews. 2022; 15(1):683-93.
  20. Castanheira M, Simner PJ, Bradford PA. Extended-spectrum  $\beta$ -lactamases: an update on their characteristics, epidemiology and detection. JAC-antimicrobial resistance. 2021 Sep 1;3(3):dlab092.
  21. David EE, Yameen MA, Igwenyi IO, Okafor AC, Obeten UN, Obasi DO, Ezeilo UR, David Ch N. The frequency of virulent genes and antimicrobial resistance patterns of diarrheagenic Escherichia coli isolated from stools of children presenting with diarrhea in a tertiary hospital in Abakaliki, Nigeria. Int J One Health. 2020 Jul 1;6(2):147-52.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/123624>