

Ureteric Transection with Jejunal Injury Secondary to Bicycle Handle Bar Penetrating Injury in a 13 Year Old Boy

Ojas Vijayanand Potdar ^{a*},
Mohammed Ayub Karamnabi Siddiqui ^a,
Vivek Shaw ^a, Kaustubh Vaidya ^a, Prashant Sarawade ^a,
Shashank Sharma ^a and Amrita Patkar ^b

^a Grant Medical College and J. J. Group of Hospitals, Mumbai, India.

^b Bharatratna Dr. Babasaheb Ambedkar Municipal General Hospital, Kandivali (W.),
Mumbai, Maharashtra, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97784>

Case Report

Received: 15/01/2023
Accepted: 22/03/2023
Published: 25/03/2023

ABSTRACT

Background: Ureteric trauma is rare, occurring in <1% of all traumas. Bicycle handlebar injury is a unique trauma mechanism especially noticed in childhood.

Case Presentation: We present a unique case of a 13-year-old male who sustained a penetrating abdominal injury from a bicycle handlebar. Upon initial examination there was herniation of bowel through the abdominal wound, so exploratory laparotomy was performed. There was transection of the jejunum at a distance of 10 cm from the duodeno-jejunal flexure with more than 75 % of the circumference of the jejunum and there was transection of the upper one third of left ureter near the left renal pelvis with foreign body (child's cloth remnant) seen which was initially managed by

*Corresponding author: Email: ojaspotdar@yahoo.com;

placement of an infant feeding tube as a bridge between the two transected ends owing to a defect of 3 cm as a temporary emergency salvage procedure. He then underwent per cutaneous nephrostomy insertion. After 3 months, patient underwent definitive repair by left sided pyeloureterostomy over a DJ stent.

Conclusion: This case demonstrates the emergency and delayed definitive management of ureteric transection and jejunal injury secondary to penetrating bicycle handlebar injury.

Keywords: Ureter; jejunal; bicycle; handlebar; penetrating.

ABBREVIATIONS

DJ stent : Double-J stent.

I.V : Intravenous.

PDS : Polydioxanone

1. INTRODUCTION

“Penetrating abdominal trauma secondary to bicycle handlebars is rare. Most reports regarding handlebar injuries in the literature are related to traumatic abdominal wall hernias, solid organ injury, or bowel injury” [1-5]. “Likewise, ureteric injury related to external trauma is also rare, with even the busiest adult trauma centres typically seeing fewer than 10 cases per year” [6]. We present a unique case of ureteric transection and jejunal injury secondary to bicycle handlebar injury.

2. CASE PRESENTATION

An otherwise healthy 13-year-old male presented to the Emergency department after a bicycle accident. Primary survey revealed tachycardia, hypotension and Glasgow Coma Score of 10. Secondary survey was remarkable for a deep 3-centimetre (cm) laceration in the left lower abdominal quadrant. There was active bleeding from the wound, with bowel herniating through a fascial defect. There was no remarkable haematuria. Initial workup involved a complete blood count, biochemistry profile in the form of Renal function tests and serum electrolytes and blood for grouping and cross-match. Further imaging was not done as the patient was haemodynamically unstable. The patient was taken to the operating room for exploration. An exploratory laparotomy was performed via midline abdominal incision. The underlying fascial defect was found to be approximately 5 cm. There was transection of the jejunum at a distance of 10 cm from the duodeno-jejunal

flexure with more than 75 % of the circumference of the jejunum cut and evidence of tear of Gerotas fascia of left kidney and on exploration of the retroperitoneum, there was transection of the upper one third of left ureter near the left renal pelvis with foreign body (child's cloth remnant) seen. There was no other solid organ injury. The jejunal injury was repaired after resecting the injured segment and then handsewn anastomosis done using vicryl 3-0. The ureteric transection was initially managed by placement of a 6 Fr infant feeding tube as a bridge between the two transected ends owing to a defect of 3 cm as a temporary emergency salvage procedure as the patient was haemodynamically unstable. The complex abdominal wall injury was repaired primarily after placing a 32 Fr abdominal drain. Postoperatively, the patient's condition improved and he underwent Computerised Intravenous Pyelography (CT-IVP) which showed a small leak near the repaired left ureter (see Figs. 1-4).

The patient underwent ultrasound guided left sided percutaneous nephrostomy insertion after cystoscopy with Retrograde pyelography also confirmed the leak from the left ureter (Fig. 5). Patient was discharged after 4 weeks of admission with percutaneous nephrostomy in situ. The patient underwent definitive surgery through a left subcostal incision and removal of the infant feeding tube, Ureteropyelostomy was done after adequate mobilisation of the left kidney and left ureter using 3-0 polydioxanone (PDS) suture over a double-J stent and a drain was left in place in the retroperitoneum (Fig. 6). Post-operatively the patient recovered well and was discharged five days later. The double-J stent was removed at six postoperative weeks. Concurrent cystoscopy and retrograde ureterogram demonstrated a normal ureter with no evidence of stricture.




Fig. 1. CT IVP image




Fig. 2. First delayed scan CT IVP



Fig. 3. Second delayed scan-CT IVP image after clamping the drain tube




Fig. 4. Reconstructed CT-IVP image showing left sided repaired ureteric injury by a bridging IFT and leakage of contrast through the distal part of the repair into the left perinephric drain




Fig. 5. Left sided RGP image confirming leakage




Fig. 6. Intraoperative image showing left ureteropyelostomy after adequate mobilisation of the left kidney and left ureter using 3-0 polydioxanone (PDS) suture over a double-J stent

3. DISCUSSION

According to our review of literature, most bicycle-related incidents described in the literature deal with head trauma. "Handlebar injury is also an important mechanism of bicycle-related trauma, and the associated morbidity is underappreciated. Most cases described in the literature are secondary to blunt trauma" [1-5]. "The most common injuries described include bowel perforation, solid organ injury, and traumatic abdominal wall hernia" [1-5]. There are no studies on penetrating injuries due to bicycle handlebar injuries. In a retrospective review by Nadler et al., [3] "31% of children sustaining direct-impact handlebar injuries required operative intervention". "Ureteric injuries account for less than 1% of all urologic traumas, [7] with the majority being a result of penetrating trauma". "Data compiled from the National Trauma Database shows the varying mechanisms of ureteric injury" [8]. "It shows that the most common mechanism of ureteric injury in trauma is penetrating injury in the form of gunshot wounds whereas in case of blunt trauma, it is motor vehicle collisions. The diagnosis and management of ureteric injury is challenging and often delayed or missed. A retrospective review of traumatic ureteric injury by Medina et al. found that only 40% of patients with ureteric injury had positive findings on preoperative imaging studies" [9]. "Computed tomography with delayed phase images must be included if there is any suspicion of ureteric injury" [9]. "Cystoscopy with retrograde ureterogram/pyelogram is the most accurate way of identifying ureteric injury, but may not be feasible if patient is hemodynamically unstable" [9]. "Due to a high rate of concomitant injuries many ureteric injuries are diagnosed intra-operatively during exploration for other reasons. This can pose a challenge for the surgeon as haemorrhage or spillage of intestinal contents can make detecting leakage of a small amount of urine difficult. The possibility of ureteric injury should be suspected if there is violation of retroperitoneum. Approximately 60% of injuries occur in the proximal ureter" [10]. In emergency setting if the patient is haemodynamically unstable, a unique approach of placing infant feeding tube as a bridge was used in our study. "The type of operative repair varies depends on the location of the injury. Regardless of the location of injury, the basic tenets of repair always involve debridement to healthy tissue and tension-free anastomosis over a stent with absorbable suture. In our patient, a unique approach of placing an infant feeding

tube as a bridge as demonstrated in this case is recommended. The type of operative repair varies depends on the location of the injury. Regardless of the location of the injury, the basic tenets of repair always involve debridement to healthy tissue and tension-free anastomosis over a stent with absorbable sutures" [10]. In our patient, a unique approach of placing an infant feeding tube to bridge the ureteric defect was used as a temporary measure as the patient was haemodynamically unstable. He did undergo a definitive repair after a period of 3 months and it was successful. Thus, this case report does add a unique perspective of emergency management of ureteric injuries.

4. CONCLUSION

A high index of suspicion is necessary for timely diagnosis of ureteric injuries. Intra-operative discovery of retroperitoneal violation warrants further exploration, with identification of ureteric trauma prompting an appropriate repair based on the location of the injury. However, in an emergency setting with hemodynamic instability, novel approaches like placing an infant feeding tube to bridge the large ureteric defect can be used to tide over the patient till definitive repair is carried out.

CONSENT

As per international standard or university standard, parental(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Debbink KP, Tashjian DB, Tirabassi MV, Gaffey R, Nahmias J. Ureteric transection secondary to penetrating handlebar injury. *Trauma Case Reports*. 2017;10:16-8. ISSN 2352-6440 Available: <https://doi.org/10.1016/j.tcr.2017.07.002>.

2. Karaman A, Karaman MK, Aslan D, Erdoğan YH, Cavuşoğlu O. Tütün. A hidden danger of childhood trauma: bicycle handlebar injuries, Surg. 2009;39(7):572–574.
3. Nadler EP, Potoka DA, Shulz BL, Morrison KE, Ford HR, Gaines BA. The high morbidity associated with handlebar injuries in children, J. Trauma. 2005;58(6):1171–1174.
4. Goliath J, Mittal V, McDonough J. Traumatic handlebar hernia: a rare abdominal wall hernia, J. Pediatr. Surg. 2004;39(10):20–22.
5. Clarnette TD, Beasley SW. Handlebar injuries in children: patterns and prevention, Aust. N. Z. J. Surg. 1997;67(6):338.
6. Brandes SB, Chelsky MJ, Buckman RF, Hanno RM. Ureteral injuries from penetrating trauma, J. Trauma. 1994;36(6):766–769
7. Presti JC, Jr., Carroll PR, McAninch JW. Ureteral and renal pelvic injuries from external trauma: diagnosis and management, J. Trauma. 1989;29(3):370–374.
8. Siram SM, Gerald SZ, Greene WR, et al., Ureteral trauma: patterns and mechanisms of injury of an uncommon condition, Am. J. Surg. 2010;199:566–570.
9. Medina D, Lavery R, Ross SE, Livingston DH. Ureteral trauma: Preoperative studies neither predict injury nor prevent missed injuries, J. Am. Coll. Surg. 1998;186:641–644.
10. Pereira BMT, Ogilvie MP, Gomez-Rodriguez JC, et al., A review of ureteral injuries after external trauma, Scand. J. Trauma. 2010;18(6).

© 2023 Potdar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97784>