



Normal and Pathological Vaginal Microflora in the Bitch-A Review

Anton Antonov^{1*}

¹*Department of Obstetrics, Reproduction and Reproductive Disorders, Faculty of Veterinary Medicine, Trakia University, 6000, Stara Zagora, Bulgaria.*

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/JAMB/2019/v15i130074

Editor(s):

- (1) Dr. Ana Claudia Correia Coelho, Department of Veterinary Sciences, University of Trás-os-Montes and Alto Douro, Portugal.
- (2) Dr. Hung-Jen Liu, Distinguished Professor, Institute of Molecular Biology, National Chung Hsing University, Taiwan.

Reviewers:

- (1) Aiyenuro Ademola Emmanuel, Federal University of Technology, Nigeria.
- (2) Alessia Mariacher, Istituto Zooprofilattico Sperimentale Regioni Lazio e Toscana Centro di Referenza Nazionale per la Medicina Forense Veterinaria, Italy.
- (3) Tsaku Paul Alumbugu, Coal City University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46825>

Received 17 November 2018

Accepted 29 January 2019

Published 26 February 2019

Review Article

ABSTRACT

The vagina is not only a place for the development of various microorganisms, but also serves as a border before reaching the uterus of all mammals. Determining the qualitative and quantitative specifics of vaginal bacterial flora is of great importance for a better understanding of genital pathology mechanism, clinical diagnosis, management and antibiotic treatment of reproductive diseases in the bitch.

Keywords: *Vagina; bitch; bacterial flora.*

1. INTRODUCTION

A comparatively limited analysis of the vaginal microflora in the bitch has been performed in the literature. The type of isolated microorganisms from healthy or bitches with reproductive

diseases is important because in the presence of some predisposing factors, the development of opportunistic bacteria may escalate and cause different pathological processes in the reproductive system and the whole organism [1].

*Corresponding author: E-mail: anton.antonov@abv.bg;

2. MATERIALS AND METHODS

Various types of microorganisms are isolated from the vagina of bitches [2-9]. Age specificity has been determined [10]. In a large percentage of non-sexually mature female dogs more coagulase-positive staphylococci have been found compared to those who have reached puberty [11]. In one study, bacteria were found in 86% of the animals before the first estrus occurrence and in 84% after the onset of puberty [12]. No differences were found between the microorganisms isolated from animals prior to the onset of puberty, spayed or intact females and those living in the cranial or caudal part of the vagina [10,13]. Other authors detected microorganisms between 89.7% [14] and 96% [3] of the vaginal samples tested. Some found the existence of monocultures [6,15,16] and others mixed isolates [12,17], both from aerobes and anaerobes [2,3,18].

3. RESULTS AND DISCUSSION

The most frequently isolated bacteria from the vagina of bitches are *E. coli*, *Staphylococcus spp.*, *Streptococcus spp.*, *Pasteurella spp.*, *Proteus spp.*, *Bacillus spp.*, *Corynebacterium spp.*, *Pseudomonas spp.* and *Micrococcus spp.* Less common isolated are *Neisseria spp.*, *Klebsiella spp.* and *Moraxella spp.* [8-10,14,15,17-26]. Some of these bacteria are isolated from both fertile and infertile dogs [2] with similar types of aerobes as part of the vaginal microflora of both groups of animals [11].

Isolation of *Mycoplasma spp.* [3,20,27,28] and *Ureaplasma spp.* [27,29,30] has been reported in 30 to 88% of the samples of bitches without reproductive problems. *Mycoplasma* isolates from the vaginas of healthy bitches did not vary during the various stages of the estrous cycle [31]. Rarely anaerobic bacteria were found in healthy bitches, such as *Bacteroides spp.* and *Peptostreptococcus spp.* [3,23].

The presence of lactic acid bacteria (*Lactobacillus* and *Enterococcus*) in the vagina of the bitch has been reported. Their importance is expressed in suppressing the development of pathogenic microorganisms in the urinary and reproductive systems by synthesizing bacteriocins and hydrogen peroxide and disrupting their adhesion to epithelial cells [32,33].

In the bitch, the presence of fungus is rarely found [2,12,30]. In a mycological study before and during the first estrus, a presence of *Malassezia spp.* [34] and *Candida spp.* [35,36] was recorded as a very rare finding. There are also reports for isolation of *Penicillium spp.* and *Malassezia pachydermatis* [37]. Other authors found that *Malassezia pachydermatis* is the most commonly isolated yeast as a part of the normal canine surface microbiota, followed by *Candida parapsilosis*, *Candida tropicalis*, *Candida albicans*, *Saccharomyces cerevisiae* and *Rhodotorula spp.* [38].

In some studies the authors argue that there may be a difference in vaginal microorganisms during the different stages of the estrous cycle in bitches [10,15,39]. According to some authors there is no such a difference [13,35,40]. Pure bacterial cultures can be isolated from the vagina of healthy bitches during proestrus and anestrus, while mixed bacterial cultures are found during estrus and diestrus [39]. *E. coli* and *Streptococcus spp.* are isolated during all the stages of the estrous cycle in the bitch, but not during pregnancy [41]. According to the same authors, β -hemolytic streptococci are often found in estrus and early diestrus, and α -hemolytic streptococci in the other stages of the cycle.

Other authors reported only an increase in the number of isolates during proestrus and estrus [1,3,14,22,42,43]. There are reports that the smallest number of bacterial isolates were found through proestrus [44,45]. According to them the reason for this is the continuous passing through the vagina of fluid from the uterus. The highest number of isolates they found during diestrus and pregnancy, which according to them is due to the high progesterone concentrations. These levels of progesterone and the pH values typical for these periods provide favorable conditions for the development of microorganisms in the sexual system [46]. It is also suggested that immunosuppression occurs during the luteal phase, which further promotes the proliferation of microorganisms [45].

It is impossible to find an etiological connection between the isolation of a particular microorganism and the occurrence of a specific genital disease in the bitch [18] and is very difficult to associate a disease with a specific bacterial isolate due to the wide variety of microorganisms into the vagina of bitches with or without reproductive diseases [10]. Only *Brucella*

canis is mentioned as a specific microorganism and etiological agent of a disease [23].

Different investigations have shown that microorganisms into the vagina of clinically healthy bitches are isolated also from patients with genital system inflammation [13,17]. The most common are *E. coli*, *Staphylococcus spp.*, *Streptococcus spp.* [17,25,47,48] and *Pasteurella spp.* [6,17]. Less common isolated are *Pseudomonas spp.* [6,22], *Proteus spp.*, *Citrobacter spp.* and *Enterobacter spp.* [6,13]. There is a higher percentage of *Staphylococcus spp.* in bitches with vaginitis compared to healthy ones [6]. They are also predominant isolates in young compared to adult animals [18]. Limited information is available regarding the vaginal microbiota of normal spayed dogs and spayed dogs with recurrent urinary tract inflammations (UTI), but the results showed that vaginal microbiota of spayed female dogs with recurrent UTI was similar to the control population of normal, spayed female dogs [49]. The microbiome of the endometrium is distinctly different from that of the vagina [50].

In a study of bitches with chronic vaginitis were isolated *Streptococcus spp.*, *Staphylococcus spp.*, *Proteus spp.* and *Bacillus spp.* and in healthy animals during estrus-*Klebsiella spp.*, *Staphylococcus spp.*, *Corynebacterium spp.* and *Proteus spp.* [13,51,52]. Some authors reported that the isolated microorganism during a disease of the reproductive system should not necessarily be considered as a cause of the disease [18].

Because most of the bacterial species in the caudal part of the genital tract in the bitch are opportunistic microorganisms, the interpretation of the results without additional information from the anamnesis and the clinical finding is generally not possible. An exception in this aspect is *Brucella canis* [53].

High bacterial growth, especially monocultures, may serve as a pathological finding associated with the presence of vaginal discharge (most commonly purulent) and finding leukocytes in exfoliative vaginal cytology, with the exception during the early diestrus [1].

4. CONCLUSION

The importance of infections in the pathogenesis of infertility and reproductive disorders in the bitch may be exaggerated. It is possible some

microorganisms to penetrate through a transcervical migration from the vagina into the uterine cavity leading to abortion and development of various pathologies and infertility, but it should not be an argument bitches to be treated prophylactically with antibiotics before insemination.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Antonov, A. Investigations of the values and the dynamics of certain vaginal parameters in bitches; 2015. Thesis.
2. Osbaldiston, G. Vaginitis in a bitch associated with *Haemophilus spp.* Am J Vet Res. 1971;32:2066-2070.
3. Baba E, Hata H, Fukata T, Arakawa A. Vaginal and uterine microflora of adult dogs. Am J Vet Res. 1983;44:606-609.
4. Olson P, Thrall M, Wykes P. Vaginal cytology: Part I: A useful tool for staging the canine estrous cycle. Compend. Contin. Educ. Pract. Vet. 1984;6:288-298.
5. Soderberg S. Canine breeding management. Vet Clin North Am Small Anim Pract. 1986;16: 419-433.
6. Kowitz G. Bakteriell bedingte Ursachen von Reproduktionsstörungen des Hundes Berlin, Freie Univ; 1998. Thesis.
7. Gunzel-Apel A, Lubke A, Rohde J. Vergleichende Untersuchungen der vaginalen und uterinen Zytologie und Keimflora im Sexualzyklus und Puerperium von Beaglehündinnen. Tierärztl. Prax. 1999;27:112-119.
8. Popova T, Todorov T. In vitro antimicrobial activity of enrofloxacin against clinical isolates from dogs and cats in Bulgaria. Folia Veterinaria. 2008;52(3-4):124-128.
9. Groppetti, D, Pecile, A, Barbero, C, Martino, P. Vaginal bacterial flora and cytology in proestrus bitches: Role on fertility. Theriogenology. 2012;77:1549-1556.

10. Olson P, Marther E. Canine vaginal and uterine bacterial flora. *J Am Vet Med Assoc.* 1978;172:708-711.
11. Feldman E. Infertility. In: *Textbook of Veterinary Internal Medicine.* Vol. 2, 3rd Ed., S.J. Ettinger (ed). Philadelphia, W.B. Saunders Co. 1989;1838-58.
12. Siesenop U, Gunzel-Apel A, Hackbarth H, Spitz G, Bohm K. Die mikrobielle Vaginalflora bei Hündinnen einer Beaglegruppe vor und im Verlauf des ersten Sexualzyklus, *Kleintierpraxis.* 1996;41:449-455.
13. Johnson C. Diagnosis and treatment of chronic vaginitis in the bitch. *Vet Clin North Am Small Anim Pract.* 1991;21:523-531.
14. Antonov A, Koleva M, Georgiev P. Vaginal aerobic bacterial flora during the sexual cycle stages in bitches of different breeds. *Bulgarian Journal of Veterinary Medicine.* 2013;16(1):69-76.
15. Bjurstrom L, Linde-Forsberg C. Long term study of aerobic bacteria of the genital tract of breeding bitches. *Am J Vet Res.* 1992;53:665-669.
16. Zahr E. Untersuchungen zur Infertilitätsdiagnostik bei der Hündin Berlin; 2001. Thesis.
17. Hirsh D, Wiger N. The bacterial flora of the normal canine vagina compared with that of vaginal exudates. *J Small Anim Pract.* 1977;18(1):25-30.
18. Olson P, Thomas T, Husted P, Nett T. Clinical evaluation of infertility in the bitch. *Clinical signs and diagnosis in small animal practice,* New York, NY 10036, USA, Churchill Livingstone Inc. 1988;631-654.
19. Platt A, Simpson R. Bacterial flora of the canine vagina. *Southwest Vet.* 1974;27:76-77.
20. Ling G, Ruby A. Aerobic bacterial flora of the prepuce, urethra and vagina of normal adult dogs. *Am J Vet Res.* 1978;39:695-698.
21. Schaefer V, Kirpal G, Pantel M. Die bedeutung des katheterisierens für das bakteriologische ergebnis der harnuntersuchung. *Kleintierpraxis.* 1978;23:181-188.
22. Allen W, Dagnall G. Some observations on the aerobic bacterial flora on the genital tract of the dog. *J Small Anim Pract.* 1982;23:325-335.
23. Van Dujkeren E. Significance of the vaginal bacterial flora in the bitch: A review. *Vet Rec.* 1992;131:367-369.
24. Gandotra V, Singla V, Kochhar H, Chauhan F, Dwivedi P. Hematological and bacteriological studies in canine pyometra. *Indian Vet J.* 1994;71:816-8.
25. Laznicka A, Huml O, Nesnalova E. Microflora of genital organs of bitches and its relationship to reproductive disorders. II. Vaginitis. *Veterinarstvi.* 1995;45:210-212.
26. Wendt K, Stellmacher H. Infektiöse Belastungen der Reproduktion bei der Hündin *Tierärztl. Prax.* 1996;24:79-84.
27. Doig P, Ruhnke, H, Bosu, W. The genital Mycoplasma and Ureaplasma flora of healthy and diseased dogs. *Can J Comp Med.* 1981;45(3):233-238.
28. Rosendahl, S. Canine mycoplasmas: Their ecologic niche and role in disease. *J Am Vet Med Assoc.* 1982;180:1212-1213.
29. Allen W. Vaginale Veränderungen von klinischer Bedeutung. In: Allen, W. E. *Fruchtbarkeit und Geburtshilfe beim Hund* Enke Verlag Stuttgart. 1994;12-18.
30. Davidson, A. Frustrating case presentations in canine theriogenology. *Vet Clin North Am Small Anim Pract.* 2001;310(2):411-420.
31. Maksimovic Z, Maksimovic A, Halibasic A, Rifatbegovic M. Genital mycoplasmas of healthy bitches. *J Vet Diag Invest.* 2018;30(4):651-653.
32. Reid G. The scientific basis for probiotic strains of *Lactobacillus.* *Appl Environ Microbiol.* 1999;65:3763-3766.
33. Delucchi L, Fraga M, Perelmuter K, Cidade E, Zunino P. Vaginal lactic bacteria in healthy and ill bitches and evaluation of in vitro probiotic activity of selected isolates. *Can Vet J.* 2008;49:991-994.
34. Bohm K, Ruffler A, Siesenop U, Beckmann G. Mikrobiologische Untersuchungen zur Zusammensetzung der Vaginalflora der Hundin. *WSAVA Weltkongress und FKDVG, Berlin.* 1993;275.
35. Cleff M, Lima A, Faria R, Meinerz A, Antunes F, Araujo F, Nascente P, Nobre M, Meireles M. Isolation of *Candida* spp. from vaginal microbiota of healthy canine females during estrous cycle. *Brazilian Journal of Microbiology.* 2005;36:201-204.
36. Gunay U, Gunay A, Ulgen M, Ozel A. Investigation of the vaginal bacterial flora at different stage of sexual cycle in the bitch. *Uludag Univ J Fac Vet Med.* 2004;23:15-19.
37. Oliveira, C De, Oliveira, C. Study of aerobic microflora and vaginal pH in healthy pubescent bitches (*Canis*

- familiaris*) during the phases of the oestrus cycle; evaluation of the sensitivity to antimicrobial agents. Thesis, Universidade de Sao Paulo, Brazil; 1991.
38. Brito E, Fontenelle R, Brilhante R, Cordeiro R, Monteiro A, Sidrim J, Rocha M. The anatomical distribution and antimicrobial susceptibility of yeast species isolated from healthy dogs. *The Veterinary Journal*. 2009;182(2):320-326.
 39. Maksimovic A, Maksimovic Z, Filipovic S, Besirovic H, Rifatbegovic M. Vaginal and uterine bacteria of healthy bitches during different stage of their reproductive cycle. *Veterinary Record*. 2012;171:1-2.
 40. Ozyurtlu, N, Yesilmen, S, Arserim Kaya N. Determination of vaginal aerobic bacterial flora view of sexual cycle in bitch in the province of Diyarbakir. *Dicle Univ Vet Fak Derg*. 2008;1:7-10.
 41. Findik M, Maral N, Keskin O, Kalender H, Erdeger J, Aslan S. The relationship between the stages of sexual cycle, the pregnancy and postpartum periods and vaginal flora in kangal breed bitches. *Turk J Vet Anim Sci*. 2003;27:761-765.
 42. Noguchi K, Tsukumi K, Urano T. Qualitative and quantitative differences in normal vaginal flora of conventionally reared mice, rats, hamsters, rabbits and dogs. *Comp Med*. 2003;53:404-412.
 43. Laurusevicius, S, Siugzdaite, J, Zilinskas, H. Correlation between different sexual cycle stages and vaginal bacterial flora in bitches of different breeds. *Veterinarija ir zootechnika*. 2008;41(63):76-79.
 44. Holst P. Vaginal cytology in the bitch. In: *Current Therapy in Theriogenology*, Vol. 2, Philadelphia, W. B. Saunders Co. 1986; 457-462.
 45. Mshelia G, Amin J, Chaudhary S. Vaginal bacterial flora of Nigeria local bitches during different stages of reproductive cycle. *IntJ Agri Biol*. 2001;3(2):183-185.
 46. Amin J, Zaria L, Malgwi R. Vaginal aerobic bacterial flora of apparently healthy cattle in various stages of the reproductive cycle in sahel of Nigeria. *Bull Anim Hlth Prod Africa*. 1996;44:15-18.
 47. Bjurstrom L. Aerobic bacteria occurring in the vagina of bitches with reproductive disorders. *Acta Vet Scand*. 1993;34(1):29-34.
 48. Ross A. Investigations in the female dog's vaginal aerobic flora and pH. Dissertation, Berlin; 2005.
 49. Hutchins R, Vaden S, Jacob M, Harris T, Bowles K, Wood M, Bailey C. Vaginal microbiota of spayed dogs with or without recurrent urinary tract infections. *J Vet Intern Med*. 2014;28(2):300-304.
 50. Lyman C, Holyoak G, Meinkoth K, Wieneke X, Chillemi K, DeSilva U. Canine endometrial and vaginal microbiomes reveal distinct and complex ecosystems. *PLoS One*. 2019;14(1):e0210157.
 51. Johnston S, Kustritz M, Olson P. *Canine and feline theriogenology*. 2nd ed. W.B Saunders Company; 2001.
 52. Gunzel-Apel A. *Fruchtbarkeitskontrolle bei der Hundin*. Vet Reihe; 1993.
 53. Johnson C, Walker R. Clinical signs and diagnosis of *Brucella canis* infection. *Compend Contin Educ Pract Vet*. 1992;14:763-773.

© 2019 Antonov; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/46825>