



## A Comparative Study of the Daily Activity Patterns of Dog Faced Baboon (*Papio anubis*) in Captivity at the Kano University Zoo and Kano Zoological Garden

Mohammed Musa Yahaya<sup>1\*</sup>, Isah Suleiman Dutse<sup>2</sup>, D. O. Oshibanjo<sup>1</sup>  
and Dan`azumi Danboyi Daya<sup>3</sup>

<sup>1</sup>Department of Animal Production, University of Jos, Jos Plateau State, Nigeria.

<sup>2</sup>Kano University of Science and Technology, Wudil, Kano State, Nigeria.

<sup>3</sup>Yobe State College of Agriculture Gijba, Yobe State, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. Author MMY designed the study, Authors ISD, DOO, and DDD performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors MMY and ISD managed the analyses of the study. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJRAF/2019/v4i130054

#### Editor(s):

(1) Dr. Nebi Bilir, Professor, Forestry Faculty, Isparta University of Applied Sciences, Turkey.

#### Reviewers:

(1) Emmanuel Danquah, Kwame Nkrumah University of Science and Technology, Ghana.

(2) James T. Anderson, West Virginia University, USA.

(3) Sandra Elizabeth Perez, Universidad Nacional del Centro de la Provincia de Buenos Aires, Argentina.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50793>

Original Research Article

Received 22 June 2019  
Accepted 30 August 2019  
Published 07 September 2019

### ABSTRACT

**Aim:** This study seeks to observed the daily activity patterns of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Wudil Zoo and Kano Zoological Garden.

**Materials and Methods:** The study was carried out daily between 6:00am to 6:00pm from December 2016 to January 2017. Digital camera was also attached to cages at the two sites. The observation in the activity patterns were recorded on recording sheet, observation is done three times a week at 20 minutes intervals.

**Results:** The findings on activity pattern of dog faced baboon (*Papio anubis*) in captivity shows that the day time activities decrease from morning to evening. Resting activities was 47.5%, movement

\*Corresponding author: E-mail: yahayam@unijos.edu.ng;

and feeding were carried out in the morning, followed by afternoon and evening with 33.3% and 19.1% activities respectively. The results from the Kano Zoological Garden, indicated that 42.7% of the activities performed by dog faced baboon in captivity are resting, this is followed by movement which accounted for 34.9% of the activities, while feeding activities account for the least with 22.4%. Similarly, about 43.2% of the recorded activities carried out by dog faced baboon in Kano University of Science and Technology Wudil, Zoo was resting, followed by movement which constituted 34.8% of the activities and feeding activity which accounted for 22%.

**Conclusion:** Due to the fact that majority of the baboons activities usually take place between morning and afternoon, it is recommended that visitors interested in baboons should plan their visitation to the Zoo pen during that time. It is also recommended that feeding and harassing of animals by the visitors should be discouraged in order to ensure consistency in their behaviour.

**Keywords:** *Papio Anubis*; feeding; movement; resting; Kano University of Science and Technology Wudil and Kano Zoological Garden.

## 1. INTRODUCTION

Activity patterns have been studied in several primate taxa including hominoids [1,2] cercopithecines [3,4,5] and colobines [6]. Time is limited for most animals [7,8]. Thus, animals are faced with the challenge of allocating the limited time to different activities. According to the optimality theory, "the amount of time that an organism spends engaged in various activities depends on the cost of the activity relative to the derived benefits in that organism's habitat" [9].

The amount of time spent on foraging activities therefore relates to the energy content of the food relative to the costs of obtaining the food plus the cost of all other activities (resting, moving or socialising). Thus, specifically, food availability and energy content are critical determinants of an animals' daily activity pattern. Therefore, factors that influence the availability of food have a strong bearing on time allocation profiles in baboons.

Due to the different costs and benefits of specific activities animals have varying time allocation profiles based on age and sex for certain activities [10]. Furthermore, since these activities cannot be performed simultaneously some individuals may allocate time between various behaviours better than others [7,8]. The costs and benefits of these activities change with changes in the ecological and social state of the environment as well as the physiological state of the animal. This gives rise to temporal and spatial variation in individual activity budgets of the animal. Baboons allocate the greater proportion of their time to foraging activities [11, 12,13,14,15]. De Hoop and Mkuzi baboon troops spent 69.8 % and 66.5 % respectively of their time foraging [14]. In a study of [12] that they

spend 69.8%, 75.2% and 43% of their time foraging, respectively. The Lodge troop spent relatively less time foraging than Alto and Hook groups.

Weather patterns have both direct and indirect influences on the activity pattern of primates. Rainfall and temperature have pervasive effects on animals [16] and so influence time allocation patterns both temporally and spatially.

This study seeks to observe the daily activity patterns of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Wudil Zoo and Kano Zoological Garden.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The study was carried out in Kano University of Science and Technology Zoo and Kano Zoological Garden. **Wudil** is located between the latitude 11°37'N and longitude 8° 58'E has a total area of 362 km<sup>2</sup> and is located within Sudan savannah region of Nigeria. The annual maximum rainfall is between 850mm-870mm with a minimum and maximum temperature of 26°C - 30°C. The relative humidity of the region is always low and ranges between 40% - 51%.

### 2.2 Materials

Standardized data collection sheet, stop clock, recording sheet and digital camera.

### 2.3 Data Collection

Sampling method was used to study the activities of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Wudil Zoo

and Kano Zoological Garden from 6:00 am to 6:00 pm between December to January 2016. The observations were recorded in standard data sheets, at 20 minutes intervals in each of the cages under study. Note: this research is limited to period when the temperature is extremely low (Hammattan period). The activity parameters recorded include: Feeding, Moving, and Resting and are described as follows:

**Feeding:** the feeding began when the animal first made contact with any part of food or other food substances, feeding bout terminated when the either moved more than one full stride, even if it was carrying some food material on its hand and mouth or stopped looking at the food material, by this definition, a switch to a new food type in the absence of either of these condition was not for bout to be consider terminated, thus a single feed bout could include more than one food type [17,9].

**Resting:** this includes behavior during which an animal was neither feeding, moving or engaged in other social behavior that include sleeping auto-grooming, looking around etc [9,10].

**Moving:** this includes all locomotion activities like walking, running, climbing, jumping and leaping but excluding short movements during feeding and locomotion during social behavior e.g when primates chased one another [9,10].

**Other activities:** other social behavior including all other activities which an animal's attention and behavior where clearly directed toward another individual. These include allo-grooming, mounting, mating, chasing, playing, aggressive or agnostic behaviours [9,10].

All the activities such as resting, movement and feeding are carried out in the morning, afternoon and evening.

## 2.4 Data Analysis

The data collected was subjected to descriptive statistics which includes frequency distribution and percentage.

## 3. RESULTS AND DISCUSSIONS

The result of the day time activities of dog faced baboon (*Papio anubis*) in captivity is presented in Table 1. The result shows that the day time activities decrease from morning to evening. Resting was 47.5% of the activities, movement

and feeding were carried out in the morning, followed by afternoon and evening with 33.3% and 19.1% activities respectively.

The result of the activities of dog faced baboon in Kano Zoological Garden shown in Tables 2 and 3, indicated that 42.7% of the activities perform by dog faced baboon in captivity are resting. This was followed by movement which accounted for 34.9% of the activities, while feeding activities account for the least with 22.4%. The result of this study is in variance with the finding of [9] who reported 50.00% for feeding and 8.50% for resting for the Kwano Forest baboons. In his study, Kwano Forest baboons spent relatively higher proportion of time feeding and lesser proportion of time resting and moving, this is probably due to the level of availability and distribution of food resources at the site compare to captive environment.

The result of the dog faced baboon activities is indicated in Table 4 and 5. It shows that 43.2% of the activities carried out by dog faced baboon in Kano University of Science and Technology Wudil, Zoo is Resting, followed by the Movement with 34.8% of the activities and the feeding activity account for 22%. This result of the dog faced baboon activities in captivity indicated in the table above however, agrees with the finding of [10] who reported highest resting period than feeding and movement period.

The Dog faced Baboon activities are higher in day time period. The baboon was observed to be very active in the morning followed by afternoon and evening. These activities which is made up of mostly movement and feeding may be due to the presence of visitors in the morning and afternoon his activity pattern morning, afternoon and evening have been commonly reported among arboreal species [17]. The daily activities of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Zoo and Kano Zoological Gerden ranged between 6:00am in the morning to 6:00pm in the evening in which most of the visitors usually pay their visit. However, [18] was of the opinion that adaptive significances of diurnal variability in primate's activities budget are poorly understood. With regards to individual activities, resting which include sleeping, looking about etc. was the most frequent activity carried out by the dog faced baboon in captivity. This may be due to the confinement in which the baboons were kept. Most of the baboon's time was spent in sitting postures, standing or playing posture. In this

position, the hind limb may be placed in variety of positions and the fore limbs of the baboon often at rest on the knees or between hind limb. However, despite the confinement, movement also constitute the substantial percentage of the dog faced baboon activities in Kano University of Science and Technology and Kano Zoological Garden. The movement which accounted for 34.9% in Kano Zoological Garden and 34.8 in the University Zoo include walking, running, climbing,

leaping and riding. This significance percentage might be due to the fact that dog faced baboon are usually regarded as one of the most entertaining animal within the Zoo.

Meanwhile, of both the animals studied, the Baboon in the Kano Zoological Garden has the slightly higher frequency of activities. This might be also due to more number of visitor.

**Table 1. Variation in day time activities of dog faced baboon (*Papio anubis*) in captivity**

Day Time	Total Number of Activities	Percentage (%)
Morning	67	47.52
Afternoon	47	33.33
Evening	27	19.15
Total	141	100.00

**Table 2. Variation in the activities of dog faced baboon (*Papio anubis*) in Kano Zoological garden/day**

Day Time	Total Number of Activities	Percentage (%)
<b>DAY 1</b>		
Feeding	32	22.7
Moving	50	35.5
Resting	59	41.8
<b>DAY 2</b>		
Feeding	32	23
Moving	49	35.2
Resting	58	41.7
<b>DAY 3</b>		
Feeding	30	21.4
Moving	52	37.1
Resting	58	41.4
<b>DAY 4</b>		
Feeding	31	22.5
Moving	46	33.3
Resting	61	44.2
<b>DAY 5</b>		
Feeding	30	21.3
Moving	53	37.6
Resting	58	41.1
<b>DAY 6</b>		
Feeding	31	22.5
Moving	47	34
Resting	60	43.5
<b>DAY 7</b>		
Feeding	32	23.3
Moving	43	31.4
Resting	62	45.2
<b>Total</b>	<b>974</b>	<b>100</b>

**Table 3. Variation in the activities of dog faced baboon (*Papio anubis*) in Kano Zoological garden/week**

Activities	Total Number of Activities	Percentage (%)
Feeding	218	22.4
Movement	340	34.9
Resting	416	42.7
Total	974	100

**Table 4. Variation in the activities of dog faced baboon in Kano University of Science and Technology Wudil, Zoo/day**

Nuber of Days/Activities	Total Number of Activities	Percentage
<b>DAY 1</b>		
Feeding	29	21
Moving	37	26.8
Resting	72	52.1
<b>DAY 2</b>		
Feeding	27	19.8
Moving	35	25.7
Resting	74	54.4
<b>DAY 3</b>		
Feeding	30	21.9
Moving	42	30.6
Resting	65	47.4
<b>DAY 4</b>		
Feeding	30	22.2
Moving	45	33.3
Resting	60	44.4
<b>DAY 5</b>		
Feeding	31	22.3
Moving	56	40.3
Resting	52	37.4
<b>DAY 6</b>		
Feeding	32	23.3
Moving	60	43.8
Resting	45	32.8
<b>DAY 7</b>		
Feeding	32	23.7
Moving	58	43
Resting	45	33.3
<b>Total</b>	<b>957</b>	<b>100</b>

**Table 5. Variation in the activities of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Zoo/week**

Activities	Total Number of Activities	Percentage (%)
Feeding	211	22
Movement	333	34.8
Resting	413	43.2
<b>Total</b>	<b>957</b>	<b>100</b>

#### 4. CONCLUSION

This study was designed to gather information on the daily activity pattern of Dog faced Baboon in

Kano University of Science and Technology and Kano Zoological Garden. From the study, the following conclusions can be made. The dog faced baboons are most active in the morning.

Also Resting constitute the most frequent activity of dog faced baboon in captivity. Most of the baboon activities have short duration.

## ACKNOWLEDGEMENTS

We thank the staff of The Kano University of Science and Technology Wudil and Kano Zoological Garden, who provided assistance by providing farm facilities. There is no sponsors involvement, the authors in this study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Goodall J. The chimpanzees of Gombe: Patterns of behaviour, Belknap Press, Cambridge;1986.
2. Clutton-Brock THC, Harvey P. Species differences in feeding and ranging behavior in primates. In: Primate Ecology (Ed. T.H.C. Clutton-Brock), New York: Academic press. 1977;557-584.
3. Post DG. Activity Patterns of Yellow Baboons (*Papio cynocephalus*) In the Amboseli National Park, Kenya. Animal Behaviour. 1981;29:357-374.
4. Van Schaik CP. Why are diurnal primates living in groups? Behaviour. 1983;87:120-144.
5. Isbell LA, Preutz JD, Young TP. Movements of vervet (*Cercopithecus aethiops*) and patas monkeys (*Erythrcebus patas*) as estimators of food resource size, density and distribution. Behaviour Ecology Sociobiology. 1993;42:123-133.
6. Strushaker TT. The red colobus monkey, University of Chicago Press, Chicago; 1975.
7. Dunbar RIM. Primate Social Systems, Cornell University Press, Ithaca;1988.
8. Dunbar RIM. Time: A hidden constraint on the behavioural ecology of baboons, Behaviour Ecology Sociobiology. 1992;31: 35-49.
9. Gessaman JA. Ecological energetics of homeotherms: A view compatible with ecological modelling, Utah University Press Monograph series. 1973;20:1-155.
10. Melle EM. socio-ecology of Olive baboons thesis department of wildlife and fisheries management, university of Ibadan; 2004.
11. Altmann J, Altmann S. Baboon Ecology: African Field Research. Karger, Basel; 1970.
12. Amboseli, Kenya, Bronikowski and Altmann. Guarding constrains foraging activity of male baboons, Animal Behaviour. 1996;51:1269–1277.
13. Barton RA, Whitten A, Strum SC, Byrne R W, Simpson AJ. Habitat and Resource Availability in Baboons, Animal Behaviour. 1992;43:831-844.
14. Gaynor D. Foraging and feeding behaviour of chacma baboons in a woodland habitat, PhD thesis, University of Natal; 1994.
15. Johnson SE, Bock J. Tradeoffs in skill acquisition and time allocation among juvenile chacma baboons. Human Nature Volume. 2004;15(1):45-62.
16. Bronikowski AM, Altmann J. Foraging in variable environment: Weather patterns and the behavioural ecology of baboons. Behaviour Ecology Sociobiology. 1996;39: 11-25.
17. Post DG. Feeding Behaviour of Yellow Baboons (*Papio cynocephalus*) In Amboseli National Park, Kenya. International Journal of Primatology. 1982; 3(4):403-430.
18. Clutton-Brock THC. Some aspect of intra-specific variation in feeding and ranging behavior in primates. In: Primate Ecology (Ed. By T.H.C. Clutton-Brock), New York: Academic press. 1977;529-536.

© 2019 Yahaya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sdiarticle3.com/review-history/50793>