



Evaluation of Mulching Methods and Legume Intercropping on Weed Control Efficiency and Agronomic Traits of Pearl Millet (*Pennisetum glaucum*)

X. Catherine Sharon ^{a*}, R. Isaac Manuel ^a, K. Sharmili ^a
and Dinesh Kumar ^a

^a Division of Agronomy, School of Agricultural Sciences, Karunya Institute of Technology and Sciences, Coimbatore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i82134

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101871>

Original Research Article

Received: 10/04/2023

Accepted: 12/06/2023

Published: 16/06/2023

ABSTRACT

Aim: To evaluate the weed control efficiencies of different mulching practices and legume intercropping in transplanted pearl millet.

Study Design: Randomized Complete Block Design (RBD).

Place and Duration of Study: The field trial was conducted during the *Rabi* season of 2022 at the school of agricultural sciences in Karunya Institute of technology and sciences, Coimbatore, Tamil Nadu.

Methodology: The field trial consists of 8 treatments viz., T₁ - PE of Atrazine 0.5kg/ ha +1 Hand Weeding at 30 DAT, T₂ - Two Hand Weeding at 15 and 30 DAT, T₃ - Paddy straw mulching at 5t/ ha

*Corresponding author: E-mail: chocisharon@gmail.com;

3 DAT, T₄ - Black polythene mulching at 3 DAT, T₅ - Black silver polythene mulching at 3 DAT, T₆ - Intercropping of pulses (cowpea) (1:1), T₇ - Intercropping of pulses (cowpea) (1:1) + PE Pendimethalin 0.75 kg/ ha, T₈ - Un weeded check.

Results: Results of the experiment proves that the intercropping of cowpea in pearl millet along with the pre-emergence application of pendimethalin proves to be beneficial and advantageous in terms of returns per rupee invested for profitable crop production, rather than going for polythene mulches. Hence adoption of legume intercropping in pearl millet, along with the usage of mild herbicides like pendimethalin, will be an economically viable and environmentally sustainable weed management practice for improving the yield of pearl millet.

Keywords: Pearl millet; plastic mulching; paddy straw mulching; weed management; Pendimethalin; atrazine.

1. INTRODUCTION

Pearl millet [*Pennisetum glaucum*] commonly known as Bajra, is one of the most important cereal crops grown worldwide after rice, wheat and maize. Pearl millet ranks as the fourth most significant staple food crop. Pearl millet stands out from other cereals for its distinctive characteristics, like exceptional photosynthetic efficiency, because of its C₄ nature of carbon fixation and a remarkable capacity for high dry matter production. The crop is highly versatile, well adaptable to stress intensive environments, and highly responsive to inputs. Pearl millet serves as an excellent source of energy, protein, as well as essential vitamins and minerals. It contains about 12.4% moisture, 11.6% protein, 5% fat, 72.2% carbohydrates, 2.7% mineral matter and gives 360 calories per 100 grams. Due to its superior nutritional composition, pearl millet has been designated as "Nutri-cereals" by the Ministry of Agriculture, Government of India. Pearl millet accounts for about 50% of the total production of millets. India is the larger producer of pearl millet. Pearl millet encompasses a land area of 7.55 million hectares; with an average annual production reaching 9.22 million tonnes and productivity level of 1374 kg/ha.

Weed infestation in pearl millet is one of the major causes for low productivity. Initial slow growth with wider spacing of the crop causes severe infestation of weeds in pearl millet. The Major weeds of pearl millet are *Trianthema protulacastrum*, *Tribulus terrestris*, *Cyprus rotundus*, *Amaranthus sp*, *Echinochloa colona*, *Cynodan dactylon*, etc. Weed infestation in pearl millet drastically reduces the grain yield and stover yield of the crop up to 40% and more [1]. Weed infestation during the early stages of the crop growth is a major constrain because it severely impacts the yield and income. If the weed infestation gets addressed advertently

during the critical crop weed competition period, the reduction in crop yield can be reversed effectively.

The predominant methods of weed management in pearl millet are hand weeding, hoeing and inter-culturing. By following these mechanical measures, the benefits obtained are loosening of soil, aeration in roots, deep root penetration and conservation of moisture. However due to the limitations like nonavailability of labour during the critical stages of crop growth, high labour cost and unpredicted continuous rains often causes difficulties in managing the weeds manually. Though the use of pre-emergence chemical herbicides like atrazine and pendimethalin have proved to be economical and effective to a large extent, continuous use of single classes herbicides over a period of time opens the doors for the evolution of herbicide resistant weed species and shift in the weed flora. Under such circumstances opting for mulching practices and adopting other management practices like intercropping can prove to be an effective viable option for managing the weeds in pearl millet. Mulches contribute to weed management by affecting the germination of weed seeds, blocking weed growth, favouring the crop by conserving soil moisture and sometimes by moderating soil temperature. Organic mulching reduces the weed competition, suppress annual weed seedlings, conserve moisture and add organic matter to soil as they break down. Synthetic mulches like black polythene mulch and black silver polythene sheets blocks the light stimulus which is the major source for seed germination and reduces the evaporation losses. On the other hand, usage of inorganic mulches has its own drawback of leaving back the non-biodegradable plastic residue in the field which drastically affects the soil health and pollutes the environment. Intercropping of pulses in pearl millet can also be adopted as an effective weed

control mechanism, it not only reduce the intensity of weeds but also gives additional yield and returns [2,3] Keeping in view with the above fact, the present investigation was carried out to study the influence of conventional weed management practices of pearl millet alongside with the different mulching methods and legume intercropping practices on the weed control efficiency and agronomic traits of transplanted pearl millet.

2. MATERIALS AND METHODS

2.1 Location

The experiment was conducted at the Instructional farm, School of Agricultural Sciences, Karunya Institute of Technology and Sciences, Coimbatore which falls in the western agro zone of the Tamil Nadu. The experimental site is geographically located at 10° 56' N latitude and 76° 44' E longitude at an elevation of 474 m above mean sea level.

2.2 Season and Crop Variety

The pearl millet crop was raised under the irrigated condition during the *Rabi* season from September 2022 to December 2022. The pearl millet hybrid CO9 released by Tamil Nadu was chosen for the trial. For intercropping treatment, cowpea variety VBN3 was selected.

2.3. Experimental Design

The field experiment was conducted in randomized block design (RBD) with three replications comprising of eight treatments. The treatments taken up for the experimental trial were : T₁ - Pre Emergence (PE) application of Atrazine 0.5kg/ ha +1 Hand Weeding at 30 Days After Transplanting (DAT), T₂ - Two Hand Weeding at 15 and 30 DAT, T₃ - Paddy straw mulching @ 5t/ ha at 3 DAT, T₄ - Black polythene mulching at 3 DAT, T₅ - Black silver polythene mulching at 3 DAT, T₆- Intercropping of cowpea (1:1), T₇- Intercropping of cowpea (1:1) + PE Pendimethalin 0.75 kg/ ha, T₈ - Unweeded check.

2.4 Cultural Practices for the Crop Management

The sowing of pearl millet seeds were taken up in raised bed nurseries and the seedlings were raised in the nursery for 15 days. The main field was prepared by ploughing the soil thoroughly

with tractor drawn rotavator to fine tilth and with the formation of ridges and furrows, which were 45 cm apart. The 15 days old seedlings were transplanted in the main field along the sides of the ridges with an intra row spacing of 15 cm (45×15 cm). In case of polythene mulching treatments, the polythene sheets were spread over the respective treatment plots on an end to end basis. Holes or punchers were made in the polythene sheets at 15 cm interval for transplanting the seedlings. Simultaneously for the intercropping treatments, 1:1 additive series of planting method was taken up by sowing the cowpea in between the pearl millet rows on the other side of the ridge at a spacing of 15 cms. Nutrient management practices were done as per the standard recommended package of practices. Irrigation was given to the crop according to the requirement.

2.5 Experimental Observations

Weed observations such as weed density, weed control efficiency were recorded at 15 days after transplanting, 45 days after transplanting and at harvest stages The crop biometric observations on yield parameters viz., number of productive tillers, crop dry matter production, grain yield, stover yield and harvest index were recorded at harvest stages.

2.6 Statistical Analysis

All the data were subjected to analysis of variance (ANOVA) at P (0.05) significance using STAR (Statistical Tool For Agricultural Research) version 2.0.1.

3. RESULTS AND DISCUSSION

3.1 Effect of Mulching Methods and Legume Intercropping on Weed Density and Weed Control Efficiency in Transplanted Pearl Millet

The measurement of weed density involves quantifying the number of weed species present within a specific area and the data recorded on weed density observed in the field trial are furnished in the Table 1. The weed density exhibits variations not only across different weed species but also across various stages of crop growth. Throughout the entire crop growth stage, the average data clearly indicates that all weed control treatments exhibited significant efficacy in managing weeds when compared to the weedy

check treatment. Particularly, noteworthy is the substantial reduction in weed density or no weed was observed in the polythene mulching treatments (T_4 & T_5) at all the stages of the crop growth. As there were no weeds observed in black polythene mulching and black silver polythene mulching treatments, eventually these treatments have reflected the highest weed control efficiency of 99.9% (Fig. 1) during all the stages of the crop growth Boutagayout et al. [4], Zhao et al. [5]. This was followed by the treatment T_7 - intercropping of cowpea along with the application of pendimethalin, which exhibited a moderately reduced total weed count of 7.19, 3.37 and 3.49 per square meter at 15, 45 and harvest stages of the crop respectively and a weed control efficiency almost 90% of was observed at all the stages of the crop growth, which was statistically on par with the paddy straw mulching (T_3).

The decreased weed density and the higher weed control efficiency recorded during all the stages of the crop growth under the polythene mulching treatments can be attributed to the prevention of light or filtering the photosynthetically active radiation from the sun falling over the soil surface by the polythene sheets, because of its opaque nature, hinders the germination of weed seeds in the soil and leading to a reduction in weed population. Similar findings of achieving complete weed management, to an extent of 100% control rate by using agricultural photo selective films were also reported by Vineet and Yadav [6], Gangaiah et al. [7] and Patel et al. [8] through their research findings. The reduction of weeds when intercropped with legumes is due to the interactive effect between the pearl millet and cowpea. The dense vegetation of cowpea creates a smothering effect on weeds. Intercropping with legumes like cowpea hinder the nutrients uptake of weeds by cutting off the weed growth and thus reduces the crop weed competition which leads to reduced weed dry matter production. Usually the most of the effective herbicides of pearl millet are not suitable under legume intercropping system as they are detrimental over broad leaved vegetation, but the pre- emergence spray of pendimethalin can be effectively used to controls the weeds. Further it improves the soil fertility through atmospheric nitrogen fixation and increases the soil microbiome activities. These findings were in agreement with the study made by Aasha et al. [9], Asodewine. [10], Meena. [11].

3.2 Effect of Mulching Methods and Legume Intercropping on the Yield Attributes of Transplanted Pearl Millet

The data on the yield attributes and the yield of pearl millet as influenced by the mulching methods and the legume intercropping are presented in Table 2. It is evident from the results that the inorganic mulching with black polythene sheets (T_4) significantly influenced the agronomic traits of pearl millet by registering a higher values of productive tillers (4.11 tillers / plant), earhead length (29.42cm), earhead girth (13.94cm), test weight (13.97g), grain yield (3553.11 kg/ha), stover yield (6473.05 kg/ha) and harvest index (35%). This was statistically on par with black silver polythene mulching treatment (T_5) This was followed by the treatment T_7 - intercropping of cowpea along with the application of pendimethalin, which was statistically on par with the paddy straw mulching treatment (T_3). The unweeded check plot (T_8) register the lower values of yield parameters among all the treatments.

The increased in yield under polythene mulching treatments T_4 and T_5 This might be due to the reason that the inorganic polythene mulching materials could have created a favourable micro environment that would have enhanced the plant growth and development by providing perfect weed control, conserving soil moisture, and maintaining stable soil temperatures. The enhanced grain yield and stover yield observed under polythene mulch could be also attributed to the elevation in soil temperature, which could have consequently facilitated an accelerated crop development and biomass accumulation with the increased photosynthetic activity, and efficient translocation of photosynthates from source to sink. This is in line with the findings of Shamla et al. [12], Timsina et al. [13] and Tiwari et al. [14] who also reported similar results from their research work. The similar findings were found in Chanu et al. [15].

3.3 Effect of Mulching Methods and Legume Intercropping on the Economics of Transplanted Pearl Millet

A non-statistical data on the economics of mulching methods and legume intercropping of transplanted pearl millet is presented in the Fig. 2.

Table 1. Weed density in response to weed management practices (No. m⁻²)

Treatment	15 DAT				45 DAT				Harvest			
	Grasses	Sedges	Broad- leaf weeds	Total	Grasses	Sedges	Broad- leaf weeds	Total	Grasses	Sedges	Broad- Leaf weeds	Total
T1- PE of Atrazine 0.5kg/ha on 3 DAT+ 1 HW AT 30 DAT	2.33 (4.92)	2.41 (5.33)	1.77 (1.46)	3.49 (11.71)	1.95 (3.31)	2.37 (5.10)	2.14 (4.08)	3.60 (12.49)	1.96 (3.34)	2.20 (4.35)	2.10 (3.92)	3.48 (11.60)
T2- Two hand weeding at 15 DAT and 30 DAT	3.54 (12.01)	3.39 (10.98)	2.76 (7.12)	5.53 (30.11)	2.20 (4.32)	2.45 (5.51)	2.18 (4.25)	3.82 (14.08)	2.21 (4.39)	2.47 (5.59)	2.26 (4.62)	3.89 (14.60)
T3- Paddy straw mulching at 5t/ha at 3 DAT	1.86 (2.97)	1.96 (3.33)	1.33 (1.27)	2.84 (7.55)	1.40 (1.46)	1.64 (2.18)	1.45 (1.60)	2.40 (5.24)	1.39 (1.42)	1.62 (2.12)	1.58 (2.01)	2.46 (5.53)
T4- Black polythene mulching	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)
T5- Black silver polythene mulching	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)
T6- Intercropping of pulses (Cowpea) (1:1)	2.93 (8.07)	2.99 (8.45)	2.37 (5.75)	4.77 (22.27)	2.92 (8.01)	3.11 (9.19)	2.65 (6.52)	4.92 (23.73)	2.76 (7.10)	3.23 (9.93)	2.67 (6.63)	4.91 (23.65)
T7- Intercropping of pulses (Cowpea) (1:1) + PE pendimethalin 0.75 kg/ha	1.81 (2.79)	1.93 (3.21)	1.30 (1.19)	2.77 (7.19)	1.25 (1.06)	1.32 (1.23)	1.26 (1.09)	1.97 (3.37)	1.27 (1.10)	1.35 (1.31)	1.26 (1.08)	2.00 (3.49)
T8- Unweeded check	3.58 (12.32)	3.45 (11.38)	2.80 (7.34)	5.62 (31.04)	3.85 (14.32)	3.66 (12.89)	3.13 (9.30)	6.08 (36.50)	4.11 (16.42)	3.89 (14.67)	3.45 (11.42)	6.56 (42.51)
SE(d)	0.29	0.21	0.08	0.13	0.22	0.2	0.17	0.37	0.23	0.2	0.18	0.39
CD (5%)	0.59	0.45	0.17	0.27	0.48	0.42	0.37	0.8	0.5	0.42	0.38	0.84

Data are root transformation values of $x + 0.5$
Data in the parenthesis are raw values

Table 2. No. of productive tillers, earhead length, earhead girth, test weight, grain yield and straw yield in response to weed management practices

Treatment Details	Number of productive tillers / Plant	Earhead length (cm)	Earhead girth (cm)	Test weight (g)	Grain yield (kg/ha)	Straw yield (kg/ha)	Harvest index (%)
T1- PE of Atrazine 0.5kg/ha on 3 DAT+ 1 HW AT 30 DAT	2.69	21.41	9.07	13.83	2629.33	5255.51	33
T2- Two hand weeding at 15 DAT and 30 DAT	2.67	20.46	9.01	13.79	2438.51	5171.72	32
T3- Paddy straw mulching at 5t/ha at 3 DAT	3.29	24.92	10.75	13.85	3025.67	5805.45	34
T4- Black polythene mulching	4.11	29.42	13.94	13.97	3553.11	6473.05	35
T5- Black silver polythene mulching	3.90	29.13	13.58	13.94	3465.00	6371.65	35
T6- Intercropping of pulses (Cowpea) (1:1)	2.17	17.72	8.05	13.76	1987.64	4624.80	30
T7- Intercropping of pulses (Cowpea) (1:1) + PE pendimethalin 0.75 kg/ha	3.32	25.32	10.86	13.87	3062.17	5821.42	34
T8- Unweeded check	1.63	14.23	7.08	13.64	1537.21	4098.63	27
SE(d)	0.22	1.10	0.41	1.21	182.72	77.93	-
CD	0.46	2.36	0.88	2.60	391.89	167.14	-




Fig. 1. Weed control efficiency (%) in response to weed management practices

The treatments are T₁ - PE of Atrazine 0.5kg/ ha +1 Hand Weeding at 30 DAT, T₂- Two Hand Weeding at 15 and 30 DAT, T₃- Paddy straw mulching at 5t/ ha 3 DAT, T₄- Black polythene mulching at 3 DAT, T₅- Black silver polythene mulching at 3 DAT, T₆- Intercropping of pulses (cowpea) (1:1), T₇- Intercropping of pulses (cowpea) (1:1) + PE Pendimethalin 0.75 kg/ ha, T₈- Un- weeded check




Fig. 2. Economics of pearl millet in response to weed management practices

The treatments are T₁ - PE of Atrazine 0.5kg/ ha +1 Hand Weeding at 30 DAT, T₂- Two Hand Weeding at 15 and 30 DAT, T₃- Paddy straw mulching at 5t/ ha 3 DAT, T₄- Black polythene mulching at 3 DAT, T₅- Black silver polythene mulching at 3 DAT, T₆- Intercropping of pulses (cowpea) (1:1), T₇- Intercropping of pulses (cowpea) (1:1) + PE Pendimethalin 0.75 kg/ ha, T₈- Un- weeded check

It is evident from the computation of economics that the higher cost of cultivation per hectare was observed in the treatment T₄ - black polythene mulching (₹ 1, 05,690 ha⁻¹), but the benefit cost ratio of polythene mulching treatment was very low to an extent of almost comparable with the unweeded control treatment T₈. On the other hand, the treatment T₇ - intercropping of Cowpea along with Pre-emergence application of Pendimethalin 0.75 kg ha⁻¹, registered a higher benefit cost ratio of 2.3. From these result it could be inferred that though the polythene mulches are effective in controlling the weeds and improving yields, they are not economical.

4. CONCLUSION

Based on the results, it can be concluded that the intercropping of cowpea in pearl millet along with the pre-emergence application of pendimethalin performs better in terms of profitable crop production. Though the polythene mulches are 99.9% effective in controlling weeds and enhancing yield of the crop, at the moment they are not cost effective for a profitable crop production and it has its own drawback of leaving back the non-biodegradable plastic residue in the field which drastically affects the soil health and pollutes the environment. Hence adoption of legume intercropping in pearl millet, along with the usage of mild herbicides like pendimethalin, will be an economically viable and environmentally sustainable weed management practice for improving the yield of pearl millet when compared with the practice of polythene mulching.

ACKNOWLEDGEMENT

The authors are grateful to the division of Agronomy, School of Agricultural sciences (SAS), Karunya Institute of Technology and Sciences (KITS), Coimbatore, Tamil Nadu – 641114.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Girase PP, Suryawanshi RT, Pawar PP, Wadile SC. Integrated weed management in pearl millet. Indian journal of weed science. 2017;41.
2. Vishalini R, Rajakumar D, Joseph M, Gomathy M. Efficient non chemical weed management strategy for irrigated finer millet (*Eleusine coracana* L.). Journal of Pharmacognosy and Phytochemistry. 2020;9(5):1210-1212.
3. Chaudhary C, Dahiya S, Rani S, Pandey S. Review and outlook of weed management in Pearl millet. IJCS. 2018;6(2):2346-2350.
4. Boutagayout A, Nassiri L, Bouiamrine E H, Belmalha S. Mulching effect on weed control and faba bean (*Vicia faba* L. Minor) yield in Meknes region, Morocco. In E3S web of conferences; 2020.
5. Zhao Hubing, Liu Jifei, Chen Xiuwen, Wang Zhaohui. Straw mulch as an alternative to plastic film mulch: Positive evidence from dryland wheat production on the Loess Plateau. Sci Environ. 2019; 676(3).
6. Dheer V and Yadav RA. Comparative efficacy of paddy straw and polythene sheet mulching on weed infestation in wheat (*Triticum aestivum* L.). The Pharma Innovation Journal. 2021;10(9): 1516-1519.
7. Gangaiah B, Babu MBB, Latha PC, Singh TV, Rao PR. Rice cultivation using plastic mulch under saturated moisture regime and its implications on weed management, water saving, productivity and profitability. Indian Journal of Weed Science. 2019; 198-202. Available:<http://dx.doi.org/10.5958/0974-8164.2019.00041.8>
8. Patel HF, Attar SK, Makwana AI, Bana JK. Effect of mulching and herbicides on weeds, yield and economics of tomato grown under drip irrigation system. Indian Journal of Weed Science. 2021;195-197.
9. Aasha MB, Amani EH, Ras H, Abusin MA, Khalil NA. Effects of intercropping pearl millet with some legumes on Striga Hermonthica emergence. SSRG Int. J. Agric. Environ. Sci. 2017;4:64-72.
10. Asodewin PA, Lamptey S, Dzomeku IK. Pearl millet-cowpea intercrop effect on Striga hermonthica and grain yield; 2021.
11. Meena R, Kaur A, Nagar R, Nagar D. Studies on weed competition in intercropping system of pearl millet (*Pennisetum Glaucum* L.) with legumes as fodder; 2022.
12. Shamlia K, Sindhu PV, Menon MV. Effect of weed management practices on growth and yield of okra (*Abelmoschus esculentus*

- (L.) Moench.). Journal of Tropical Agriculture. 2017;55(1):57-62.
13. Timsina D, Marahattha S, Sah SK, Adhikari JB. Evaluation of different types of mulching practices on weed management and productivity of winter maize in Chitwan, Nepal. Journal of Research in Weed Science. 2019;2(1):65-77.
 14. Tiwari DK, Gangwar SK, Singh RP, Mishra PK, Singh RP. On-field assessment of mulching materials for weed control and its impact on yield and economics in okra (*Abelmoschus esculentus* (L.) Moench). Indian Journal of Extension Education. 2021;57(2): 157-161.
 15. Chanu NB, Kanaujia S, Changkiri M, Singh A. Response of weed management practices in okra (*Abelmoschus esculentus*) under foothill condition of Nagaland. Annals of Plant and Soil Research. 2018;20(4):324-329.

© 2023 Sharon et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/101871>