

## Asian Journal of Case Reports in Surgery

2(1): 13-16, 2019; Article no.AJCRS.46202

# Accidental Ingestion of Spoon Presenting as Gastric Outlet Obstruction – An Unusual Case Report

Sumit Pathania<sup>1</sup>, Ashish Vora<sup>1\*</sup>, Rachan Kathpal<sup>1</sup>, Dipankar Naskar<sup>1</sup>, Nikhil Gupta<sup>1</sup> and C. K. Durga<sup>1</sup>

<sup>1</sup>PGIMER & Dr R.M.I. Hospital, BKS Marg, New Delhi-110001, India.

#### Authors' contributions

This work was carried out in collaboration between all authors. Author AV designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors SP and RK managed the analyses of the study. Author RK managed the literature searches.

All authors read and approved the final manuscript.

#### Article Information

DOI: 10.9734/AJCRS/2019/46202

Editor(s)

(1) Dr. Georgios Tsoulfas, Associate Professor, Aristoteleion University of Thessaloniki, Thessaloniki,

Greece. Reviewers:

(1) Nagahito Saito, Nemuro City Hospital, Japan.

(2) Einar Arnbjörnsson, Lund University, Sweden.

Complete Peer review History: http://www.sciencedomain.org/review-history/28178

Case Study

Received 09 October 2018 Accepted 21 December 2018 Published 09 January 2019

## **ABSTRACT**

We report a case of a 60 year old male, who presented to our emergency with history of accidental spoon ingestion causing gastric outlet obstruction. Patient had undergone endoscopic extraction which was unsuccessful. He underwent exploratory laparotomy to retrive the spoon and was discharged on 8<sup>th</sup> postoperative day.

Keywords: Spoon; gastric outlet obstruction; exploratory laparotomy.

#### 1. CASE

A 60 year old male presented to our emergency with complaints of pain abdomen associated with large amount of non-bilious foul smelling

vomiting since 20 days following accidental ingestion of spoon while cleaning his tongue with it, he also gave history of failed endoscopic extraction of the spoon in some other hospital. There was no significant medical or surgical but

patient had psychiatric history. In spite of recurrent vomiting patient was taking food in order to push the spoon distally.

On examination, patient was dehydrated with pulse rate of 110 bpm and blood pressure of 100/70 mmHg. On abdominal examination patient had tender fullness over the epigastrium tympanic note on percussion succussion splash on auscultation. Arterial Blood Gas showed hypochloraemic, hypokalaemic metabolic alkalosis, rest investigations were unremarkable. A naso-gastric tube was insterted and large amount of foul smelling undigested food particles were aspirated and, after which, a X-ray erect abdomen was done (Fig. 1) that showed right paravertebral radio-opaque shadow of a spoon. ULTRA SONOGRAPHY whole abdomen showed intraluminal foreign body lodged partly inside distal stomach and proximal duodenum. Based on the history, presentation and investigations a diagnosis of gastric outlet obstruction secondary to accidental spoon ingestion was made and patient was shifted to operation theatre. An emergency exploratory laparotomy was done and with a juxta-pyloric gastrotomy, a large amount of undigested foul smelling food was removed. the spoon was



Fig 1. X-ray erect abdomen showing right paravertebral radio-opaque shadow of spoon.

found to be tightly occluding the pylorus and proximal duodenum with the pylorus and first and second part of duodenum falling in one straight line with food particles cementing the lodgement. The 14.5 cms spoon was maneuvered with due precautions to minimize futher trauma to stomach and duodenum (Fig. 2). Gastrotomy was closed in two layers after through wash and placement of naso-gastric tube in-situ. A 28 French intra-abdominal drain was placed and abomen was closed in layers. Post operative period was uneventful and the patient was discharged on post-operative day 8 on full diet. The patient was followed for next 3 months which were uneventful.

## 2. DISCUSSION

Most of the cases of foreign body ingestion that present to the emergency are in the paediatric age group, whereas in healthy adults this is rare and the patients are either psychiatric or under the influence of alcohol [1].

The impaction of foreign body is also determined by its dimensions, any foreign body longer than 6 cms and wider than 2 cms are less likely to pass spontaneously [2].



Figure 2. 14.5 cms spoon after extraction.

Table 1. Anatomical sites of natural constrictions along upper gastrointestinal tract

| SI. no. | Site                                    | Distance from incisors (in cms) |
|---------|---|---------------------------------|
| 1       | Pharyngo-oesophageal junction           | 9                               |
| 2       | Crossing of the arch of aorta           | 22.5                            |
| 3       | Crossing of the left principal bronchus | 27.5                            |
| 4       | Level of the diaphragm                  | 40                              |

Types of foreign bodies can be Radio-opaque viz. coins, spoon, fish bones, magnets, etc. or Radiolucent viz. food bolous, trichobezoars, phytobezoars, etc. [3,4]. Body packing refers to smuggling of drugs by swollowing it in packets which could obstruct or burst causing toxicity.

Any foriegn body that has been intentionally or accidentally ingested might lodge at few anatomical narrowings of the oesophagus.

Out of total about 80 % to 90% of foreign bodies do not need any intervention and can be managed conservatively, arround 10% to 20% might need Endoscopic intervention and less than 1% will need surgery in form of Exploratory Laparotomy [5,6,7,8]. Laparoscopic or Laparoscopic assisted romoval of the foreign body can be attempted to avoid large laparotomy incision in young patients [6,8].

Diagnosis mainly is by history, which is supported by investigations like CT abdomen (sensitivity of 100% and specificity of 91%) and X-ray and ULTRA SONOGRAPHY whole abdomen, out of which latter is of less diagnostic value [9].

Complications due foreign body ingestions can be:

- 1. Obstruction.
- 2. Haemorrhage.
- Perforation (especially if the foreign body is sharp), leading to peritonitis or mediastinitis.
- 4. Pressure necrosis at the site of impaction.

## 3. CONCLUSION

Foreign body ingestion and further obstruction is commonly encountered in emergency department. The main idea is to tailor the treatment according to type and size of the foreign body and site of obstruction.

#### CONSENT

As per international standard or university standard, patient's consent has been collected and preserved by the authors.

## **ETHICAL APPROVAL**

It is not applicable.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### **REFERENCES**

- Pfau PR. Removal and management of Esophageal foreign bodies. Techniques in Gastrointestinal Endoscopy. 2014;16:32-20
  - DOI:https://doi.org/10.1016/j.tgie.2013.10.004
- Munter DW, Dronen SC. Medscape (Internet) 2012. May 24, [Last cited on 2013 Aug 25].
  - Available:http://emedicine.medscape.com/article/776566-clinical
- Kay M, Wyllie R. Pediatric foreign bodies and their management. Curr Gastroenterol Rep. 2005;7(3):212–218. Crossref, Medline, Google Scholar
- Rothmann BF, Boeckman CR. Foreign bodies in the larynx and tracheobronchial tree in children: A review of 225 cases. Ann Otol Rhinol Laryngol. 1980;89(5 Pt 1):434–436. Crossref, Medline, Google Scholar
- Dalal PP, Otey AJ, McGonagle EA, Whitmill ML, Levine EJ, McKimmie RL, et al. Intentional foreign object ingestions: Need for endoscopy and surgery. Journal of Surgical Research. 2013;184:145-149. DOI:https://doi.org/10.1016/j.jss.2013.04.0 78.

- 6. Gajendran M, Muniraj T, Gelrud A. A challenging case of gastric outlet obstruction (Bouveret's Syndrome): A case report. Journal of Medical Case Reports. 2011;5:497-401.
- 7. Yao CC, Wu IT, Lu LS, Lin SC, Liang CM, Kuo YH, et al. Endoscopic management of foreign bodies in the upper gastrointestinal tract of adults. BioMed Research International. 15, 658602; 2015.
- 8. Beldholm BR, Lee AU. Simple endoscopic technique for retrieving a long foreign body from the stomach. ANZ J Surg. 2007;77:560–561. [PubMed].
- 9. Marco De Lucas E, Sadaba P, Lastra Garcia-Baron P, et al. Value of helical computed tomography in the management of upper esophageal foreign bodies. Acta Radiol. 2004;45:369–374. [PubMed].

© 2019 Pathania et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
http://www.sciencedomain.org/review-history/28178