



Effects of Addition of Dried Sweet Potato Peels Powder on the Quality Characteristics of Cupcakes

Mst. Sorifa Akter^{a*}, Md. Asaduzzaman^a and Mahbub Rahman^a

^a *Department of Food Engineering and Technology, State University of Bangladesh, Dhanmondi, Dhaka 1205, Bangladesh.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2022/v37i1030535

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86165>

Original Research Article

Received 20 February 2022
Accepted 23 April 2022
Published 19 September 2022

ABSTRACT

The study was conducted to develop baked cakes using different pretreated sweet potato peels powders (such as fresh, boiled and citric acid) with various concentration (5%, 10% & 20%) and to evaluate the changes of quality characteristics. Sweet potato peel powder was investigated for the different nutritional and physiochemical properties were boiling and pretreatment effects also evaluated. Fresh peel, 0.5% citric acid treated and boiled sweet potato peel powder was utilized as sample powder for the studies. There were significant differences ($p < 0.05$) in fiber and ash content and Hunter color values between powders from fresh peeled, 0.5% citric acid treated and boiled sweet potato peels powders. However, fresh and boiled sweet potato peel powder had higher fiber content than treated. The highest amount of fiber content was found in cakes incorporated with 10% boiled treated sweet potato peels powders content highest fiber. Pretreated and boiled with citric acid samples were obtained overall acceptability in the hedonic rating test.

Keywords: Sweet potato; peels; baked cake.

1. INTRODUCTION

Sweet potato (*Ipomoea batatas*) is one of the major root crops which owing to present of an ample amount of numerous health-beneficial compounds. It has high nutritional value and a vital source of carbohydrates, protein, vitamins, and minerals [1]. It is ranked the seventh most important food crop around the world which fulfilled the highest amount of edible energy [2]. Due to limited production cost, it was widely cultivated in most developing countries [3]. In Bangladesh 60000 acreage of land is under sweet potato cultivation within an annual production of 236000 metric tons in 2018-2019 where 59000 acreage of land is under sweet potato cultivation within an annual production of 247000 metric tons in 2017-2018 [4]. Due to higher production rate and lower price sweet potato flour can be made and used as wheat flour substitute all the year round [1]. Generally, more than fifty percent of the sweet potato produced is consumed boiled and fried and the rest is used for the processing of different processed products. The processed products are bread, crisps, puff-puff, buns, cake, chips, jam, starch, chin-chin, cookies and pasta, formulated infant foods, puree, fermented lacto juice, and lacto pickles, fries [5-6]. Nowadays people are searching about new, low cost and healthful food to adjust their changed lifestyles. Therefore, the new taste of lifestyles also reflects on their food production as well as food consumption. Lifestyle has changed along the way, from food production to food consumption.

Day by day baked goods are attracted considerable attention due to their excellent taste, ease of preparation, and nutritional value. Cake is a worldwide known baked item prepared from the mixture of different flours, butter, sugar, oil, egg. Previously it was found that incorporation of dried fruits and vegetable powder in cakes can contributed additional nutritional values especially antioxidants, vitamins, minerals and fiber content as well as improved the textural and sensorial properties of cakes [7]. Nowadays it is transformed in different shaded of tastes as the ingredients are slightly modified. For example, whole grain, fiber fortified, less salt, gluten free baked items. As a result, valuable fruits and vegetables by products are incorporated into baked items such as cookies and muffins to reduce carbohydrate intake as well as increase fiber content [8]. In developing countries, wheat flours are partially combined with flours made from other locally

grown crops as composite flours as a substitute for wheat flour or to encourage the use/application of locally grown crops. Composite flours actually improve the nutritional quality of the wheat flour. Previously it was shown that composite flours are attracting much attention for the preparation of baked goods [9].

Every year the huge quantity of sweet potatoes is grown in Bangladesh. During peak season its production is very high in Bangladesh. In this connection, huge amount of sweet potato peels is discarded as wasted. Usually, sweet potato peels are not degraded into the soil. Therefore, these peels could be cause environmental pollution. Usually, sweet potato is used to make various baked products as already said but there are no products prepared from sweet potato peels. Although it was proven that sweet potato peel flour contains higher amount of nutrients as compared to their flesh flour [10]. Therefore, Sweet potato peel flour may be used in the preparation of cake making.

After fruits and vegetables processing there are several waste materials are generating such as seeds, peels, leaves, roots, and stems which can be utilized as by-products. These by products have lots of nutritional value and inexpensive. Fruit peels powders have high bioactive compounds and functional properties for food formulations [11]. There are various fruits and vegetables by-products have been identified for being used in the production of baked cupcakes such as cakes from apple pomace, pea hulls, pineapple waste [12-14]. It has already been established those sweet potatoes are rich in vitamin A and other bioactive compounds. Although it was proven that sweet potato peel had higher nutritional compositions as compared to its flesh [10]. There is no formulated food using sweet potato peel yet in Bangladesh. So, Sweet potato peel could be used as health booster due to source of different health beneficial compounds. In Bangladesh, people are very fond of sweet potato as boiled or fried and discarded a huge amount of peel as waste. However, sweet potato peel products are not available in local market in Bangladesh. Therefore, present research will open new era in food sector as well as reduce environment pollution or utilization of by-products in Bangladesh. Therefore, the aims of the present study to develop value-added cakes formulated by incorporating sweet potato peels at different ratios with wheat flour and to find out optimal condition of the cake and analyze various

nutritional, physicochemical properties and sensory evaluation of backed cakes.

2. MATERIALS AND METHODS

2.1 Raw Materials, Chemicals and Other Ingredients

Sweet potatoes (*Ipomoea batatas*) were collected from a local supplier at Dhaka in Bangladesh and selected on the basis of uniform size and shape, good color, free from insects, and without any mechanical damage. The packaged wheat flour, oil, and sugar of "Pushti" brand, skimmed milk powder of "diploma" brand, and butter of "Akij Food and beverage" brand, eggs, baking powder, salt, powder milk, margarine and vanilla essence were collected from the local market. Citric acids and packaging materials were also collected from local company. All materials were kept at room temperature except butter and eggs which were stored in refrigerator. High density polyethylene bags were used for storage of peel powders and cakes samples.

2.2 Preparation of Sweet Potato Peels Powder

In this research, we utilized sweet potato peels in three groups where the first group, a sharp stainless-steel knife utilizes to peeled the fresh sweet potato and the peels collected in a fresh bowl. Then peels cut into small pieces and dried by an oven at $55\pm 5^{\circ}\text{C}$ until moisture content reach below 10%. Then the dried peels were blended by blender machine and sieved through 50 mesh size to obtain a fine powder. For the second group, according to Pao and Petracek, 1997, after collecting the peels by knife, peels were dipped in citric acid aqueous solution (0.5% w/v) at ambient temperature for 2 min [15]. After that drain out excess water and then dried and ground to make peel powder same as before. For the third group, peels were collected from boiled sweet potatoes and dipped in dipped in citric acid aqueous solution (0.5% w/v) at ambient temperature for 2 min, dried and ground to make treated boiled peel powders.

2.3 Formulations of Cake from Sweet Potato Peels Powders

The basic formulations of developed cakes incorporated with sweet potato peels powders using different treatments and ratios are outlined in Table 1. Sweet potato peels powders were

incorporated into cakes in the following proportions: Sample S_1 = Control (Market Flour); Sample S_2 = 05% Fresh sweet potato peel powder; Sample S_3 = 05% Treated sweet potato peel powder (0.5% citric acid); Sample S_4 = 05% Boiled Treated sweet potato peel powder (0.5% citric acid); Sample S_5 = 10% Fresh sweet potato peel powder; Sample S_6 = 10% Treated sweet potato peel powder (0.5% citric acid); Sample S_7 = 10% Boiled Treated sweet potato peel powder (0.5% citric acid); Sample S_8 = 20% Fresh sweet potato peel powder; Sample S_9 = 20% Treated sweet potato peel powder (0.5% citric acid); Sample S_{10} = 20% Boiled Treated sweet potato peel powder (0.5% citric acid). At first, grounded the sugars to make them fine powders. Then it was properly mixed with butter and dalda mixed. The eggs were beaten for 5 minutes with the hand bitter. Dried ingredients such as wheat flour, baking powder, salt, powdered milk and peels powder were mixed properly and then mixed to the butter to get soft mixer. At last vanilla essence was added. Then it was poured batter into cake mould box and baked in an oven at 170°C for 17 min. Then it was cooled to room temperature and was packaged with high density polyethylene bags and stored in room temperature for further analysis.

2.4 Proximate Composition

Association of Official Agricultural Chemists (AOAC) official methods are utilized as determining the proximate analysis for sweet potato peels powders and baked cakes [16]. Carbohydrate was determined by the difference from other nutrition such as fat, moisture, ash, and protein. The caloric value was calculated to conversion factors of four kcal per gram for proteins and carbohydrates, and nine kcal per gram for fats [17]. All the analyses were done in triplicates.

2.5 Determination of Crude Fiber

The crude fiber contents of different sweet potato peel powders and cakes were determined by the method of AOAC Method: 991.43 [18].

2.6 Determination of Total Polyphenol Content

Determination of the total polyphenol content was done by Ahmed *et al.*, (2010) and Saikia *et al.* (2012) method with some modifications where absorbance at 765 nm was measured using a spectrophotometer (T80 U/VIS, United Kingdom)

[19,20]. Total phenols were calculated based on standard curves of gallic acid and expressed as milligram per gram.

2.7 Cake Specific Volume

Cake-specific volume determined by according to method described by AACC (2000) with adaptations to the different conditions of the product [21]. A slide calipers utilized to determine the shape dimensions of the cakes.

2.8 Hunter Color Values

A spectrophotometer (CM-3500d, Minolta, Japan) was utilized to determine the hunter values such as L*, a*, and b* which are the color attributes of a product. Color change was calculated utilizing the following equation.

$$\Delta E = [(\Delta L^*)^2 + (\Delta a^*)^2 + (\Delta b^*)^2]^{1/2}$$

2.9 Sensory Evaluation

Sensory evaluations of the entire sample baked cake were done by taste testing panel utilizing hedonic rating test where ten test panelists present. They were asked to evaluate different sensory parameters and scoring rate on a nine-point hedonic scale where 09= Like extremely, 08= Like very much, 07= Like moderately, 06= like slightly, 05 = neither like nor dislike, 04= Dislike slightly, 03= Dislike moderately, 02 = Dislike very much and 01= Dislike extremely. The preference differences were evaluated by statistical analysis of the data for variance and

consequently, Duncan's Multiple Range Test (DMRT). Procedures of the Statistical Analysis System (SAS, 1985) were used for statistical analysis.

2.10 Statistical Analysis

The results were expressed as the mean, standard error of the mean (SEM), and coefficient of variation of each species for each parameter was determined. Data were statistically analyzed (R statistical software 3.4.1) by one-way analysis of variance (ANOVA). Mean comparisons were performed using DMRT for significant effect at $P < 0.05$.

3. RESULTS AND DISCUSSION

3.1 Physical and Chemical Properties

Table 2 represented the values of proximate analysis of control (market wheat flour) and different sweet potato peels powders such as fresh, treated, and boiled. The moisture content of all peel powders fluctuated from 7.30 % to 7.96% while the moisture content of market wheat flour was 7.94%. The extent of drying and compositional differences might be reasons for the differences of peels powders with the market wheat flour. The protein content of all peel's powders ranged from 6.77% to 9.45% protein and 74.88% to 80.66% carbohydrate whereas protein content and carbohydrate of wheat flour was 13.70% and 70.51% respectively. All sweet potato peels powders contained lesser protein content and higher amount of superior carbohydrate than wheat flour.

Table 1. Percentage compositions of control and sweet potato peels powders incorporated cakes

Ingredients	Sample			
	Control	05%	10%	20%
Sweet potato peels powders (gram)	0	1.10	2.20	4.40
Flour (gram)	22	20.90	19.80	17.60
Sugar (gram)	25.00	25.00	25.00	25.00
Egg (gram)	25.63	25.63	25.63	25.63
Fat (gram)	25.00	25.00	25.00	25.00
Milk Powder (gram)	1.50	1.50	1.50	1.50
Baking Powder (gram)	0.50	0.50	0.50	0.50
Salt (gram)	0.10	0.10	0.10	0.10
Vanilla essence (gram)	0.27	0.27	0.27	0.27
Total	100	100	100	100

Table 2. Proximate compositions and total polyphenol content of the peel's powders

Powder sample	Moisture content (%)	Protein content (%)	Fat Content (%)	Ash content (%)	Carbohydrate Content (%)	Total crude fiber content (%)	Total Polyphenol content (mg gallic acid equivalent/1g)
Control (Market Flour)	7.94±0.01 ^a	13.70±0.01 ^a	1.87±0.01 ^a	5.98±0.01 ^b	70.51±0.01 ^a	1.15±0.01 ^b	0 ^d
Fresh sweet potato peel powder	7.86±0.01 ^a	6.77±0.01 ^b	1.71±0.01 ^a	6.86±0.01 ^a	76.86±0.01 ^a	13.64±0.39 ^a	14.42±0.02 ^a
Treated sweet potato peel powder (0.5% citric acid)	4.96±0.01 ^b	7.02±0.02 ^b	1.23±0.01 ^b	6.13±0.01 ^a	80.66±0.01 ^a	4.86±0.06 ^a	10.03±0.01 ^b
Boiled Treated sweet potato peel powder (0.5% citric acid)	7.30±0.01 ^a	9.45±0.01 ^a	1.48±0.01 ^{ab}	6.89±0.01 ^a	74.88±0.01 ^a	8.28±0.05 ^a	8.35±0.01 ^c

All values are express as mean ± SD.

Mean followed by different superscript letters in each column are significantly different (p<0.05)

From the Table 2 we also found the fat content of all peel's powders ranged from 1.23% to 1.71% which was slightly lower than that of wheat flour (1.87%). Maruf, Sorifa, and Jong-Bang (2010) mentioned 0.59% to 1.29% fat content in peeled and unpeeled sweet potato powder [22]. This variation in fat content might be due to the varietal difference, peel content, fat extraction by different chemicals and different milling procedure and machines. The ash content of peels powders ranged from 6.13% to 6.89% whereas wheat flour was 5.98%. The ash contents of all peel's powders significantly higher than market wheat flour. Different milling machines and environment may be responsible for this fluctuation.

3.2 Total Crude Fiber Content

Table 2 represents the total crude fiber content of peels powders which ranged from 4.86% to 13.65% where as the crude fiber of wheat flour was 1.15%. The total crude fiber content of fresh peel had the highest value (13.65%) as compared to that of citric acid treated and boiled peels powders. This might be due to the loss of components which molecular weight comparatively lower than other. The lower molecular weight molecules such as sugars, vitamins, minerals and soluble dietary fiber loss during processing of treated and boiled peels powders.

3.3 Total Polyphenol Content of Peels Powders

The total polyphenol content of sweet potato peels powders ranged from 8.35 to 14.42 mg gallic acid equivalent/ 1g (Table 2). Due to peels are rich in different phenolic content, unpeeled sweet potato powder had better phenolic content than peeled flour which has similar observations were found by Maruf, Sorifa, and Jong-Bang [22]. The total polyphenol content was highest in fresh peels powders and lowest in boiled peels powders. Similar observations were obtained from persimmon peel where blanched samples had lower total phenolic content. Different oxidation such as enzymatic or chemically or water leaching responsible for this massive loss of the phenolic content [19,23].

3.4 Proximate Compositions of Baked Cakes

The data obtained from the Table 3. proximate analysis of baked cakes integrating three different concentrations such as (5%, 10%, 20%)

sweet potato peels powders with citric acid treated and fresh (untreated) samples are shown in Table 3. The moisture content of the control cake (wheat flour cake) was 22.75% whereas all cakes samples were ranged from 12.06% to 21.15%. The protein content and fat content of wheat flour cake were 8.53% and 28.83% respectively where as all peels powders cakes samples had 7.46% to 9.25% protein content and 26.59% to 30.60%. However, the values were not significantly different from each others for all samples. The ash and carbohydrate percentage of control cake was 0.85% and 39.04%, respectively and these percentages increased with the increased of sweet potato peels powders concentrations (5%, 10%, 20%) and with the different treatments. In addition, the ash content increased with increase in peel powder concentration and the higher concentrations (20%) had higher amount of ash and carbohydrate content also.

3.5 Total Polyphenol and Total Crude Fiber Content of Baked Cakes Samples

Total crude fiber contents of backed cake from sweet potato peel powders ranged from 0.51 to 7.85 g per 100 g (Table 3). Sample S3 contained highest amount of total crude fiber and sample S9 had the lowest level of protein. There were significant differences in total crude fiber contents of all samples at different backed cake which was fortified by different concentrations sweet potato peels powders.

Table 3 represents the total polyphenol content of control cake sample (wheat flour cake) was 1.13 mg gallic acid equivalent/1g. On the other hand, the total polyphenol content of all cake samples ranged from 1.53 to 4.69 mg gallic acid equivalent/1g. It was observed that the total phenolic content of cake samples was increased with increase in the concentrations of sweet potato peels powders and it was higher in cakes prepared from boiled peels powders. However, the content of polyphenol content of peels powders is much superior to any kind of baked cakes because polyphenol degrade due to different processing conditions and baking temperature.

3.6 Hunter Color Values

Hunter color was represented by L*, a*, b*, and ΔE where L* is a measure of lightness, a*

Table 3. Proximate compositions and Total Polyphenol content of the baked cake sample

Cake type	Moisture content (%)	Protein content (%)	Fat Content (%)	Ash content (%)	Carbohydrate Content (%)	Calorie value (Kcal/g)	Total crude fiber content (%)	Total Polyphenol content (mg gallic acid equivalent/1g)
Sample S ₁	22.75±0.346 ^a	8.53±0.01 ^{ab}	28.83±0.01 ^{abc}	0.85±0.0042 ^f	39.04 ^d	449.75	6.9±1.84 ^{ab}	1.13±0.02 ^e
Sample S ₂	20.26±0.111 ^{abc}	9.02±0.01 ^a	26.59±0.01 ^c	0.89±0.0002 ^{ef}	43.24 ^{bc}	448.35	3.8±0.23 ^{abc}	1.63±0.01 ^{de}
Sample S ₃	19.46±0.330 ^{abc}	9.07±0.01 ^a	27.32±0.01 ^c	0.90±0.0004 ^{cdf}	43.25 ^{bc}	455.16	7.9±0.77 ^a	1.99±0.01 ^{de}
Sample S ₄	15.97±0.018 ^{cde}	9.25±0.01 ^a	27.99±0.01 ^{abc}	0.99±0.0001 ^{cde}	45.80 ^{ab}	472.11	0.7±0.02 ^c	2.9±0.01 ^{bc}
Sample S ₅	21.15±0.015 ^{ab}	8.32±0.02 ^{abc}	29.01±0.01 ^{abc}	1.01±0.0004 ^{cde}	40.51 ^{cd}	456.41	1.3±0.96 ^{bc}	1.53±0.01 ^{de}
Sample S ₆	18.53±0.378 ^{abcd}	7.93±0.02 ^{bc}	30.28±0.02 ^{ab}	1.03±0.0002 ^{cd}	42.23 ^{bcd}	473.16	2.5±0.60 ^{abc}	2.18±0.01 ^{cd}
Sample S ₇	14.77±0.094 ^{de}	8.82±0.01 ^{ab}	30.60±0.02 ^a	1.12±0.0001 ^c	44.69 ^{ab}	489.44	6.2±1.20 ^{ab}	3.01±0.01 ^{bc}
Sample S ₈	12.06±0.010 ^e	9.05±0.02 ^a	29.24±0.01 ^{abc}	1.2676±0.0001 ^b	48.38 ^a	492.89	0.8±0.40 ^c	3.08±0.02 ^{bc}
Sample S ₉	17.87±0.100 ^{bcd}	8.73±0.02 ^{ab}	27.15±0.01 ^c	1.4271±0.0001 ^a	44.82 ^{ab}	458.56	0.5±0.34 ^c	3.57±0.02 ^b
Sample S ₁₀	16.77±0.094 ^{bcd}	7.46±0.02 ^c	26.81±0.01 ^c	1.5406±0.0001 ^a	47.42 ^a	460.81	1.3±0.33 ^{bc}	4.69±0.01 ^a

All values are express as mean ± SD.

Mean followed by different superscript letters in each column are significantly different ($p < 0.05$)

Sample S₁ = Control (Market Flour); Sample S₂ = 05% Fresh sweet potato peel powder; Sample S₃ = 05% Treated sweet potato peel powder (0.5% citric acid); Sample S₄ = 05% Boiled Treated sweet potato peel powder (0.5% citric acid); Sample S₅ = 10% Fresh sweet potato peel powder; Sample S₆ = 10% Treated sweet potato peel powder (0.5% citric acid); Sample S₇ = 10% Boiled Treated sweet potato peel powder (0.5% citric acid); Sample S₈ = 20% Fresh sweet potato peel powder; Sample S₉ = 20% Treated sweet potato peel powder (0.5% citric acid); Sample S₁₀ = 20% Boiled Treated sweet potato peel powder (0.5% citric acid)

Table 4. Hunter color values of the powder sample

Hunter color values	L*	a*	b*	ΔE
Fresh sweet potato peel powder	46.14±1.900 ^a	5.69±0.147 ^b	12.03±0.569 ^a	48.02±1.644 ^a
Treated sweet potato peel powder (0.5% citric acid)	40.11±8.331 ^a	6.833±0.722 ^b	11.64±0.758 ^a	42.34±8.188 ^a
Boiled Treated sweet potato peel powder (0.5% citric acid)	36.17±4.532 ^a	8.74±0.840 ^a	9.71±1.086 ^b	38.46±4.727 ^a

All values are express as mean ± SD.

Mean followed by different superscript letters in each column are significantly different ($p < 0.05$)

redness, b* yellowness and ΔE color difference. The Hunter color parameters L*, a*, b*, and ΔE have been widely used to describe color changes during dehydration of fruits and vegetable products. These values of sweet potato peels powders with different pretreatments such as citric acid and boiling and cakes samples prepared with various concentrations (5%, 10%, and 20%) of treated peels powders were measured (Table 4 and Table 5). Fresh peels powders had higher lightness (L* values) than treated and boiled samples. L* values were positively correlated with phenolic content. Although all the values of lightness (L* values) of peels powders were not significantly different from each other. This is due to heat exposure during drying because fresh peels need lower time as compared to treated peels. There is no significant difference in L* values of the all cakes samples made with peels powders. The hunter a* values (redness) were significantly higher in boiled treated peels powders as compared to the fresh and treated peels powders. The browning which was greater in boiled treated samples might be responsible for this. This observation was similar to that obtained previously [19,22]. However, the a* values were not significantly different from each other for the cakes samples except 20% peels powder incorporated samples. Hunter b* values (yellowness) were significantly lower in boiled treated peels powders than those of treated and untreated samples, which might be due to losses of solid content during boiling. There were no significant differences were observed for all the cakes samples. The flour colors can best be described by the change in total color difference, ΔE values. The lower ΔE values (38.46) were found in treated boiled peels powders as compared to fresh and treated peels powders. The lower ΔE values may be due to the loss of phenolics and oxidation during boiling treatment [11,19]. There were no significant

differences in ΔE values among all cakes samples.

3.7 Specific Volumes of Baked Peels Cake

Different cakes specific volumes and expansion rate of all cakes samples were estimated and the results are shown in Table 5. It was found that the addition of various concentrations (5%, 10%, and 20%) of sweet potato peels powders has affected on specific volume and expansion rate and the values were higher than control cake. In addition, it was observed that the specific volume and expansion rate were higher for the sweet potato peels powders addition of 10% concentration for all peels powders such as boiled, treated with citric acid and untreated peels powders had higher specific volume of the baked cake had higher expansion rate as compared to control (wheat flour cake). Due to high protein and low moisture content in the market flour which is highly recommended for cake manufacturing can be minimized it in a certain point by utilizing sweet potato peels powder. As a result, higher moisture content in cakes samples made from sweet potato peels powders. The cakes made from citric acid treated 10% sweet potato peels powders was evaluated the highest specific volume (0.26 inch³/g), lowest density (3.91 inch³/g) and the highest expansion rate is associated with the excellent texture and other quality attributes of the baked cakes. In addition, the specific volume of baked cakes and expansion rate were increased up to the 10% of sweet potato peels powders substitution/ incorporation with boiled treated with citric acid and untreated samples. Hence, for all samples containing 20% sweet potato peels powders had the weak/less gluten for cake formulations and therefore affecting the product cake quality especially specific volume, expansion rate and sensory qualities.

Table 5. Hunter color values, Expansion rate (%), specific volumes (inch³ g⁻¹) and density (g inch⁻³) of the rough dough and cakes samples

Cake type	L*	a*	b*	ΔE	Specific volume of the rough dough (inch ³ /g)	Specific volume of the backed cake (inch ³ /g)	Density of the rough dough (g/inch ³)	Density of the baked cake (g/inch ³)	Expansion rate (%)
Sample S ₁	36.36±5.302 ^{ab}	17.31±2.765 ^{ab}	21.40±6.895 ^a	45.70±8.422 ^a	0.02	0.13	40.76	7.55	342.77±49.97 ^b
Sample S ₂	32.64±2.305 ^{ab}	16.16±1.115 ^{abc}	17.87±2.666 ^a	40.58±3.454 ^a	0.02	0.16	70.76	6.28	437.43±74.35 ^b
Sample S ₃	37.72±1.695 ^{ab}	18.07±0.660 ^a	23.16±1.784 ^a	47.83±2.099 ^a	0.02	0.21	41.81	4.87	606.9±77.62 ^{ab}
Sample S ₄	38.73±4.000 ^{ab}	15.95±0.586 ^{abc}	22.88±3.067 ^a	47.77±4.507 ^a	0.03	0.19	35.64	5.17	485.90±34.49 ^b
Sample S ₅	36.82±1.916 ^{ab}	17.76±0.206 ^a	22.01±1.905 ^a	46.44±2.480 ^a	0.02	0.21	40.76	4.83	639.50±173.49 ^{ab}
Sample S ₆	35.84±3.782 ^{ab}	14.55±1.115 ^{bcd}	20.21±3.463 ^a	43.71±4.341 ^a	0.02	0.26	40.76	3.91	810.93±212.87 ^a
Sample S ₇	29.44±8.9172 ^b	13.41±1.988 ^{cde}	20.47±0.806 ^a	38.61±6.788 ^a	0.03	0.21	35.64	4.86	557.54±87.23 ^{ab}
Sample S ₈	40.65±4.231 ^a	12.40±1.187 ^{de}	21.53±2.529 ^a	47.67±4.795 ^a	0.02	0.16	40.76	6.32	447.4±32.14 ^b
Sample S ₉	33.68±1.025 ^{ab}	10.84±0.394 ^{ef}	17.39±1.118 ^a	39.43±1.401 ^a	0.02	0.13	40.76	7.98	364.06±98.25 ^b
Sample S ₁₀	40.53±4.120 ^a	9.19±0.055 ^f	18.07±0.596 ^a	45.33±3.886 ^a	0.03	0.17	35.64	5.95	442.74±150.02 ^b

All values are express as mean ± SD.

Mean followed by different superscript letters in each column are significantly different ($p < 0.05$)

Sample S₁ = Control (Market Flour); Sample S₂ = 05% Fresh sweet potato peel powder; Sample S₃ = 05% Treated sweet potato peel powder (0.5% citric acid); Sample S₄ = 05% Boiled Treated sweet potato peel powder (0.5% citric acid); Sample S₅ = 10% Fresh sweet potato peel powder; Sample S₆ = 10% Treated sweet potato peel powder (0.5% citric acid); Sample S₇ = 10% Boiled Treated sweet potato peel powder (0.5% citric acid); Sample S₈ = 20% Fresh sweet potato peel powder; Sample S₉ = 20% Treated sweet potato peel powder (0.5% citric acid); Sample S₁₀ = 20% Boiled Treated sweet potato peel powder (0.5% citric acid)

Table 6. Mean score for texture, aroma, color, taste and overall acceptability of backed cake

Cake type	Color	Aroma	Taste	Texture	Overall acceptability
Sample S ₁	7.8 ^a	7.2 ^a	7.2 ^a	7.2 ^b	7.8 ^a
Sample S ₂	5.6 ^d	6.6 ^c	6.8 ^c	5.4 ^e	6.4 ^d
Sample S ₃	7.2 ^c	7.2 ^a	7.0 ^b	7.0 ^c	7.2 ^c
Sample S ₄	7.6 ^b	7.2 ^a	7.2 ^a	7.8 ^a	7.8 ^a
Sample S ₅	6.6 ^c	6.8 ^c	6.4 ^c	6.4 ^d	7.4 ^b
Sample S ₆	6.8 ^c	7.2 ^a	7.0 ^b	7.2 ^b	7.4 ^b
Sample S ₇	7.2 ^c	7.0 ^b	7.2 ^a	7.2 ^b	7.4 ^b
Sample S ₈	4.4 ^d	5.8 ^d	5.4 ^d	4.2 ^d	5.0 ^d
Sample S ₉	4.4 ^d	6.6 ^c	5.2 ^d	4.4 ^d	5.2 ^d
Sample S ₁₀	4.6 ^d	6.6 ^c	5.2 ^d	4.8 ^d	5.4 ^d
LSD (p<0.05)	22.68	1.57	7.57	9.68	14.70

All values are express as mean \pm SD;

Mean followed by different superscript letters in each column are significantly different ($p < 0.05$)

Sample S₁ = Control (Market Flour); Sample S₂ = 05% Fresh sweet potato peel powder; Sample S₃ = 05% Treated sweet potato peel powder (0.5% citric acid); Sample S₄ = 05% Boiled Treated sweet potato peel powder (0.5% citric acid); Sample S₅ = 10% Fresh sweet potato peel powder; Sample S₆ = 10% Treated sweet potato peel powder (0.5% citric acid); Sample S₇ = 10% Boiled Treated sweet potato peel powder (0.5% citric acid); Sample S₈ = 20% Fresh sweet potato peel powder; Sample S₉ = 20% Treated sweet potato peel powder (0.5% citric acid); Sample S₁₀ = 20% Boiled Treated sweet potato peel powder (0.5% citric acid)

3.8 Sensory Evaluation

A panel of 10 judges tested the texture, aroma, color, Taste, and overall acceptability of baked cake made from different sweet potato powder in the various ration. The mean scores for texture, aroma, color, Taste, and overall acceptability of different types of cakes sample presented in Table 6. A two-way analysis of variance ANOVA was carried out for color preference, and results revealed that there was significance ($p < 0.05$) difference in color acceptability among the baked cakes. The results of DMRT showed that there was no significance for texture difference among the formation of S₁, S₆ and S₇ (Table 6). In the case of color preference among the sample, the sample S₄ was more acceptable than other samples. Sample S₄ secured the highest score of 7.8 and ranked as "Like very much." Sample S₁, S₆ and S₇ are rated as "Like moderately" and obtaining rating 7.2. The presence of sweet potato peels powder in baked cake was 0.5% which 05% Boiled and Treated by 0.5% citric acid. In case of aroma preference among the sample, ANOVA analysis showed that there was significance ($p < 0.05$) difference in flavor acceptability among the baked cake. From Table 6, it is seen that sample S₁, S₃, S₄ & S₆ secured the highest score 7.2 for flavor and was ranked as "Like very much" and followed by the sample S₇ obtaining score 7.0. In the case of color among the sample showed that there was significance ($p < 0.05$) difference in color, as shown in Table 6. Sample S₁ secured the highest

score of 7.8 for composition and was ranked as "Like very much." The sample S₄ scored 7.6 and listed as "Like moderately." In the case of taste preference among the sample showed that there was significance ($p < 0.05$) difference in taste, as shown in Table 6. Sample S₁ & S₇ secured the highest score of 7.2 for taste and ranked as "Like very much." The sample S₉ & S₁₀ scored 5.2 and posed the lowest score. It was apparent from the results of the ANOVA there was significance ($p < 0.05$) difference in overall acceptability of the sample tested as the calculated F value (14.70) is higher than the tabulated F value (2.98). This indicates that so far as overall acceptability is a concern, the samples were equally acceptable. It can be seen from Table 06 that the sample S₁ & S₄ is the most acceptable product receiving 7.8 out of 9.0 composed to the other sample and ranked as "Like very much." The sample S₅, S₆ & S₇ securing 7.2 and was ranked as "Like moderately." After sample S₁ sample S₄ secured the highest score for texture, aroma, color, taste and overall acceptability among all the samples and was closely followed by sample S₃. So, the sample S₄ product may be regarded as the best product.

4. CONCLUSION

The nutritional and sensory attributes can be enriched by utilizing sweet potato peels in different confectionary products which can be employing any kind of manufacturing factory in any region. In this study physicochemical,

sensory attributes and nutritional features are investigated in different conditioned and treated sweet potato peels which represent it is an excellent source of nutrition that can be utilized in the manufacturing area in different concentrations where the effect of boiling and citric acid-treated samples represents better results than control and fresh samples. From the study, we found different quantitative and qualitative advantages from market flour which is significant to ash and fiber content. In nine points scale hedonic rating test results "Good" scored 78% by different panelists. In this study, it can be concluded that sweet potato peels are an excellent source that can be utilized as an alternative nutritional source which was cheap, easy to add, and better for our healthy and nutritional life.

ACKNOWLEDGEMENT

This study was supported by the Ministry of Science and Technology, Government of the Peoples' Republic of Bangladesh Special Allocation (Grant number ES 395; 2019-2020).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Silva JS, Marques TR, Simão AA, Corrêa AD, Pinheiro AC, Silva RL. Development and chemical and sensory characterization of pumpkin seed flour-based cereal bars. *Food Science and Technology*. 2014; 34(2):346-52.
2. Loebenstein G. Control of sweet potato virus diseases. In *Advances in virus research* 2015;91:33-45. Academic Press.
3. Van Hal M. Quality of sweetpotato flour during processing and storage. *Food Reviews International*. 2000;16(1):1-37.
4. Statistics BB. Statistical yearbook of Bangladesh. Statistics Division, Ministry of Planning, Dhaka, Government of the People's Republic of Bangladesh; 2019.
5. Okorie SU, Onyeneke EN. Production and quality evaluation of baked cake from blend of sweet potatoes and wheat flour. *Academic Research International*. 2012; 3(2):171.
6. Dash SK, Rayaguru K. Post-harvest processing and utilization of sweet potato: A review. *Food Reviews International*, 2019;35(8):726-762.
7. Salehi F, Aghajanzadeh S. Effect of dried fruits and vegetables powder on cakes quality: A review. *Trends in Food Science & Technology*. 2020;95:162-72.
8. Taiyeba N, Gupta A, Verma T. Utilization of Sweet Potato Peels and Potato Peels for the Department of Value Added Food Products. *Int. J. Curr. Microbiol. App. Sci*. 2020;9(10):546-53.
9. Noorfarahzilah M, Lee JS, Sharifudin MS, Fadzelly MA, Hasmadi M. Applications of composite flour in development of food products. *International Food Research Journal*. 2014;21(6):2061.
10. Maruf A, Sorifa MA, Jin-cheol L, Jong-Bang E. Encapsulation by spray drying of bioactive components, physicochemical and morphological properties from purple sweet potato. *LWT-Food Science and Technology*. 2010;43(9):1307-12.
11. Ahmed M, Sorifa AM, Eun JB. Effect of pretreatments and drying temperatures on sweet potato flour. *International journal of food science & technology*. 2010;45(4): 726-32.
12. Masoodi FA, Sharma B, Chauhan GS. Use of apple pomace as a source of dietary fiber in cakes. *Plant Foods for Human Nutrition*. 2002;57(2):121-8.
13. Kaack K, Pedersen L. Low energy chocolate cake with potato pulp and yellow pea hulls. *European Food Research and Technology*. 2005;221(3):367-75.
14. Segovia Gómez F, Almajano Pablos MP. Pineapple waste extract for preventing oxidation in model food systems. *Journal of food science*. 2016;81(7):C1622-8.
15. Pao S, Petracek PD. Shelf life extension of peeled oranges by citric acid treatment. *Food Microbiology*. 1997;14(5):485-91.
16. AOAC. Official Methods of Analysis. Fourteenth Edition. Association of official analytical chemists. Washington, D.C; 1998.
17. Navacchi MF, de Carvalho JC, Takeuchi KP, Danesi ED. Development of cassava cake enriched with its own bran and *Spirulina platensis*. *Acta Scientiarum. Technology*. 2012;34(4):465-72.
18. AOAC Official Methods of Analysis. Method 991.43, total, soluble, and insoluble dietary fiber in foods. Association

- of Official Analytical Chemists International, Gaithersburg; 1998.
19. Ahmed M, Akter MS, Eun JB. Peeling, drying temperatures, and sulphite-treatment affect physicochemical properties and nutritional quality of sweet potato flour. *Food chemistry*. 2010; 121(1):112-8.
 20. Saikia S, Dutta H, Saikia D, Mahanta CL. Quality characterisation and estimation of phytochemicals content and antioxidant capacity of aromatic pigmented and non-pigmented rice varieties. *Food Research International*. 2012;46(1):334-40.
 21. Zavareze ED, Moraes KS, Salas-Mellado MD. Technological and sensory quality of cakes produced with milk whey. *Food Science and Technology*. 2010;30:100-5.
 22. Akter MS, Ahmed M, Eun JB. Effect of blanching and drying temperatures on the physicochemical characteristics, dietary fiber composition and antioxidant-related parameters of dried persimmons peel powder. *International Journal of Food Sciences and Nutrition*. 2010;61(7):702-12.
 23. Ahmed M, Akter MS, Eun JB. Optimisation of drying conditions for the extraction of β -carotene, phenolic and ascorbic acid content from yellow-fleshed sweet potato using response surface methodology. *International Journal of Food Science & Technology*. 2011;46(7):1356-62.

© 2022 Sorifa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86165>