



# Periodic Oscillation of the Solutions for a Parkinson's Disease Model

Chunhua Feng <sup>a\*</sup>

<sup>a</sup> Department of Mathematics and Computer Science, Alabama State University, Montgomery, USA.

*Author's contribution*

*The sole author designed, analysed, interpreted and prepared the manuscript.*

*Article Information*

DOI: 10.9734/ARJOM/2023/v19i10743

**Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105739>

*Received: 15/07/2023*

*Accepted: 17/09/2023*

*Published: 22/09/2023*

**Original Research Article**

## Abstract

In this paper, the oscillation of the solutions for a Parkinson's disease model with multiple delays is discussed. By linearizing the system at the equilibrium point and analyzing the instability of the linearized system, some sufficient conditions to guarantee the existence of periodic oscillation of the solutions for a delayed Parkinson's disease system are obtained. It is found that under suitable conditions on the parameters, time delay affects the stability of the system. The present method does not need to consider a bifurcating equation. Some numerical simulations are provided to illustrate our theoretical prediction.

*Keywords: Delayed Parkinson's disease model; instability; periodic solution.*

**AMS Mathematical Subject Classification: 34K13.**

*\*Corresponding author: E-mail: cfeng@alasu.edu;*

*Asian Res. J. Math., vol. 19, no. 10, pp. 217-226, 2023*

## 1 Introduction

It is known that the beta oscillations in the basal ganglia may be induced by anomalous interaction of circuits that consist of the subthalamic nucleus (STN) and the globus pallidus pars externa (GPe). The STN and the GPe form an inhibitory-excitatory coupling loop. Holgado et al. provided the following oscillation model [1]:

$$\begin{cases} \tau_S S'(t) = F_S(-W_{GS}G(t - T_{GS}) + W_{TS}Ct) - S(t), \\ \tau_G G'(t) = F_G(-W_{SG}S(t - T_{SG}) - W_{GG}G(t - T_{GG}) - W_{XG}Str) - G(t), \end{cases} \quad (1)$$

where  $S(t)$  and  $G(t)$  represent the average discharge rates of STN and GPe, respectively;  $S(t - T)$ ,  $G(t - T)$  represent the corresponding delay discharge rates.  $F_S$  and  $F_G$  are the activation functions of population  $S$  and  $G$ , showing the effect of synaptic input on the average discharge rate.  $W_{AB}$  are the weights of the connections from neural population  $A$  to neural population  $B$ .  $T_{AB}$  is the transmission delay of connections from population  $A$  to population  $B$ . The authors identified a simple set of necessary conditions on model parameters that guarantee the existence of beta oscillations in the model (1). Many authors have investigated the nonlinear dynamics in Parkinson's disease by means of the experimental method or analytical method [2-27]. For example, Wang et al. described the model by mean-field firing rate equations with discrete and distributed delays [12]:

$$\begin{cases} \tau_S S'(t) = F_S(-W_{GS}G(t - T_{GS}) + W_{CS} \int_{-\infty}^t K_1(t-s)E(s)ds) - S(t), \\ \tau_G G'(t) = F_G(W_{SG}S(t - T_{SG}) - W_{GG} \int_{-\infty}^t K_2(t-s)G(s)ds - Str) - G(t), \\ \tau_E E'(t) = F_E(-W_{SC} \int_{-\infty}^t K_3(t-s)S(s)ds - INN + c) - E(t), \end{cases} \quad (2)$$

where  $E(t)$  represents the firing rate of cortical excitatory pyramidal neurons (EXN). The authors studied the absolutely stable, conditional stable, conditional oscillation, and absolutely oscillation for model (2), which can explain different mechanisms of oscillation origin. Wang et al. systematically studied Parkinson's oscillation origin mechanism, oscillation amplitude and frequency characteristics in an improved cortex-basal ganglia (EXN-INN-STN-GPe) resonance model as follows [14]:

$$\begin{cases} \tau_S S'(t) = F_S(-W_{GS}G(t - T_{GS}) + W_{CS}E(t - T_{CS})) - S(t), \\ \tau_G G'(t) = F_G(W_{SG}S(t - T_{SG}) - Str) - G(t), \\ \tau_E E'(t) = F_E(-W_{CC}I(t - T_{CC}) - W_{SC}S(t - T_{SC}) + C) - E(t), \\ \tau_I I'(t) = F_I(W_{CC}E(t - T_{CC})) - I(t), \end{cases} \quad (3)$$

where  $F_Y(x) = \frac{M_Y}{1 + ((M_Y - B_Y)/B_Y) \exp(-4x/M_Y)}$  ( $Y = S, G, E, I$ ) are activation functions,  $I(t)$  represents the firing rate of inhibitory nuclei (INN). Assume that  $\tau_S = \tau_G = \tau_E = \tau_I = 15, T_{SG} = T_{GS} = T_{CS} = T_{CC} = T_{EE} = T$ , the Hopf bifurcation of system (3) was considered. For a modified model of the system (3) as follows:

$$\begin{cases} \tau_S S'(t) = F_S(-W_{GS}G(t - T_{GS}) + W_{CS}E(t - T_{CS})) - S(t), \\ \tau_G G'(t) = F_G(W_{SG}S(t - T_{SG}) - Str) - G(t), \\ \tau_E E'(t) = F_E(-W_{CC}I(t - T_{CC}) + W_{EE}E(t - T_{EE}) + C) - E(t), \\ \tau_I I'(t) = F_I(W_{CC}E(t - T_{CC}) - W_{II}I(t - T_{II})) - I(t). \end{cases} \quad (4)$$

Assume that

$$\tau_S = \tau_G = \tau_E = \tau_I = 10, T_{SG} = T_{GS} = T_1, T_{CS} = T_{CC} = T_{EE} = T_{II} = T_2 \quad (5)$$

The Hopf bifurcation critical condition of the system (4) was provided. However, no matter what  $\tau = 15, T_{SG} = T_{GS} = T_{CS} = T_{CC} = T_{SC} = T$ , or condition (5), those always are special cases for the parameter values. According to the simulation result in [16], the parameters  $\tau_S = 12.80ms, \tau_G = 20ms, \tau_E = 10-20ms$ , and  $\tau_I = 10-20ms$ . Therefore, the results in [14] and [16] are only for special parameters. In this paper we study the dynamic behavior for model (4) under general parameter values and do not use the bifurcation method. For convenience, we rewrite model (4) as the following:

$$\begin{cases} S'(t) = -r_1 S(t) + r_1 F_S(-W_{GS}G(t - T_{GS}) + W_{CS}E(t - T_{CS})), \\ G'(t) = -r_2 G(t) + r_2 F_G(W_{SG}S(t - T_{SG}) - Str), \\ E'(t) = -r_3 E(t) + r_3 F_E(-W_{CC}I(t - T_{CC}) + W_{EE}E(t - T_{EE}) + C), \\ I'(t) = -r_4 I(t) + r_4 F_I(W_{CC}E(t - T_{CC}) - W_{II}I(t - T_{II})), \end{cases} \quad (6)$$

where  $r_1 = \frac{1}{\tau_S}, r_2 = \frac{1}{\tau_G}, r_3 = \frac{1}{\tau_E}, r_4 = \frac{1}{\tau_I}$ . From  $F_Y(x) = \frac{M_Y}{1+((M_Y-B_Y)/B_Y)\exp(-4x/M_Y)}$  we know that  $F_S < M_S, F_G < M_G, F_E < M_E$ , and  $F_I < M_I$ . Therefore,

$$\begin{cases} S'(t) < -r_1S(t) + \tau_1M_S, \\ G'(t) < -r_2G(t) + \tau_2M_G, \\ E'(t) < -r_3E(t) + \tau_3M_E, \\ I'(t) < -r_4I(t) + \tau_4M_I. \end{cases} \tag{7}$$

System (7) implies that  $|S(t)| < \frac{r_1M_S}{r_1} = M_S, |G(t)| < \frac{r_2M_G}{r_2} = M_G, |E(t)| < \frac{r_3M_E}{r_3} = M_E$ , and  $|I(t)| < \frac{r_4M_I}{r_4} = M_I$ , in other words, all of the solutions of system (4) are boundedness. According to the parameter values in [16]:  $M_S=300$  spk/s,  $B_S=17$  spk/s,  $M_G=400$  spk/s,  $B_G=75$  spk/s,  $M_E=71.77$  spk/s,  $B_E=3.62$  spk/s,  $M_I=276$  spk/s,  $B_I=7.18$  spk/s, we know that  $F_Y(x)$  are monotone increasing functions for  $Y = S, G, E$ , and  $I$ . Therefore system (4) has a unique equilibrium point  $(S^*, G^*, E^*, I^*)^T$ . Make the change of variables  $S(t) \rightarrow S(t) - S^*, G(t) \rightarrow G(t) - G^*, E(t) \rightarrow E(t) - E^*, I(t) \rightarrow I(t) - I^*$ , the Taylor expansion of system (4) at the equilibrium point is the following:

$$\begin{cases} S'(t) = -r_1S(t) + a_{12}G(t - T_{GS}) + a_{13}E(t - T_{CS}) \\ \quad + \sum_{i+j \geq 2} \frac{[G(t-T_{GS})]^i [E(t-T_{CS})]^j}{i!j!} \cdot \frac{\partial^{i+j} F_S}{\partial G^i \partial E^j} |_{(G^*, E^*)}, \\ G'(t) = -r_2G(t) + a_{21}S(t - T_{SG}) + F_G''|_{S^*} S^2(t - T_{SG}) + \dots, \\ E'(t) = -r_3E(t) + a_{33}E(t - T_{EE}) + a_{34}I(t - T_{CC}) \\ \quad + \sum_{i+j \geq 2} \frac{[E(t-T_{EE})]^i [I(t-T_{CC})]^j}{i!j!} \cdot \frac{\partial^{i+j} F_E}{\partial E^i \partial I^j} |_{(E^*, I^*)}, \\ I'(t) = -r_4I(t) + a_{43}E(t - T_{CC}) + a_{44}I(t - T_{II}) \\ \quad + \sum_{i+j \geq 2} \frac{[E(t-T_{CC})]^i [I(t-T_{II})]^j}{i!j!} \cdot \frac{\partial^{i+j} F_I}{\partial E^i \partial I^j} |_{(E^*, I^*)}, \end{cases} \tag{8}$$

where  $a_{12} = r_1 \frac{\partial F_S}{\partial G} |_{(G^*, E^*)}, a_{13} = r_1 \frac{\partial F_S}{\partial E} |_{(G^*, E^*)}, a_{21} = r_2 F_G' |_{S^*}, a_{33} = r_3 \frac{\partial F_E}{\partial E} |_{(E^*, I^*)}, a_{34} = r_3 \frac{\partial F_E}{\partial I} |_{(E^*, I^*)}, a_{43} = r_4 \frac{\partial F_I}{\partial E} |_{(E^*, I^*)}, a_{44} = r_4 \frac{\partial F_I}{\partial I} |_{(E^*, I^*)}$ . The linearized system of system (8) is the follows:

$$\begin{cases} S'(t) = -r_1S(t) + a_{12}G(t - T_{GS}) + a_{13}E(t - T_{CS}), \\ G'(t) = -r_2G(t) + a_{21}S(t - T_{SG}), \\ E'(t) = -r_3E(t) + a_{33}E(t - T_{EE}) + a_{34}I(t - T_{CC}), \\ I'(t) = -r_4I(t) + a_{43}E(t - T_{CC}) + a_{44}I(t - T_{II}), \end{cases} \tag{9}$$

Let  $s = \min\{T_{GS}, T_{CS}, T_{SG}, T_{EE}, T_{CC}, T_{II}\}$ . Consider a special case of the system (9):

$$\begin{cases} S'(t) = -r_1S(t) + a_{12}G(t - s) + a_{13}E(t - s), \\ G'(t) = -r_2G(t) + a_{21}S(t - s), \\ E'(t) = -r_3E(t) + a_{33}E(t - s) + a_{34}I(t - s), \\ I'(t) = -r_4I(t) + a_{43}E(t - s) + a_{44}I(t - s), \end{cases} \tag{10}$$

The matrix form of the system (10) is as follows:

$$u'(t) = Cu(t) + Au(t - s), \tag{11}$$

where  $u(t) = [S(t), G(t), E(t), I(t)]^T, u(t - s) = [S(t - s), G(t - s), E(t - s), I(t - s)]^T$ , and

$$A = (a_{ij})_{4 \times 4} = \begin{pmatrix} 0 & a_{12} & a_{13} & 0 \\ a_{21} & 0 & 0 & 0 \\ 0 & 0 & a_{33} & a_{34} \\ 0 & 0 & a_{43} & a_{44} \end{pmatrix},$$

$$B = (b_{ij})_{4 \times 4} = \text{diag}(-r_1 - r_2 - r_3 - r_4).$$

## 2 The Existence of Periodic Solution

Since the system (9) is a linearized system of (8). Thus, we can see that the system (8) is a disturbed system of (9). If the trivial solution of system (9) is unstable, then the trivial solution of system (8) is also unstable. In what follows, we first consider the instability of the zero equilibrium point of the system (10) (or (11)). So we have the following Theorems.

**Theorem 1.** Assume that the system (11) has a unique trivial solution,  $\alpha_1, \alpha_2, \alpha_3, \alpha_4$  are characteristic values of matrix  $A$ . If there is a characteristic value, say  $\alpha_1$ ,

- (i)  $Re(\alpha_1) = 0, Im(\alpha_1) \neq 0$ , and  $\alpha_1 = \omega i$ ;
- (ii)  $Re(\alpha_1) > 0$ , and  $Re(\alpha_1) > \max\{r_1, \dots, r_4\}$ ,
- (iii)  $Im(\alpha_1) = 0, \alpha_1 > 0$  and  $\alpha_1 > \max\{r_1, \dots, r_4\}$ . Then the trivial solution of system (11) (thus the system (8)) is unstable, implying that there exists a limit cycle in the system (4), namely, system (4) has a periodic solution.

**Proof.** We will show that the trivial solution of the system (11) is unstable. When  $Re(\alpha_1) = 0, Im(\alpha_1) \neq 0$ , and  $\alpha_1 = \omega i$ , then  $e^{\omega it} = \cos \omega t + i \sin \omega t$ . Since  $\cos \omega t$  is a periodic function, therefore, the trivial solution of system (11) is unstable. Obviously, all characteristic values of matrix  $B$  are  $-r_1, -r_2, -r_3, -r_4$ . Since  $\alpha_1, \alpha_2, \alpha_3, \alpha_4$  are characteristic values of matrix  $A$ , then the characteristic equation of the system (11) is the following:

$$\prod_{i=1}^4 \lambda + r_i - \alpha_i e^{-\lambda s} = 0. \tag{12}$$

So we have

$$\lambda + r_1 - \alpha_1 e^{-\lambda s} = 0. \tag{13}$$

If  $Re(\alpha_1) > 0$ , and  $Re(\alpha_1) > \max\{r_1, \dots, r_4\}$ , or  $\alpha_1 > 0$ , and  $\alpha_1 > \max\{r_1, \dots, r_4\}$ , this means that the equation (13) has a positive real part characteristic value or a positive characteristic value. It suggests that the trivial solution of the system (11) is unstable. According to the basic theory of delayed differential equation, if the trivial solution is unstable for a small delay, then the trivial solution is still unstable as the delay increases [28]. In other words, the trivial solution of system (9) is unstable. This implies that the equilibrium point  $(S^*, G^*, E^*, I^*)^T$  of system (8) is unstable. Equivalently, the unique equilibrium point of the system (4) is unstable. This instability of the unique equilibrium point together with the boundedness of the solutions will force system (4) to generate a limit cycle, namely, a periodic solution according to the extended Chafee's criterion [29, 30].The proof is completed.

Let  $m = \max\{|a_{21}|, |a_{12}|, a_{33} + |a_{13}| + |a_{43}|, a_{44} + |a_{34}|, \}$  then we have

**Theorem 2.** Assume that the system (11) has a unique trivial solution. If the following condition holds

$$m - \tau > 0. \tag{14}$$

where  $\tau = \min\{r_1, \dots, r_4\}$ . Then the trivial solution of system (11) is unstable, implying that there exists a limit cycle of system (4), namely, system (4) has a periodic solution.

**Proof.** To prove the instability of the trivial solution of the system (11), Let  $z(t) = |S(t)| + |G(t)| + |E(t)| + |I(t)|$ . Therefore,  $z(t) > 0$  for  $t > 0$ , and

$$z'(t) \leq -\tau z(t) + mz(t-s). \tag{15}$$

Specifically, consider a scalar equation



$$v'(t) = -\tau v(t) + mv(t-s). \tag{16}$$

According to the comparison theory of differential equation, we have  $z(t) \leq v(t)$ . We claim that the trivial solution of equation (16) is unstable. Indeed, the characteristic equation of (16) is as follows:



$$\lambda = -\tau + m e^{-\lambda s}. \tag{17}$$



Consider a function  $\varphi(\lambda) = \lambda + \tau - me^{-\lambda s}$ . Then  $\varphi(\lambda)$  is a continuous function of  $\lambda$ . Noting that  $\varphi(0) = \tau - m = -(m - \tau) < 0$ . Obviously, there exists a  $L > 0$  such that  $\varphi(L) = L + \tau - me^{-Ls} > 0$ . By the Intermediate Value Theorem, there exists a  $\lambda_0 \in (0, L)$  such that  $\varphi(\lambda_0) = 0$ . In other words, there exists a positive characteristic root of the equation (16), which means that the trivial solution of equation (15) is unstable, implying that the trivial solution of system (11), thus (4) is unstable. Similar to Theorem 1, the system (4) has a periodic solution. The proof is completed.

**Fig.1** Periodic oscillation of the solutions,  $r_1=0.05, r_2=0.1, r_3=0.1, r_4=0.08$ ,  
delays:  $T_{SG}=15.2, T_{GS}=15.3, T_{CS}=T_{CC}=T_{EE}=15.4, T_{II}=15.5$ .



**Fig.2** Periodic oscillation of the solutions,  $r_1=0.08, r_2=0.05, r_3=0.08, r_4=0.0625$ ,  
delays:  $T_{SG}=15.2, T_{GS}=15.3, T_{CS}=T_{CC}=T_{EE}=15.4, T_{II}=15.5$ .









### 3 Computer Simulation Result

In model (4), according to the parameters in [16], setting  $M_S = 300, M_G = 400, M_E = 72, M_I = 276, W_{GS} = 3, W_{CS} = 6, W_{SG} = 2.5, W_{II} = 0.1, W_{EE} = 1, W_{CC} = 3, B_G = 75, B_S = 17, B_E = 3.6, B_I = 7, C = 277, Str =$

**Fig.5** Periodic oscillation of the solutions,  $r_1=0.1, r_2=0.2, r_3=0.2, r_4=0.05$ ,  
delays:  $T_{SG}=24.8, T_{GS}=25.5, T_{CS}=T_{EE}=26.4, T_{CC}=24.2, T_{II}=24.5$ .



**Fig.6** Periodic oscillation of the solutions,  $r_1=0.05, r_2=0.1, r_3=0.16, r_4=0.1$ ,  
delays:  $T_{SG}=24.8, T_{GS}=25.5, T_{CS}=T_{EE}=26.4, T_{CC}=24.2, T_{II}=24.5$ .



40, when we select the time delay  $T_{SG} = 15.2, T_{GS} = 15.3, T_{CS} = T_{CC} = T_{EE} = 15.4, T_{II} = 15.5$ , firstly,  $\tau_S = 20, \tau_G = 10, \tau_E = 10, \tau_I = 12.5$ , namely,  $r_1 = 0.05, r_2 = 0.1, r_3 = 0.1, r_4 = 0.08$ , then the unique positive equilibrium point  $(S^*, G^*, E^*, I^*) = (61.2085, 167.5346, 60.7945, 72.0425)$ . Thus,  $a_{12} = r_1 \frac{\partial F_S}{\partial G} |_{(G^*, E^*)} = -0.0055, a_{13} = r_1 \frac{\partial F_S}{\partial E} |_{(G^*, E^*)} = 0.0112, a_{21} = r_2 F'_G |_{S^*} = 0.1213, a_{33} = r_3 \frac{\partial F_E}{\partial E} |_{(E^*, I^*)} = -0.0963, a_{34} = r_3 \frac{\partial F_E}{\partial I} |_{(E^*, I^*)} = 0.0328, a_{43} = r_4 \frac{\partial F_I}{\partial E} |_{(E^*, I^*)} = 0.2186, a_{44} = r_4 \frac{\partial F_I}{\partial I} |_{(E^*, I^*)} = -0.0076$ . The characteristic values of matrix  $A_1$  are  $0.0436, -0.1475, 0.0258i, -0.0258i$ . Since there exists a characteristic value  $0.0258i$ , and the conditions of Theorem 1 are satisfied. There exists a periodic oscillatory solution (see Fig.1). Then setting  $\tau_S = 12.5, \tau_G = 20, \tau_E = 12.5, \tau_I = 16$ , namely,  $r_1 = 0.08, r_2 = 0.05, r_3 = 0.08, r_4 = 0.0625$ , the other parameters are the same as Fig. 1, then  $(S^*, G^*, E^*, I^*) = (64.3124, 164.6548, 62.4903, 75.8162)$ .

There exists a periodic oscillatory solution (see Fig.2). When we select  $\tau_S = 10, \tau_G = 12.5, \tau_E = 8, \tau_I = 5$ , namely,  $r_1 = 0.1, r_2 = 0.08, r_3 = 0.125, r_4 = 0.2$ , the other parameters are the same as Fig. 1, then  $(S^*, G^*, E^*, I^*) = (51.9186, 142.0682, 54.1432, 64.9544)$ . There exists a periodic oscillatory solution (see Fig.3). When we select the time delay  $T_{SG} = 24.8, T_{GS} = 25.5, T_{CS} = T_{EE} = 26.4, T_{CC} = 24.2, T_{II} = 24.5, \tau_S = 20, \tau_G = 10, \tau_E = 10, \tau_I = 12.5$ , namely,  $r_1 = 0.05, r_2 = 0.1, r_3 = 0.1, r_4 = 0.08$ , then the unique positive equilibrium point  $(S^*, G^*, E^*, I^*) = (79.7544, 143.0359, 49.5957, 66.1679)$ . Thus,  $a_{12} = -0.0623, a_{13} = 0.1246, a_{21} = 0.2487, a_{33} = -0.0641, a_{34} = 0.0213, a_{43} = 0.1352, a_{44} = -0.0046$ . The characteristic values of matrix  $A_2$  are  $0.0271, -0.0975, 0.1245i, -0.1245i$ , and the conditions of Theorem 1 are satisfied. There exists a periodic oscillatory solution (see Fig.4). Then setting  $\tau_S = 10, \tau_G = 5, \tau_E = 5, \tau_I = 10$ , namely,  $r_1 = 0.1, r_2 = 0.2, r_3 = 0.2, r_4 = 0.05$  the other parameters are the same as Fig. 4, then  $(S^*, G^*, E^*, I^*) = (88.6328, 209.1742, 56.4323, 68.6519)$ . There exists a periodic oscillatory solution (see Fig.5). When we select  $\tau_S = 20, \tau_G = 10, \tau_E = 6.25, \tau_I = 10$ , namely,  $r_1 = 0.05, r_2 = 0.1, r_3 = 0.16, r_4 = 0.1$ , the other parameters are the same as figure 4, then  $(S^*, G^*, E^*, I^*) = (42.4116, 178.3734, 52.2841, 62.1911)$ . There exists a periodic oscillatory solution (see Fig.6). From the figures, we see that time delays affect the oscillatory frequency.

## 4 Conclusion

The present paper discusses the oscillation of the solutions for a Parkinson's disease model with multiple delays by means of the extended Chafee's criterion. Two sufficient conditions are provided to guarantee the existence of periodic solutions. We change the nonlinear model to an equivalent system. The instability of the solution of the equivalent system implies the instability of the equilibrium point of the original system. Our criterion is only a sufficient condition.

## Competing Interests

Author has declared that no competing interests exist.

## References

- [1] Holgado AJ, Terry JR, Bogacz R. Conditions for the generation of beta oscillations in the subthalamic nucleus-globus pallidus network. *J. Neuroscience*. 2010;30:12340-12352.
- [2] Chen Y, Wang J, Kang Y, Ghori MB. Emergence of beta oscillations of a resonance model for Parkinson's disease. *Neural Plast*. 2020;2020:8824760.
- [3] Al-Tuwairqi SM, Badrah AA. Modeling the dynamics of innate and adaptive immune response to Parkinson's disease with immunotherapy. *AIMS Mathematics*. 2023;8(1):1800-32.
- [4] Tang CC, Poston KL, Dhawan V, Eidelberg D. Abnormalities in metabolic network activity precede the onset of motor symptoms in Parkinson's disease. *Journal of Neuroscience*. 2010;30(3):1049-56.
- [5] Schapira AH, Jenner P. Etiology and pathogenesis of Parkinson's disease. *Movement disorders*. 2011;26(6):1049-55.
- [6] Gubellini P, Kachidian P. Animal models of Parkinson's disease: An updated overview. *Revue neurologique*. 2015 Nov 1;171(11):750-61.
- [7] Raza C, Anjum R. Parkinson's disease: Mechanisms, translational models and management strategies. *Life sciences*. 2019 Jun 1;226:77-90.
- [8] Jane YN, Nehemiah HK, Arputharaj K. A Q-backpropagated time delay neural network for diagnosing severity of gait disturbances in Parkinson's disease. *Journal of Biomedical Informatics*. 2016;60:169-76.
- [9] Rubin JE, Terman D. High frequency stimulation of the subthalamic nucleus eliminates pathological thalamic rhythmicity in a computational model. *Journal of computational neuroscience*. 2004;16:211-35.



- [10] Terman D, Rubin JE, Yew AC, Wilson CJ. Activity patterns in a model for the subthalamopallidal network of the basal ganglia. *Journal of Neuroscience*. 2002 Apr 1;22(7):2963-76.
- [11] Hu B, Xu M, Zhu L, Lin J, Wang Z, Wang D, Zhang D. A bidirectional Hopf bifurcation analysis of Parkinson's oscillation in a simplified basal ganglia model. *Journal of Theoretical Biology*. 2022 Mar 7;536:110979.
- [12] Wang Z, Hu B, Zhou W, Xu M, Wang D. Hopf bifurcation mechanism analysis in an improved cortex-basal ganglia network with distributed delays: An application to Parkinson's disease. *Chaos, Solitons & Fractals*. 2023 Jan 1;166:113022.
- [13] Chen Y, Wang J, Kang Y, Ghorl MB. Emergence of Beta Oscillations of a Resonance Model for Parkinson's Disease. *Neural Plasticity*. 2020;2020:8824760.
- [14] Wang Z, Hu B, Zhu L, Lin J, Xu M, Wang D. The possible mechanism of direct feedback projections from basal ganglia to cortex in beta oscillations of Parkinson's disease: A theoretical evidence in the competing resonance model. *Communications in Nonlinear Science and Numerical Simulation*. 2023;120:107142.
- [15] Hu B, Xu M, Wang Z, Jiang D, Wang D, Zhang D. The theoretical mechanism of Parkinson's oscillation frequency bands: a computational model study. *Cognitive Neurodynamics*. 2021 Aug;15(4):721-31.
- [16] Wang Z, Hu B, Zhu L, Lin J, Xu M, Wang D. Hopf bifurcation analysis for Parkinson oscillation with heterogeneous delays: A theoretical derivation and simulation analysis. *Communications in Nonlinear Science and Numerical Simulation*. 2022 Nov 1;114:106614.
- [17] Van Wijk BC, Beudel M, Jha A, Oswal A, Foltyniec T, Hariz MI, Limousin P, Zrinzo L, Aziz TZ, Green AL, Brown P. Subthalamic nucleus phase-amplitude coupling correlates with motor impairment in Parkinson's disease. *Clinical Neurophysiology*. 2016 Apr 1;127(4):2010-9.
- [18] Burke RE, O'Malley K. Axon degeneration in Parkinson's disease. *Experimental neurology*. 2013 Aug 1;246:72-83.
- [19] Lobb CJ. Abnormal bursting as a pathophysiological mechanism in Parkinson's disease. *Basal ganglia*. 2014 Apr 1;3(4):187-95.
- [20] Wang J, Hirschmann J, Elben S, Hartmann CJ, Vesper J, Wojtecki L, Schnitzler A. High-frequency oscillations in Parkinson's disease: Spatial distribution and clinical relevance. *Movement disorders*. 2014;29(10):1265-72.
- [21] Darcy N, Lofredi R, Al-Fatly B, Neumann WJ, Hübl J, Brücke C, Krause P, Schneider GH, Kühn A. Spectral and spatial distribution of subthalamic beta peak activity in Parkinson's disease patients. *Experimental neurology*. 2022 Oct 1;356:114150.
- [22] Pang X, Pan X, Xu Y. Multi-parameter safe sample elimination rule for accelerating nonlinear multi-class support vector machines. *Pattern Recognition*. 2019;95:1-1.
- [23] Wilson H, Niccolini F, Pellicano C, Politis M. Cortical thinning across Parkinson's disease stages and clinical correlates. *Journal of the Neurological Sciences*. 2019;398:31-8.
- [24] Geng X, Xu X, Horn A, Li N, Ling Z, Brown P, Wang S. Intra-operative characterisation of subthalamic oscillations in Parkinson's disease. *Clinical Neurophysiology*. 2018;129(5):1001-10.
- [25] Eugeni M, Dessi D, Mastroddi F. A Normal Form analysis in a finite neighborhood of a Hopf bifurcation: on the Center Manifold dimension. *Nonlinear Dynamics*. 2018;91:1461-72.
- [26] Grado LL, Johnson MD, Netoff TI. Bayesian adaptive dual control of deep brain stimulation in a computational model of Parkinson's disease. *PLoS computational biology*. 2018;14(12):e1006606.
- [27] Holt AB, Netoff TI. Origins and suppression of oscillations in a computational model of Parkinson's disease. *Journal of computational neuroscience*. 2014;37:505-21.

- [28] Hale JK, Verduyn SM. Introduction to functional differential equations. New York, Springer; 1993.
- [29] Chafee N. A bifurcation problem for a functional differential equation of finitely retarded type. Journal of Mathematical Analysis and Applications. 1971 Aug 1;35(2):312-48.
- [30] Feng C, Plamondon R. An oscillatory criterion for a time delayed neural ring network model. Neural Networks. 2012;29:70-9.

---

© 2023 Feng; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Peer-review history:**

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/105739>