



# Study of Foliar Nutrition Applied at Different Growth Stages on Niger (*Guizotia abyssinica* L.)

Dipika G. Chaudhari <sup>a++\*</sup>, Swapnil P. Deshmukh <sup>b#</sup>,  
Narendra Singh <sup>c†</sup>, V. Surve <sup>d†</sup>, B. M. Mote <sup>e†</sup>  
and Gajera B. M. <sup>a++</sup>

<sup>a</sup> Department of Agronomy, N.M. College of Agriculture, Navsari Agricultural University, Navsari- 396 450, Gujarat, India.

<sup>b</sup> Niger Research Station, Navsari Agricultural University, Vanarasi- 396 580, Gujarat, India.

<sup>c</sup> Department of Soil Science, Navsari Agricultural University, Navsari- 396 450, India.

<sup>d</sup> Department of Agronomy, College of Agriculture, Navsari Agricultural University, Bharuch- 392 012, India.

<sup>e</sup> Office of Director of Research, Navsari Agricultural University, Navsari- 396 450, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i193653

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104921>

Original Research Article

Received: 08/06/2023

Accepted: 14/08/2023

Published: 01/09/2023

## ABSTRACT

A field experiment was conducted during *Rabi* season of 2021-22 at Niger Research Station, NAU, Vanarasi, Gujarat to study the effect of foliar spray of nutrients applied at different growth stages on *rabi niger* (*Guizotia abyssinica* L.). The experiment was laid out in randomized block design with

<sup>++</sup> M.Sc. Scholar;

<sup>#</sup> Assistant Research Scientist;

<sup>†</sup> Assistant Professor;

\*Corresponding author: E-mail: [chaudharidipika1999@gmail.com](mailto:chaudharidipika1999@gmail.com);

three replications. The treatments consisted of seven treatments including foliar spray of 19:19:19 and Novel organic liquid (A Patented NAU product) applied at different growth stages on niger crop. The experimental results revealed that foliar application 1% 19:19:19 applied at flowering and capitula formation stage recorded significantly higher number of branches per plant. The number of seeds per capitulum and seed yield also remained significantly higher with the application 1% 19:19:19 applied at flowering and capitula formation stage which was found at par with treatment receiving foliar spray of 1% 19:19:19 at flowering stage and foliar spray of 1 % novel organic liquid at flowering stage.

**Keywords:** Niger crop; foliar spray; 19:19:19; novel organic liquid.

## 1. INTRODUCTION

Oilseeds occupy a key place in agricultural economy of India, constituting an important group of crops next to cereals. Niger (*Guizotia abyssinica* L.) is believed to have originated from Ethiopia where it was domesticated from about 2000 BCE, and then spread to India. In India, Niger is grown on an area of 2.61 lakh ha mainly during *khari*. Andhra Pradesh, Assam, Chhattisgarh, Gujarat, Jharkhand, Karnataka, Madhya Pradesh, Maharashtra, Odisha and West Bengal are the other states where niger is grown. The average yield in India is 3.21 q/ ha. Niger seeds contains edible oil (38 to 43%), protein (18 to 24%), sugar (12%) and minerals essential for human and animal meals. The main aim of foliar fertilization is to eliminate "hidden hunger", which is a very important consideration in high-input production systems and during critical stages of growth when roots are unable to assimilate the required amounts of nutrients, even in fertile soils. One of the best foliar fertilizers 19:19:19 for crop like pulses, vegetables, paddy for green house cultivation, nurseries, kitchen garden and for all types of field crop. The use of 19:19:19 resulted in cost saving and at the same time significantly increases the yield. Economic benefit ultimately matters for farmers and novel organic liquid nutrient plays an important role to improved it, [1]. Novel organic liquid nutrient is a product of Navsari Agricultural University, Navsari, Gujarat which was patented in the year of 2012. Since foliar nutrient usually penetrate the leaf cuticle or stomata and enter the cell facilitating easy and rapid utilization of nutrients, so foliar nutrition on niger helps in achieving the optimum grain yield of niger.

## 2. MATERIALS AND METHODS

The field experiment entitled effect of foliar nutrition applied at different growth stages on *rabi* niger (*Guizotia abyssinica* L.) was conducted during *Rabi* season under irrigated

condition of the year 2021-22 at Niger Research Station, Navsari Agricultural University, Vanarasi. The soil of the experimental field was black color and clayey in texture with medium in available nitrogen, low in available phosphorus and high in available potassium and moderately alkaline in reaction. The experiment was laid out in randomized block design with three replications. The treatments consisted of seven level of foliar nutrition applied at different growth stages of niger crop viz., F<sub>1</sub>: foliar spray of water (Control), F<sub>2</sub>:1% 19:19:19 at flowering stage, F<sub>3</sub>:1% 19:19:19 at capitula formation stage, F<sub>4</sub>:1% 19:19:19 at flowering and capitula formation stage, F<sub>5</sub>:1% Novel organic liquid at flowering stage, F<sub>6</sub>:1% Novel organic liquid at capitula formation stage, F<sub>7</sub>:1% Novel organic liquid at flowering and capitula formation stage. The niger cultivar GNNIG-3 was fertilized with recommended dose 20 kg N and 20 kg P<sub>2</sub>O<sub>5</sub>. At the time of sowing Nitrogen was supplied through urea while phosphorus supplied through single super phosphate.

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Parameters in Niger as Influenced by Foliar Nutrition Applied at Different Growth Stages

Data presented in Table 1 indicated that plant height at 60 DAS it was recorded higher with foliar spray of 1% 19:19:19 at flowering stage (F<sub>2</sub>) (51.07cm). Although plant height did not differ statistically, but numerical values showed higher plant height in all the foliar nutrition treatments at harvest stage compared to control (F<sub>1</sub>). Foliar spray of 1% 19:19:19 at flowering and capitula formation stage (F<sub>4</sub>) showed significantly higher number of branches per plant at harvest stage (6.13) across all the treatments. Treatment F<sub>4</sub> was found at par with treatment 1 % 19:19:19 at flowering stage (F<sub>2</sub>) and treatment 1% novel organic liquid at flowering stage (F<sub>5</sub>).

It might be due to hastening various metabolic process *viz.*, photosynthesis, symbiotic biological N<sub>2</sub> fixation process due to nutrient availability at the initial stage of the crop. Further, phosphorus present in 19:19:19 fertilizer absorbed directly by the plant might have increased cell division and cell development leading to more number of branches. Similar results were obtained by Shwetha et al. [2] in groundnut crop Banasode and Math [3]. Basu et al. [4] in Sesamum crop and Deshmukh et al. [5] in soybean.

### 3.2 Yield Attributes and Economics in Niger as Influenced by Foliar Nutrition Applied at Different Growth Stages

#### 3.2.1 Yield attributes

The Study revealed that significantly higher number of seeds per capitulum (27.76) was recorded with foliar spray of 1% 19:19:19 at flowering and capitula formation stage (F<sub>4</sub>) and was found at par with rest of foliar nutrition treatments except 1% novel organic liquid at capitula formation stage (F<sub>6</sub>) and foliar spray of water (control) (F<sub>1</sub>). The lowest number of seeds per capitula (20.03) were recorded with foliar spray of water (control) (F<sub>1</sub>). significantly higher seed yield (514kg/ha) was recorded with foliar spray of 1% 19:19:19 at flowering and capitula formation stage (F<sub>4</sub>) and was found at par with 1% 19:19:19 at flowering stage (F<sub>2</sub>), 1% Novel organic liquid at flowering and capitula formation stage (F<sub>7</sub>) and 1% Novel organic liquid at flowering stage (F<sub>5</sub>). While, significantly the lower seed yield (379kg/ha) was recorded with

foliar spray of water (F<sub>1</sub>). The increase in yield due to foliar nutrition treatment in F<sub>4</sub>, F<sub>2</sub>, F<sub>7</sub> and F<sub>5</sub> as compared to control (F<sub>1</sub>) was 35%, 27%, 23% and 21%, respectively.

It might be due to the foliar spray of 19:19:19 that increased number of branches, capitula per plant and seeds per capitulum which directly related to seed yield per plant and production per unit area. Further, foliar spray may have increased the photosynthetic activity and delays the senescence of leaves, which enhance the supply of photosynthate available for grain filling, resulting in more number of capitula per plant and ultimately seed yield. The results are in confirmation with finding of Sethi et al. [6] in Indian mustard, Krishnaveni et al. [7] Monika et al. [8] and Sarika et al. [9] in sesame crop; Bodke et al. [10] and Deshmukh et al. [5] in soybean crop. Novel organic liquid nutrients which provide nutrient supply as well as growth promoting hormones. Application of novel at early growth stages thus might have improved growth and development of the crop resembling in number of branches, capitula per plant and seeds per capitula, which is further reflected in increase of seed yield. Similar finding was reported by Singhal et al. [1] and Champaneri et al. [11].

#### 3.2.2 Economics

The study revealed that maximum net return of ₹17676/ha with BCR of 1.96 were accrued with treatment 1% 19:19:19 at flowering and capitula formation stage (F<sub>4</sub>)

**Table 1. Study of growth parameters in niger as influenced by foliar nutrition applied at different growth stages**

Treatments	Plant height (cm)			Number of branches
	30 DAS	60 DAS	At harvest	
F <sub>1</sub> : Foliar spray of water (control)	13.02	50.33	73.17	4.63
F <sub>2</sub> : 1% 19:19:19 at flowering stage	12.73	51.07	78.77	5.87
F <sub>3</sub> : 1% 19:19:19 at capitula formation stage	12.38	47.90	76.73	5.27
F <sub>4</sub> : 1% 19:19:19 at flowering and capitula formation stage	13.05	50.48	97.07	6.13
F <sub>5</sub> : 1% Novel organic liquid at flowering stage	13.40	50.89	79.20	5.93
F <sub>6</sub> : 1% Novel organic liquid at capitula formation stage	11.93	46.20	75.10	5.20
F <sub>7</sub> : 1% Novel organic liquid at flowering and capitula formation stage	12.98	50.20	78.93	5.80
SEM±	0.38	1.88	2.91	0.23
CD at 5%	NS	NS	NS	0.73
CV%	5.3	6.6	6.5	7.5

**Table 2. Study of yield attributes and economics in niger as influenced by foliar nutrition applied at different growth stages**

Treatments	Number of seed per capitulum	Seed yield (kg/ha)	Net return (₹/ha)	B:C ratio
F <sub>1</sub> : Foliar spray of water (control)	20.03	379	10602	1.76
F <sub>2</sub> : 1% 19:19:19 at flowering stage	25.87	482	16872	2.00
F <sub>3</sub> : 1% 19:19:19 at capitula formation stage	22.72	426	12956	1.86
F <sub>4</sub> : 1% 19:19:19 at flowering and capitula formation stage	27.76	514	17676	1.96
F <sub>5</sub> : 1% Novel organic liquid at flowering stage	25.13	462	15812	1.95
F <sub>6</sub> : 1% Novel organic liquid at capitula formation stage	21.74	418	12732	1.87
F <sub>7</sub> : 1% Novel organic liquid at flowering and capitula formation stage	25.65	468	15096	1.85
SEm±	1.05	24		
CD at 5%	3.25	74		
CV%	7.6	9.2		

followed by foliar spray of 1% 19:19:19 at flowering stage (F<sub>2</sub>) with net return of 16872 ₹/ha with BCR of 2.00 and 1% novel organic liquid at flowering stage (F<sub>5</sub>) with net return of 15812 ₹/ha with BCR of 1.95, where as the lowest net return of ₹10602/ha were obtained with foliar spray of water (control) (F<sub>1</sub>) with BCR 1.76.

#### 4. CONCLUSION

On the basis of one year field experiment, it can be concluded that spraying of 1% 19:19:19 or 1% novel organic liquid at flowering stage may be done for getting profitable niger production.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Singhal VK, Patel GG, Patel DH, Kumar U, Saini LK. Effect of foliar application of water soluble fertilizers on growth, yield and economics of vegetable cowpea production. *Ecosan*. 2015; 7: 79-83.
- Shwetha BN, Anupama C, Sowmya TM, Yaligar R. Effect of foliar nutrition on productivity of groundnut crop. *Journal of Pharmacognosy and Phytochemistry*. 2018;7 (1S):2357-2360.
- Banasode C, Math KK. Effect of foliar fertilization of water-soluble fertilizers on growth and economics of soybean in a vertisol. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(2):2391-2393.
- Basu BY, Wathi PS, Rao NS, and Naik VS. Enhancement of productivity through soil and foliar application of nutrients in summer sesamum (*Sesamum indicum* L.). *Journal of Pharmacognosy and Phytochemistry*. 2019;8(4):3574-3577.
- Deshmukh SP, Rathva RS, Bambhaneeya SM, Jagtap PK, Mistry PM. Effect of foliar nutrition applied at different growth stages soybean (*Glycine Max.*) under rainfed condition. *Journal of Oilseed Research*. 2023;39(special issue):467-469.
- Sethi IB, Singh H, Kumar S, Jajoria M, Jat LK, Braod N K, and Mali HR. (2021). Effect of water-soluble fertilizers on yield and economics of Indian mustard (*Brassica juncea* (L.). *Trends in Biosciences*. 2021;13 (9):532-538.
- Krishnaveni SA, Supriya C, Sridhar SM. Impact of foliar nutrition on the yield and economics of green gram (*Vigna radiata*). *International Journal of Chemical Studies*. 2021;9(2):11-13.
- Monika P, Sumathi V, Reddy PM, Naidu M, Sagar GK. Impact of fertilizers and foliar sprays on productivity of summer sesame (*Sesamum indicum* L.) Andhra Pradesh *Journal of Agricultural Sciences*. 2021;7(3): 173-177.
- Sarika TW, Prasanna J, Sonawane DA, Ghodke PB, and Ubale SP. Effect of fertilizer levels and foliar nutrition on growth and yield of soybean (*Glycine max* (L.) Merrill). *International Journal of Plant and Soil Science*. 2022; 34(14):13-18.

10. Bodke VS, Hanwate GR, Gorde NB, Jaybhaye BB, Pidurkar PK. Studies on influence of foliar application of water-soluble fertilizers on growth and yield of soybean (*Glycine max* L.). The Pharma Innovation Journal. 2022;11(12):1682-1686.
11. Champaneri DD, Patel NK, Desai CS, Desai DH. Efficacy of novel organic liquid nutrient and novel plus organic liquid nutrient on quantitative traits of Indian bean [*Lablab purpureus* (L.) Sweet]. International Journal of Plant and Soil Science. 2021;33(17):105-115.

© 2023 Chaudhari et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/104921>*