



White Tongue as a Manifestation of Diverse Skin Diseases

Khalifa Sharquie ^{a##}, Faris O. Hadi ^{b^o} and Raed I. Jabbar ^{c^o}

^a Department of Dermatology, College of Medicine, University of Baghdad, Medical City Teaching Hospital, Baghdad, Iraq.

^b Department of Dermatology, Suq Alshuyukh General Hospital, Thiqr Health Directorate, Thiqr, Iraq.

^c Department of Dermatology, Fallujah Teaching Hospital, Al-Anbar Health Directorate, Anbar, Iraq.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Giuseppe Murdaca, University of Genova, Italy.

Reviewers:

(1) Sehar Afshan Naz, Federal Urdu University of Arts, Science and Technology, Pakistan.

(2) Sreeshyla HS, JSSDC&H (JSS Dental College and Hospital), India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/78170>

Original Research Article

Received 06 October 2021
Accepted 13 December 2021
Published 20 December 2021

ABSTRACT

Background: The mouth, along with tongue, represent a site for important dermatologic disorders and could be the earliest involved site in many systemic conditions.

Objective: To delineate and evaluate the possible etiologies in patients presenting with white tongue, early or during the course of their predisposed diseases and providing a path for dealing with this condition.

Patients and Methods: In this observational descriptive study that was carried out during the period from 2013 to 2021, twenty three patients with white tongue were included. Full history and clinical, systemic and cutaneous, examination were performed aiming to delineate the specific diagnosis. Additional general investigations and skin biopsy were done when it was needed.

Results: Twenty three patients, 13 (56.5%) males and 10 (43.5%) females, suffering from white tongue were enrolled in the present study. Their age at presentation was ranged from 16-50 years with mean 37.4±9 years. Symptomatic white tongue was the presenting complaint in 19 (82.6%) patients and the symptoms ranged from mild oral discomfort to sever oral burning and soreness

Professor;

^o Dr.;

*Corresponding author: E-mail: ksharquie@ymail.com;

that interferes with oral functions. This study highlighted that lichen planus was the major disease predisposed to the white tongue in (56.52%) of patients followed by Behcets disease (13.04%), psoriasis (8.7%), oral candidiasis (8.7%), acrodermatitis enteropathica (8.7%) and pemphigus in (4.34%) cases. Associated cutaneous lesions were found in 78.3% of patients where in many of them, the rash was mild and discovered accidentally on routine examination.

Conclusion: White tongue is not uncommon complaint with possible important and serious underlying causes. The current study showed that lichen planus, Behcets disease, psoriasis, oral candidiasis, acrodermatitis enteropathica and pemphigus can manifested as white tongue either initially and/or during the course of the disease. Hence, dermatologists and oral physicians should be aware for this distinguished clinical observation.

Keywords: White tongue; lichen planus; Behcets disease; psoriasis; oral candidiasis.

1. INTRODUCTION

The mouth is a mirror reflecting health or disease. Along with tongue, the mouth represents a site for dermatologic disorders and could be the earliest involved site in many systemic conditions [1].

Normally, the tongue is pink in color and discolored tongue had been reported in many disorders. Complete or partial white coating of the tongue can be seen in furred tongue which is a common benign condition commonly observed in smokers and those with poor oral hygiene [2]. Also, in geographic tongue, mixed red and white irregular and sharply demarcated patches appeared on the dorsum of the tongue [3]. Infection, vitamin deficiency and psoriasis are some of the reported association [4].

Oral candidiasis is an opportunistic infection of the oral cavity caused by overgrowth of *Candida albicans*. It can be acute or chronic. Acute pseudomembranous candidiasis is manifested by loosely adherent, white pseudomembrane coating which can be easily scraped off with gauze while chronic hyperplastic candidiasis can manifest as whitish plaques that cannot wiped off, and are usually located on the buccal mucosa and tongue [5].

Lichen planus is a chronic disease with proposed autoimmune etiology affecting skin, hair, nail and mucosal tissues in different clinical forms [6]. The plaque-like variant can be presented as irregular, raised white plaque on the dorsum of the tongue [7]. Behçet disease is a chronic relapsing multi systemic disorder which can affects any body organ [8]. Oral aphthous ulcer covered with a white-yellow pseudo membrane that commonly occurred in buccal mucosa, and tongue, are usually the earliest sign of Behçet disease [9].

Psoriasis is a common cutaneous disease with an incidence of 2 to 3% in the general population [10]. Although oral lesions of psoriasis are matter of controversy, an association had been postulated between psoriasis with geographic, and fissured tongue [11]. Zinc deficiency can be divided into congenital so called acrodermatitis enteropathica and acquired. Orally, zinc deficiency can be manifested as erythematous, erosive and crusted lesions on the mouths and tongues [12]. Pemphigus vulgaris is a rare, chronic autoimmune disease with substantial morbidity and mortality in which easy ruptured blisters causing painful erosion and ulceration which can affect the buccal mucosa and tongue [13].

Other condition that may be presented as white tongue is oral hairy leukoplakia presenting as asymptomatic hyperkeratotic white plaque commonly occurring on lateral tongue in human immunodeficiency virus (HIV) patients and oral squamous cell carcinoma [14].

Oral manifestations may be the first presenting sign for the serious systemic and dermatologic diseases ranging from autoimmune blistering disorders with the associated substantial morbidity and mortality particularly in pemphigus vulgaris and cicatricial pemphigoid, to many chronic gastrointestinal diseases with their debilitating and chronic courses such as Crohn's disease and ulcerative colitis. In addition, the significance of muco cutaneous manifestation is quite obvious in many hematologic diseases ranging from mild anemias to more serious malignancies such as leukemia and multiple myeloma. Renal diseases, diabetes mellitus and nutritional deficiencies other than vitamin c can also present themselves in the mouth with different clinical presentation Thus, the mouth, along with mouth, provides easy accessible window for diagnosing of many systemic

diseases, the clinician should be aware about it [1].

So the aim of the present study to delineate and evaluate the possible etiologies in patient presenting with white tongue early or during the course of their predisposed diseases.

2. PATIENT AND METHODS

In this observational descriptive study, twenty-three patients suffering from white tongue were enrolled during the period from 2013-2021 years. The study followed the Declaration of Helsinki Principles and informed consent was obtained from each patient after explanation of the nature of the study. Close-up photographs were taken at the same place with constant distance and illumination. A full history for every patient covering the duration and progression of disease with the associated cutaneous and systemic diseases was obtained. In addition, Personal oral hygiene history and family history of autoimmune diseases were also reviewed with patients.

In each patient, the mouth was carefully examined with appropriate light source. When needed, tongue depressor and sterilized gauze were used for tongue manipulation and stability. Careful inspection of oral mucosal site including tongue, buccal mucosa and palate for any abnormality in form of discoloration, swelling, ulceration, and vesiculation were performed. Other mucosal site including eyes and genital mucosa, and skin, hair and nails were also involved in examination in best effort for revealing any associated lesions to reach diagnosis. The lymph nodes were also inspected and palpated. Biopsy was done in doubtful cases aiming to determine the specific diagnosis. Additionally, 10% potassium hydroxide (KOH) examination was done either to rule out or to confirm the diagnosis of oral candidiasis.

Statistical package for social science (SPSS) version 23 was used for data input and analysis. Data were statistically described in terms of mean, frequencies (no. of cases), standard deviation (SD), male to female ratio and percentage (%).

3. RESULTS

Twenty three patients, 13 (56.5%) males and 10 (43.5%) females, suffering from white tongue during eight years' time period were enrolled in the present study. Their age at presentation was

ranged from 16-50 years with mean 37.4±9 years. The frequency of the observed diseases associated with white tongue presentation in correlation with other oral mucosal and cutaneous involvement was illustrated in Table 1. Symptomatic white tongue was the presenting complaint in 19 (82.6%) patients and the symptoms ranged from mild oral discomfort with largely cosmetic impact to severe oral burning and soreness that interferes with oral functions.

Solitary white tongue discoloration without other oral mucosal lesions was observed in 17(74%) patients while in the remaining 6(26%) patients, minimal lesions distributed elsewhere in their mouths. Lichen planus was the most frequent disease presented as white tongue in 13(56.52%) patients where white tongue precedes cutaneous involvement in 7(53.8%) patients and was the sole manifestation of the lichen planus without other oral or skin involvement in 4 (30.8%) patients (Fig. 1). Scaly discolored lips and other characteristic skin involvement was found in 9 (69.2%) supporting the diagnosis of lichen planus in many patients. Behcets disease was the second cause predisposing for white tongue in 3 (13.04%) patients, the tongue was the only oral mucosal site involved in all 3 presenting patients, the other oral mucosal sites were almost clear (Fig. 2). In all 3 patients, recurrent oro-genital ulceration was documented. The average duration of Behcets disease was 5 years and the white tongue developed during the course of disease on frequent follow up of those patients.

Psoriasis as a cause of white tongue was observed in 2 (8.7%) patients (Fig. 3). Typical but accidentally discovered psoriatic lesions, while neglected by those patients, had been detected on the scalp, extensors or flexors on full skin examination yielding a path to the diagnosis. The average duration of white tongue in those patients was 3 years period and was the only oral mucosal site involved. The patient unsure which one start earlier, the cutaneous disease or the white tongue although both patients state the white tongue start from their early lives. The patients received different diagnoses and tried different pharmacologic treatment with non-satisfactory response.

Oral Candidiasis was recorded in 2 (8.7%) of the presented white tongue patients, the buccal and gingival mucosa were also involved in one patient. The full history revealed pemphigus in one patients on large dose oral prednisolone with

minimal skin lesions while the other patient, with poor oral hygiene had chronic white tongue course, no underlying diseases had been detected and KOH scraping showed pseudo hyphae.

Acrodermatitis enteropathica was recorded as a cause of white tongue discoloration in 2 (8.7%) patients (Fig. 4). The disease presented since early childhood and white tongue with angular stomatitis appeared during the course of the

disease. In addition to tongue involvement, gingivitis, alopecia, dry exerotic skin, thinning of the finger nails and psoriasiform, and burn like dermatitis lesions had been observed.

Lastly, one (4.34%) case of pemphigus as a cause of white tongue was noticed (Fig. 5). Upon careful skin examination, minute vesicle and ulceration were seen elsewhere in the skin which paid little attention by the patient. The white precedes the cutaneous manifestations.

Table 1. The frequency of white tongue among different skin diseases

Disease	Patients			Site		
	Male No.(5)	Female No.(%)	Total No.(%)	Tongue only No.(%)	Tongue +Other oral sites No.(%)	Cutaneous No.(%)
Lichen planus	6(46.2)	7(53.8)	13(56.52)	11(84.6)	2(15.4)	9(69.2)
Behcets disease	2(66.7)	1(33.3)	3 (13.04)	3(100)	none	3(100)
Psoriasis	2(100)		2 (8.7)	2(100)	none	2(100)
Candidiasis	1(50)	1(50)	2(8.7)	1(50)	1(50)	1 (50)
Acrodermatitis enteropathica	2(100)		2(8.7)		2(100)	2(100)
Pemphigus		1(100)	1(4.34)		1(100)	1(100)
Total	13(56.5)	10(43.5)	23 (100)	17(74)	6(26)	18(78.3)




Fig. 1. Fifty years old male with lichen planus showing white tongue (A) and the characteristic buccal reticular network (B)



Fig. 2. Thirty-seven years old male with Behcets disease showing white tongue and aphthous ulcers on the lateral tongue border



Fig. 3. Thirty-eight years old female with psoriasis showing white tongue



Fig. 4. Sixteen years young male with acrodermatitis enteropathica showing white tongue with periorificial dermatitis



Fig. 5. Forty-seven years old female with pemphigus vulgaris showing white tongue and crusted lip

4. DISCUSSION

For centuries, in traditional Chinese medicine, the tongue color and topography were paramount area reflecting the health status of the

individual and provide a path for diagnosis and prognosis of many diseases [15]. Thus, Examination of the tongue can provide a useful clinical evidence to several systemic diseases and local abnormalities. Hence, it is important for the clinician to be familiar with disorders affecting the tongue [2]. Normally, the dorsum of the tongue has variable appearance which is either pinkish or covered with thin white coating [16].

Too many terms in oral medicine which is not a diagnosis and only applied to search for etiologies behind, like desquamative gingivitis, erthroplakia and leukoplakia, similarly, white tongue can be applied for generalized whitish discoloration of the dorsum of the tongue representing an umbrella that cover many important and serious etiologies.

To best of our knowledge, the present study evaluating white tongue as a symptom developing early and / or during the course of predisposed diseases is the first published study with informative approach in dealing with commonest causes of not uncommon complaint in dermatology and oral medicine clinics.

In the present study, the diseases predisposed to white tongue were in the following orders: lichen planus (56.52%), Behcets disease (13.04%), psoriasis (8.7), oral candidiasis (8.7%), acrodermatitis enteropathica (8.7%) and pemphigus (4.34%).

Lichen planus is an inflammatory disorder with chronic course and suggested autoimmune etiology. It has different oral presentations with reticular pattern is the commonest type [17]. In the present study, white tongue was the only oral manifestation of lichen planus in most patients, a finding that is in contrary with many previously reported studies where the buccal mucosa and gingiva were the mostly involved sites [18,19].

Behcets disease is not uncommon in Iraqi population. It is a chronic autoimmune disorder with recurrent oral and genital ulceration, skin lesions inform of erythema nodosum and acneform lesions, and eye lesions inform of uveitis and hypopyon, with ability to affect many organ systems including cardiovascular, nervous and gastrointestinal systems [20,21]. The mainly reported oral manifestation of Behcets disease is the oral aphthous ulcer covered with a white-yellow pseudo membrane that commonly occurred in buccal mucosa, and tongue [9]. As far as we know, white tongue developing early or

during the course of the Behcets disease represents a distinct manifestation that had never been described in previous studies in which the patients present with complete whitish discoloration of the dorsum of the tongue, this finding should be kept in mind when evaluating a patient with white tongue.

Psoriasis is a common skin disorder with heterogeneous clinical presentations and predisposing factors [22]. Rare cases of intraoral psoriasis had been reported. The first case of biopsy proven oral psoriasis was diagnosed by Oppenheim in 1903 [23]. The oral manifestations of psoriasis are arguable subject because many common oral inflammatory disorders have similar clinico-pathologic criteria. Also, it may appear as coincidental finding [24]. In 2 controlled studies evaluating tongue lesions in 200 and 207 psoriatic patients from 2 different ethnic groups, fissured tongue and benign migratory glossitis were the two most commonly recorded findings [24,25]. None of these studies reported white tongue as oral manifestation of psoriasis. Oral psoriasis presented as persistent whitish lesion of the tongue had been reported in association with cutaneous lesions in single case report [26]. So, the current study presents the white tongue as unique intra oral psoriatic lesions that should be looked for and added to other oral manifestation of psoriasis.

In the present study, oral candidiasis was observed as a cause of white tongue in 2 patients. This finding correlates with other report as oral candidiasis considered as a well-recognized cause of white coated tongue in both acute pseudomembranous and chronic hyperplastic type [5]. As many oral diseases can be complicated by oral Candidiasis, either due to immune suppressive treatment or due to the associated pain while trying proper oral hygiene, the diagnosis of oral candidiasis was made by exclusion with careful history and examination that reveal one of the presenting patients had pemphigus who continues to receive large dose of prednisolone that predispose the patient to candidal infection in agreement with another report [5].

Acrodermatitis enteropathica is rare, congenital disorder inherited in autosomal dominant pattern caused by impaired absorption of zinc characterized by scaly crusted plaque with ulceration in acral and periorificial distribution, alopecia and nail abnormalities [27]. In the present study, burn like dermatitis, that observed

in the acrodermatitis enteropathica patients, had also been described in association with zinc deficiency in earlier study by sharquie et al. [28]. Although zinc deficiency was found to be associated with many oral problems such as recurrent aphthous stomatitis, burning mouth syndrome, atrophic glossitis, xerostomia and, erythematous, erosive and crusted lesions on the mouth and tongue [29] but none of the available studies described white tongue in acrodermatitis enteropathica. Thus, this study can send a message that white tongue can be a feature of acrodermatitis enteropathica even if it's not early presentation but could be developed later and could be sign of noncompliance to treatment.

Well known presentation of pemphigus vulgaris is painful and slow to heal buccal or palatal mucosal erosion while intact blisters are difficult to be found as the blisters are flaccid and easily ruptured [30]. Many rare presentations of pemphigus vulgaris had been described in the literatures including multiple oral pustules involving the tongue, buccal mucosa, palate, and the gingiva, [31] multiple cutaneous pustules [32] and alopecia [33]. White tongue as presentation of oral pemphigus vulgaris represent a unique manifestation that should alert the physician for this serious disorder.

The present work had introduced the tongue as a mirror reflecting serious dermatologic disorders. With exception of oral candidiasis and, the white tongue should be considered as new descriptive term similar to many other non-diagnostic term and the physician should be able to specify the etiologies gone behind it.

5. CONCLUSION

White tongue is not uncommon problem in dermatology clinic with little emphasis on its etiologies. The present study recorded that lichen planus, Behcets disease, psoriasis, oral candidiasis, acrodermatitis enteropathica, and pemphigus were the main responsible disorders causing white tongue. It is important for both dermatologist and oral physician to be carefully looked for and analyzed the predisposing disease for white tongue.

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mehrotra V, Devi P, Bhovi TV, Jyoti B. Mouth as a mirror of systemic diseases. *Gomal J Med Sci.* 2010;8:235-41.
2. Rogers RS 3rd, Bruce AJ. The tongue in clinical diagnosis. *J Eur Acad Dermatol Venereol.* 2004;18(3):254-9.
3. Joseph BK, Savage NW. Tongue pathology. *Clin Dermatol.* 2000;18(5):613-8.
4. Vahedi M, Abdolsamadi HR, Mortazavi H, Abdollahzadeh SH. Evaluation of the therapeutic effects of zinc sulfate in patients with geographic tongue. *Avicenna J Dent Res.* 2009;1(1):11– 4.
5. Sharma A. Oral candidiasis: An opportunistic infection: A review. *Int J Appl Dent Sci.* 2019;5(1):23-27.
6. Sharquie KE, Noaimi AA, Zeena AA. Upsurge of Cases of Lichen Planus in Iraqi Population in Baghdad City with Frequency of Hepatitis Viruses. *IOSR J Dent Med Sci.* 2015;14(12):78–81.
7. Edwards PC, Kelsch R. Oral lichen planus: Clinical presentation and management. *J Can Dent Assoc.* 2002;68(8):494-9.
8. Eguía A, Villarroel M, Martínez-Conde R, Echebarría MA, Aguirre JM. Adamantiades-Behçet disease: An enigmatic process with oral manifestations. *Med Oral Patol Oral Cir Bucal.* 2006; 11:E6-11.
9. Bulur I, Onder M. Behçet disease: New aspects. *Clin Dermatol.* 2017;35(5):421-434.
10. Al-Hamamy HR, Al-Turfy IA, Abdul-Reda FS. Infliximab therapy in iraqi patients with moderate to severe psoriasis. *Journal of Cosmetics, Dermatological Sciences and Applications.* 2015;5:78-85.
11. Suliman NM, Åstrøm AN, Ali RW, Salman H, Johannessen AC. Oral mucosal lesions in skin diseased patients attending a dermatologic clinic: A cross-sectional study in Sudan. *BMC Oral Health.* 2011;11:24.
12. Kumar S, Thakur V, Choudhary R, Vinay K . Acrodermatitis Enteropathica. *J Pediatr.* 2020;220:258-259.
13. Dagistan S, Goregen M, Miloglu O, Cakur B. Oral *Pemphigus vulgaris*: A case report with review of the literature. *J Oral Sci.* 2008;50(3):359-62.
14. Mangold AR, Torgerson RR, Rogers RS 3rd. Diseases of the tongue. *Clin Dermatol.* 2016;34(4):458-69.
15. Anastasi JK, Chang M, Quinn J, Capili B. Tongue Inspection in TCM: Observations in a Study Sample of Patients Living with HIV. *Med Acupunct.* 2014;26(1):15-22.
16. Danser MM, Gómez SM, Van der Weijden GA. Tongue coating and tongue brushing: a literature review. *Int J Dent Hyg.* 2003; 1(3):151-8.
17. Mollaoglu N. Oral lichen planus: A review. *Br J Oral Maxillofac Surg.* 2000;38(4):370-7.
18. Tak MM, Chalkoo AH. Demographic, Clinical Profile of Oral Lichen Planus and its Possible Correlation with Thyroid Disorders: A Case-Control Study. *Int J Sci Stud.* 2015;3(8):19-23.
19. Kaushik D, Gupta B. Demographic and clinical profile of oral lichen planus in patients attending a tertiary care hospital in North East India. *IP Indian J Clin Exp Dermatol.* 2021;7(1):24-29.
20. Sharquie KE, Najim RA, Al-Dori WS, Al-Hayani RK. Oral zinc sulfate in the treatment of Behcet's disease: a double blind cross-over study. *J Dermatol.* 2006 ;33(8):541-6.
21. Al-Rawi ZS, Sharquie KE, Khalifa SJ, Al-Hadithi FM, Munir JJ. Behçet's disease in Iraqi patients. *Ann Rheum Dis.* 1986; 45(12):987-90.
22. Sharquie, K.E., Noaimi, A.A. and Alobaidi, M.H. A New Regimen in the Treatment of Psoriasis Using Oral Methotrexate. *Journal of Cosmetics, Dermatological Sciences and Applications.* 2019;9:165-176.
23. Brice DM, Danesh-Meyer MJ. Oral lesions in patients with psoriasis: Clinical presentation and management. *J Periodontol.* 2000;71(12):1896-903.
24. Hernández-Pérez F, Jaimes-Avelaño A, Urquiza-Ruvalcaba Mde L, Díaz-Barcelot M, Irigoyen-Camacho ME, Vega-Memije ME, Mosqueda-Taylor A. Prevalence of oral lesions in patients with psoriasis. *Med*

- Oral Patol Oral Cir Bucal. 2008;13(11): E703-8.
25. Daneshpazhooh M, Moslehi H, Akhyani M, Etesami M. Tongue lesions in psoriasis: A controlled study. BMC Dermatol. 2004; 4(1):16.
 26. Lier GC, Mrowietz U, Wolfart M, Warnke PH, Wiltfang J, Springer IN. Psoriasis of the tongue. J Craniomaxillofac Surg. 2009; 37(1):51-3.
 27. Perafán-Riveros C, França LF, Alves AC, Sanches JA Jr. Acrodermatitis enteropathica: Case report and review of the literature. Pediatr Dermatol. 2002; 19(5):426-31.
 28. Sharquie KE, Al-Jaralla FA. Burn- like dermatoses: A neglected group of skin diseases with a new classification. Am J Derm Vener. 2019;8:88-93.
 29. Uwitonze AM, Ojeh N, Murererehe J, Atfi A, Razzaque MS. Zinc adequacy is essential for the maintenance of optimal oral health. Nutrients. 2020;12(4):949.
 30. Niyaz A, Viswanath BK. Atypical presentation of pemphigus vulgaris - A rare case report. Our Dermatol Online. 2019; 10(1):88-90.
 31. Rimal J, Sumanth KN, Ongole R, George T, Chatterjee S. A rare presentation of oral pemphigus vulgaris as multiple pustules. Kathmandu Univ Med J (KUMJ). 2007; 5(4):541-5.
 32. Yang Y, Lin M, Huang SJ, Min C, Liao WQ. A rare presentation of pemphigus vulgaris as multiple pustules. Indian J Dermatol. 2010;55(3):293-5.
 33. Veraitch O, Ohyama M, Yamagami J, Amagai M. Alopecia as a rare but distinct manifestation of pemphigus vulgaris. J Eur Acad Dermatol Venereol. 2013;27(1):86-91.

© 2021 Sharquie et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78170>*