



Assessing the Impact of Indole Butyric Acid Application & Varietal Difference on Various Parameters of Hardwood Cutting in Pomegranate (*Punica granatum L.*) Varieties

Harshit Jangid^{a++*}, Jyoti Kanwar^{b#}, Ratnapriya Tiwari^{c†},
Sunil Khandoliya^{d†} and Jitendra Gurjar^{e†}

^a Department of Horticulture (Fruit Science), KNK College of Horticulture, Mandsaur. Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, Madhya Pradesh, India.

^b Department of Fruit Science, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, KNK College of Horticulture, Mandsaur, Madhya Pradesh, India.

^c Department of Horticulture (Fruit Science), Mahatma Gandhi University of Horticulture and Forestry, Patan, Durg, Chhattisgarh, India.

^d Department of Horticulture (Fruit Science), Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, Madhya Pradesh, India.

^e Department of Horticulture, SKRAU, Bikaner, Rajasthan, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i71044>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115234>

Original Research Article

Received: 01/02/2024

Accepted: 02/04/2024

Published: 22/06/2024

++ MSc;

Assistant Professor;

† PhD Scholar;

*Corresponding author: E-mail: hjangid771@gmail.com;

Cite as: Jangid, Harshit, Jyoti Kanwar, Ratnapriya Tiwari, Sunil Khandoliya, and Jitendra Gurjar. 2024. "Assessing the Impact of Indole Butyric Acid Application & Varietal Difference on Various Parameters of Hardwood Cutting in Pomegranate (*Punica Granatum L.*) Varieties". *Journal of Advances in Biology & Biotechnology* 27 (7):843-52. <https://doi.org/10.9734/jabb/2024/v27i71044>.

ABSTRACT

Aim: To determine the best combination of Indole-3-butyric acid & Variety for better growth of pomegranate cuttings.

Place and Duration of Study: Experiment was carried out during the year 2022-23 at the Instructional cum Research Department of Fruit Science, RVSKVV- K.N.K. College of Horticulture, Mandasaur (M.P).

Study Design: The experiment was laid out in Factorial Completely Randomized Design (FCRD) comprising of seven varieties hardwood cuttings (Mridula, Phule Arakta, Saharanpur, Jodhpur Red, Jalore Seedless, G-137, Bhagwa) and two IBA concentrations (3000 & 2500 ppm), replicated thrice.

Result: The interaction of cultivar and growth regulator significantly affected Days taken to 50% sprouting (15.34 Day), Number of shoots per cutting (9.88 & 10.43), Length of leaves (3.83 & 4.98 cm), width of leaves (1.76 & 2.10 cm) at 45 & 60 DAP respectively was recorded maximum value in Bhagwa cutting treated with 3000 ppm IBA, While SPAD Value observe non-significant.

Conclusion: The current study found that different dosages of IBA and different varieties cuttings significantly influenced in different parameters. Bhagwa treated with 3000 ppm IBA is best in different shoot parameters at both 45 and 60 DAP.

Keywords: Hardwood cuttings; indole butyric acid (IBA); *punica granatum L*; Varieties; apple Pomegranate juice.

1. INTRODUCTION

Pomegranate, scientifically known as *Punica granatum L.* belong to Punicaceae family. It thrives in arid and semiarid climates and is a highly popular fruit crop worldwide. The pomegranate tree is deciduous in subtropical regions and evergreen in tropical regions Ozugan [1]. It is believed to have originated in Iran and has been cultivated since ancient times.

"Pomegranate" is a Latin word that means "apple with numerous seeds". According to Smith [2] *Punica granatum* has $2n = 16$ or 18 chromosomes. The pomegranate is a naturally dense, deciduous, bushy, multi-stemmed shrub that typically grows to heights of 10 to 12 feet and produces highly coloured fruit filled with many juicy seeds. In most regions, pomegranates are trained into a multi-stem system, with slender, somewhat thorny branches and glossy, dark green leaves. The tree produces colourful orange-red flowers in the spring and summer, which are either bell-shaped (female) or vase-shaped (hermaphroditic), with the latter type being sterile. The edible part of the fruit, called an aril, consists of hundreds of seeds enclosed by juicy pigments, each surrounded by a seed coat Smith, [2].

It has long been a symbol of wealth and ambition [3]. Pomegranate juice has numerous benefits for those suffering from leprosy. In Azerbaijan, citric

acid extract and sodium citrate are derived from pomegranate juice. Additionally, the barks and skins of the fruit are commonly utilized for treating dysentery, diarrhea, and other digestive issues Gil et al. [4].

The pomegranate has been recognized for centuries as a potent food medicine with a diverse range of clinical applications. According to Gil et al. [4], pomegranate juice has an antioxidant capacity three times greater than red wine or green tea and 2-6 times higher than grape, grapefruit and orange juices, when tested individually [5]. Pomegranate has gained popularity due to its health-promoting qualities, including its antimutagenic, antiatherogenic, antioxidant, antihypertensive properties, and high anthocyanin content [6].

Pomegranate can be propagated through both sexual and asexual means. Cheema et al. (1954) and Nagpal [7] have recommended asexual or vegetative methods of propagation to eliminate the high degree of variability observed in seed plantations. Pomegranate is commonly propagated for commercial purposes using hardwood stem cuttings. This method is considered the most convenient and can produce a fully developed and strong plant in significantly less time. The most commonly used growth substances for better rooting of various plant parts are IAA, IBA, and NAA. Of these, IBA is preferred over the others [8-14].

The ability of cuttings to root varies depending on the cultivar, location, season, and age of the branch used. The success of pomegranate cuttings depends on various factors, such as the conditions of the mother plants, the part of the tree from which the cuttings are taken, the timing of the operation, rainfall, temperature fluctuations, and aftercare. Moreover, different environmental conditions and growth regulators also play an essential role in the rooting and growth of pomegranate cuttings. The main aim of this study is to access the effect of combination of IBA and different varieties on different parameters of pomegranate hardwood cuttings Rao et al, Tanwar et al [14,15].

2. MATERIALS AND METHODS

2.1 Experiment Detail

The experiment was laid out in factorial complete randomized design with a IBA and different varieties of pomegranate as fourteen treatment and three replications. Treatment combination are as under:

2.2 Growth Regulator Formulation

The required concentration of indole butyric acid (IBA) 2500 & 3000 ppm was prepared by dissolving 2.5 & 3.0 g IBA in 10 ml of ethyl alcohol (90%) and final volume was made to 1000 ml using distilled water.

2.3 Study Location

Study was conducted during the year 2022-2023 at the Experimental Field of Department of Fruit Science, KNK College of Horticulture, Mandsaur (MP). Mandsaur is situated at 23.450 to 24.130 N latitude and 74.440 to 75.180 E longitudes at an altitude of 435 m mean sea level. Mandsaur temperature rises up to 46 °C during summer and falls to 3.6 °C during winter with an occasional

occurrence of frost. The average annual rainfall is 797.6 mm, most of which occurred during July to September, winter and summer rains are uncommon.

2.4 Experimental Data Collection

Days Taken to 50% Sprouting: The days taken to 50% sprouting of total cutting were counted from date of planting till the 50% sprout comes out of total cutting planted and recorded as days taken to 50% sprouting.

Number of shoots per cutting at 45 and 60 DAP: The number of shoots per cutting was recorded randomly selected plants in each treatment and replication at 45 and 60 DAP.

Length of longest shoot (cm) at 45 and 60 DAP: The length of the longest shoot was measured on each sample of cutting and mean was recorded after 45 and 60 days after planting of cuttings.

Length of leaf (cm) at 45 and 60 DAP: Length in (cm) under each treatment was recorded after 45 and 60 days of planting with the help of scale and mean length of leaves per cutting was worked out.

Width of leaf (cm) at 45 and 60 DAP: width in (cm) under each treatment was recorded after 45 and 60 days of planting with the help of scale and mean width of leaves per cutting was worked out.

SPAD value at 45 and 60 DAG: The cuttings were randomly selected to measure the total chlorophyll content using a digital chlorophyll meter (SPAD-502). The mean value of the leaves was calculated, and the results were expressed in SPAD units (an acronym for Soil Plant Analysis Division).

List 1. Treatment combinations

T ₁ - Mridula + IBA 2500 ppm	T ₈ - Mridula + IBA 3000 ppm
T ₂ - Phule Arakta + IBA 2500 ppm	T ₉ - Phule Arakta + IBA 3000 ppm
T ₃ - Saharanpur + IBA 2500 ppm	T ₁₀ - Saharanpur + IBA 3000 ppm
T ₄ - Jodhpur Red + IBA 2500 ppm	T ₁₁ - Jodhpur Red + IBA 3000 ppm
T ₅ - Jalore seedless+ IBA 2500 ppm	T ₁₂ - Jalore Seedless+ IBA 3000 ppm l ₂
T ₆ - G-137+ IBA 2500 ppm	T ₁₃ - G-137+ IBA 3000 ppm
T ₇ - Bhagwa + IBA 2500 ppm	T ₁₄ - Bhagwa + IBA 3000 ppm

3. RESULTS AND DISCUSSION

3.1 Days Taken to 50% Sprouting

Among seven distinct varieties, results showed that cultivars had considerable variations in days taken to 50% sprouting. Pomegranate cv. Bhagwa was shown to be the best variety, with the minimum days to 50% sprouting (15.47 days) than cvs. Mridula (15.88 days), Phule Arakta (16.60 days), Jodhpur Red (17.53 days), G137 (18.53 days), Jalore seedless (19.42 days) as opposed to the Saharanpur, which had the maximum days to 50% sprouting (20.10 days). In case of effect of different levels of growth regulator, it was seen that the earliest 50% sprouting (17.14 days) was observed with 3000 ppm IBA, whereas, IBA 2500 ppm took (18.15 days) days to 50 % of sprouting.

As regards different varieties and levels of growth regulator were reported significant to this attribute. However, the minimum days to 50 % of sprouting was reported in Bhagwa (15.34 days) treated with 3000 ppm followed by Mridula (15.35 days) treated with 3000 ppm and Bhagwa (15.60 days) treated with 2500 ppm. Whereas, maximum days (20.50) taken to fifty per cent sprouting of cuttings were recorded in Saharanpur treated with 2500 ppm IBA.

The minimum number of days required for first sprouting (12.97 days), minimum days taken to 50 % of sprouting (15.34) and greatest number of cuttings sprouted (7.67) found in Bhagwa treated with IBA 3000 ppm. Better and earlier sprouting might be due to an accumulation of endogenous growth stimulating chemicals present in the

tissue of the cuttings reported by Kaur et al. [16] in pomegranate cv. Bhagwa. The minimum number of days took to 50 % sprouting, it might be due to the higher level of auxins resulted in the completion of physiological processes involved cutting roots and sprouting sooner. Similarly, Gautheret, Haissing [17,18] observed that the first root initial cell division is dependent on either administered or endogenous auxin.

3.2 Number of Shoots Per Cutting

Among the different varieties tested, variety Bhagwa exhibited the maximum number of shoots at 45 and 60 DAP, with values of 9.79 & 10.17 respectively then Mridula (9.26 & 9.68), Phule Arakta (8.87 & 9.14), Jodhpur Red (8.08 & 8.59), G-137 (4.24 & 8.21), Jalore Seedless (3.97 & 7.88), Saharanpur (3.92 & 7.69) at 45 and 60 DAP was recorded. In concentration of IBA maximum number of shoot (8.68 & 9.10) at 45 and 60 DAP respectively was observed in 3000 ppm IBA concentration while the minimum number of shoots per cuttings (7.96 & 8.43) at 45 and 60 DAP was recorded under 2500 ppm IBA.

The combining impact of varieties and IBA concentrations indicated significant difference in number of shoots per cutting at 45 and 60 DAP. The maximum number of shoot per cuttings (9.88 & 10.43) at 45 and 60 DAP in Bhagwa treated with 3000 ppm IBA concentration followed by Mridula (9.83 & 10.18) treated with 3000 ppm IBA, at 45 and 60 DAP respectively. While, Saharanpur (6.91 & 7.57) treated with 2500 ppm IBA recorded minimum number of shoots per cuttings at 45 & 60 DAP respectively.

Table 1. Effect of varieties and IBA concentrations on Day taken to 50% sprouting in pomegranate cuttings

Variety (V)	IBA concentrations (I)		
	2500 ppm	3000 ppm	Mean
Mridula	16.4	15.35	15.88
Phule Arakta	17.1	16.1	16.6
Saharanpur	20.5	19.7	20.1
Jodhpur Red	18.1	16.95	17.53
Jalore Seedless	20.12	18.73	19.42
G 137	19.25	17.8	18.53
Bhagwa	15.6	15.34	15.47
Mean	18.15	17.14	
	S.Em.±	C.D. @ 5%	
V	0.0723	0.209	
I	0.0386	0.112	
VxI	0.102	0.296	




Fig. 1. Effect of varieties & IBA concentration on Day taken to 50% sprouting




Fig. 2. Effect of Varieties & IBA concentration on Number of shoots per cuttings

Table 2. Effect of varieties and IBA concentrations on number of shoots per cutting in pomegranate cuttings

Variety (V)	IBA concentration (I)					
	45 DAP			60 DAP		
	2500 ppm	3000 ppm	Mean	2500 ppm	3000 ppm	Mean
Mridula	8.68	9.83	9.26	9.18	10.18	9.68
Phule Arakta	8.08	9.67	8.87	8.50	9.78	9.14
Saharanpur	6.91	7.31	7.11	7.57	7.81	7.69
Jodhpur Red	7.74	8.41	8.08	8.24	8.93	8.59
Jalore Seedless	7.11	7.68	7.40	7.61	8.16	7.88
G 137	7.51	8.01	7.76	8.01	8.40	8.21
Bhagwa	9.70	9.88	9.79	9.90	10.43	10.17
Mean	7.96	8.68		8.43	9.10	
	S.Em.±	C.D. @ 5 %		S.Em.±	C.D. @ 5%	
V	0.0689	0.199		0.1150	0.333	
I	0.0368	0.107		0.0615	0.178	
Vxl	0.097	0.282		0.163	0.471	

Table 3. Effect of varieties and IBA concentrations on length of leaves in pomegranate cuttings

Variety (V)	IBA concentration (I)					
	45 DAP			60 DAP		
	2500 ppm	3000 ppm	Mean	2500 ppm	3000 ppm	Mean
Mridula	3.58	3.78	3.68	4.71	4.91	4.81
Phule Arakta	3.18	3.65	3.42	4.31	4.81	4.56
Saharanpur	2.20	2.58	2.39	3.30	3.71	3.51
Jodhpur Red	3.07	3.33	3.20	4.20	4.46	4.33
Jalore Seedless	2.43	2.88	2.66	3.56	4.01	3.79
G 137	2.83	3.08	2.96	3.96	4.21	4.09
Bhagwa	3.78	3.83	3.81	4.91	4.98	4.95
Mean	3.01	3.31		4.14	4.44	
	S.Em.±	C.D. @ 5 %		S.Em.±	C.D. @ 5 %	
V	0.0442	0.128		0.0405	0.117	
I	0.0236	0.068		0.0217	0.063	
VxI	0.062	0.181		0.057	0.166	

It was observed that cuttings of cv. Bhagwa treated with IBA 3000 ppm sprouted significantly more number of shoots (10.43) compared to remaining treatments. The highest number of shoot development during higher auxin treatment may be due to reason that auxins support in the establishment of a strong root system, which may have improved nutrient absorption under the effect of IBA [19]. Ghosh et al., [20] also reported that the use of IBA and NAA may have caused an increase in the number of shoots per cutting.

3.3 Length of Leaves (cm)

Among varieties, the maximum length of leaves formed on the shoot of Bhagwa (3.81 & 4.95 cm) at 45 & 60 DAP was reported significantly higher than remaining cultivars. It was followed by Mridula (3.68 & 4.81 cm), Phule Arakta (3.42 & 4.56 cm), Jodhpur Red (3.20 & 4.33 cm), G-137 (2.96 & 4.09 cm), Jalore Seedless (2.66 & 3.79 cm) while, least length of leaves was reported in cv. Saharanpur (2.39 & 3.51 cm) at 45 & 60 DAP.

A significant relationship between various cultivar and growth regulator interactions has been identified. IBA 3000 ppm applied to Bhagwa (3.83 & 4.98 cm) had the longest leaf followed by Mridula (3.78 & 4.91 cm). In contrast, Saharanpur treated with 2500 ppm IBA had the shortest length (2.2 & 3.30 cm) at 45 & 60 DAP respectively.

Cuttings of Bhagwa treated with 3000 ppm produced significantly maximum number of leaves (35.50), highest leaf length (4.98 cm), maximum width (2.10 cm) and maximum leaf area (10.10 cm²). The growth regulators' ability to stimulate strong roots system, which helps

to cuttings absorb more water and nutrients and produce more leaves may be the cause of the increase in leaf count. Additionally, as auxins encourage the growth of primary shoots and their quantity, the number of leaves may also have raised [21]. The findings match those declared public by Singh [22] in pomegranate, Dahale et al. [23] in Fig fruit plants, Ghosh et al. [20] in Phalsa and Mehta et al. [24] in pomegranate.

4. WIDTH OF LEAVES (CM)

It is evident from the data that Bhagwa, which greatly outgrew in terms of width of leaves (1.74 & 2.07 cm) followed by Mridula (1.71 & 2.03 cm), Phule Arakta (1.63 & 1.95 cm), Jodhpur Red (1.52 & 1.87 cm), G137 (1.40 & 1.74 cm) and Jalore Seedless (1.31 & 1.63 cm). The smallest leaf width (1.27 & 1.59 cm) was discovered in Saharanpur. IBA at 3000 ppm was shown to be much superior than other levels of growth regulator in the case of growth regulator, as seen by the greatest breadth (1.57 & 1.90 cm) of leaf that could be recorded with IBA 3000. While IBA at 2500 ppm was used to measure the minimum breadth of leaves (1.46 & 1.78 cm) at 45 & 60 DAP respectively.

It was discovered that there was a considerable interaction between different varieties and growth regulators. However, Bhagwa (1.76 & 2.10 cm) treated with 3000 ppm IBA and Mridula (1.72 & 2.04 cm) treated with 3000 ppm IBA were found to have the widest leaves respectively 45 & 60 DAP. In contrast, IBA 2500 ppm was found to have the smallest breadth (1.22 & 1.55 cm) in Saharanpur at 45 & 60 DAP respectively.

Table 4. Effect of varieties and IBA concentrations on Width of leaves in pomegranate cuttings

Variety (V)	IBA concentration (I)					
	45 DAP			60 DAP		
	2500 ppm	3000 ppm	Mean	2500 ppm	3000 ppm	Mean
Mridula	1.70	1.72	1.71	2.02	2.04	2.03
Phule Arakta	1.54	1.71	1.63	1.86	2.03	1.95
Saharanpur	1.22	1.31	1.27	1.55	1.63	1.59
Jodhpur Red	1.42	1.62	1.52	1.76	1.97	1.87
Jalore Seedless	1.23	1.39	1.31	1.55	1.71	1.63
G 137	1.35	1.45	1.40	1.67	1.81	1.74
Bhagwa	1.72	1.76	1.74	2.04	2.10	2.07
Mean	1.46	1.57		1.78	1.90	
	S.Em.±	C.D. @ 5%		S.Em.±	C.D. @ 5%	
V	0.0130	0.038		0.0181	0.053	
I	0.0070	0.020		0.0097	0.028	
VxI	0.018	0.053		0.026	0.074	




Fig. 3. Effect of varieties & IBA concentration on Width & length of leaves

Greater leaf size may have been the result of the plant's rapid growth as a result of the greater auxin content. Kaur and Kaur [25] in fig and Patel et al. [26] in Kagzi lime reported findings that were comparable.

5. CHLOROPHYLL OR SPAD VALUE

The findings showed that there were considerable variations in total chlorophyll concentration among cultivars. Bhagwa cuttings outperformed the other cultivars in terms of chlorophyll concentration per cutting (27.87) next to Mridula (27.80), Phule Arakta (26.51), Jodhpur Red (25.72), G 137 (25.11), Jalore Seedless (24.44) at 45 DAP. While, Mridula (31.10) outperformed in terms of chlorophyll concentration next to Bhagwa (30.37), G 137 (30.11), Phule Arakta (29.32), Jodhpur Red

(28.84) and then Jalore Seedless (27.61) at 60 DAP. The minimum chlorophyll concentration was found in cv. Saharanpur (23.95 & 27.50) at 45 & 60 DAP. Regardless of cultivar, IBA 3000 ppm (26.55 & 29.79) outperformed on IBA 2500 ppm (25.28 & 28.73) at 45 & 60 DAP respectively.

There was no significant interaction between cultivars and IBA concentrations on total leaf chlorophyll amount per.

The increased concentration of auxins has been found to stimulate leaf growth, resulting in a higher leaf area. This, in turn, activates more photosynthates, leading to an increased chlorophyll content in the leaves of cuttings. Growth hormones, as demonstrated by Galston and Davies in 1969 [27], play a crucial role in

Table 5. Effect of varieties and IBA concentrations on SPAD value in pomegranate cuttings

Variety (V)	IBA concentration (I)					
	45 DAP			60 DAP		
	2500 ppm	3000 ppm	Mean	2500 ppm	3000 ppm	Mean
Mridula	26.62	28.99	27.80	30.27	31.93	31.10
Phule Arakta	25.59	27.44	26.51	28.71	29.92	29.32
Saharanpur	23.82	24.08	23.95	27.86	27.13	27.50
Jodhpur Red	25.29	26.14	25.72	28.41	29.26	28.84
Jalore Seedless	23.78	25.10	24.44	27.00	28.22	27.61
G 137	24.74	25.48	25.11	29.26	30.96	30.11
Bhagwa	27.15	28.59	27.87	29.61	31.13	30.37
Mean	24.97	26.55		28.73	29.79	
	S.Em.±	CD_(0.05)		S.Em.±	CD_(0.05)	
V	0.2298	0.666		0.3382	0.980	
I	0.1228	0.356		0.1807	0.524	
VxI	0.325	0.941		0.478	1.385	




Fig. 4. Effect of Varieties & IBA on SPAD Value

regulating the distribution and quantity of assimilates in plants. Sahab et al. in 2013 also observed that cuttings with a greater number of leaves had improved nutrient uptake, leading to increased production of photosynthates and providing an ample food supply for the metabolic activities of the plants. These findings align with the results of Kumari's study in 2014 [28], which focused on pomegranate propagation in the cv. Ganesh. The results of this study were found to be closer to the findings of Kaur et al. [16] in grapevine and Hakim et al. [29] in pomegranate that raising IBA concentration improves total chlorophyll content, Ghani et al. [30] in pomegranate cv. Phule Bhagwa Super [31-35].

6. CONCLUSION

The current study found that different dosages of IBA and different varieties cutting significantly

influenced properties of pomegranate cuttings. The result revealed that among all the treatments Bhagwa treated with 3000 ppm best in 50% sprouting, width & length of leaves, number of shoot per cutting, Length of longest shoot of cuttings. Out of 7 cultivars investigated, cvs. Bhagwa, Mridula, and Phule Arakta performed best in respect of all root and shoot parameters, followed by cvs. Jodhpur Red, G 137, Jalore Seedless, and Saharanpur. In practically all root and shoot characteristics, IBA @ 3000 ppm outperformed followed by 2500 ppm.

ACKNOWLEDGEMENT

This research would not have been possible without the support and guidance of several individuals and institutions. We would like to express our sincere gratitude to teachers and colleagues their invaluable guidance,

encouragement, and expertise throughout the research process. Their critical feedback and constructive suggestions significantly improved the quality of this work. I also express my deep feeling of affection and love to all my family members for their everlasting love, day by day help and bearing of hardship without which this would not have been reality.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ozugan AI. World production status of pomegranate (*Punica granatum L.*). In: 2nd MESFIN Meeting on Plant Genetic Resources, Madeira, Portugal. 1997; -5-8.
2. Smith PM. Minor crops, In Evolution of Crop Plants (Simmonds, N.W-ed.), Longman, London, 1976;301-324.
3. Duman AD, Ozgen M, Dayisoylu KS, Erbil N and Durgac C. Antimicrobial activity of six pomegranate (*Punica granatum L.*) varieties and their relation to some of their pomological and phytonutrient characteristics. *Molecules*. 2009;14:808-1817.
4. Gil MI, Tomas-Barberan FA, Hess-Pierce B, Holcroft DM, Kader AA. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *J. Agric. Food Chem.* 2000;48(10) 4581-4589.
5. Rosenblat M, Aviram M. Antioxidative properties of pomegranate: In vitro studies. In: Seeram NP, Heber D. (eds.) *Pomegranate: Ancient roots to modern medicine*. 2006;31-43.
6. Basu A, Penugonda K. Pomegranate Juice: A heart healthy fruit juice. *Nutritional Reviews*, 2009; 67(1): 49-56.
7. Nagpal RL. Pomegranate Cultivation in India. *Farm Bull. ICAR*, New Delhi. 1954;22.
8. Avery GS, Johnson EB. Kind of Hormones that stimulate rooting. *Ited Honnonnes and Horticulture*. 1947;13-15.
9. Pearse HL. Experiment with growth promoting substances-I. Reaction leafless cuttings to treatment with root promoting substance. *Anni. Bot. New Series-II*. 1948;5:1938.
10. Garnner RJ. Propagation by cuttings and layers. Recent work and its application with special reference to preen and stone fruits. *Tech. Common. Bur. Hort. Meeting.*, 14:19-20. *Indian Hort.* 1958;19(2):1962-720.
11. Leopold AC. Auxin and plant growth. McGraw Hill Book Co., New York. 1958;203.
12. Loreti F, Hartman HT. Propagation of Olive tree by rooting leafy cuttings under mist. *Proc. Am. Soc. Hort. Sci.* 1962;85:257-264.
13. Sarma SD, Sarma VK. Rooting pattern of hard wood and semi hard wood cuttings of wild pomegranate. *Indian J Hort.* 1991;44:188-193.
14. Rao GS, Bisati IA, Shirma A, Kosser S & Bhat SA. Effect of IBA concentration and cultivars on number of leaves, leaf area and chlorophyll content of leaf in Pomegranate (*Punica granatum L.*) cuttings under temperate conditions of Kashmir. *Journal of Pharmacognosy and Phytochemistry* 2020;SP6:86-90.
15. Tanwar DR, Bairwa HL, Lakhawat SS, Mahawer LN, Jat RK, Choudhary RC. Effect of IBA and rooting media on hardwood cuttings of Pomegranate (*Punica granatum L.*) CV. Bhagwa. *International Journal of Environment and Climate Change*. 2020;10(12):609-17.
16. Kaur SS, Cheema B, Chhabra R, Talwar KK. Chemical induction of physiological changes during adventitious root formation and bud break in grapevine cuttings. *Plant Growth regulators*. 2002;37(1): 63 – 68.
17. Gautheret RJ. Investigation on the root formation in the tissues of *Helianthus tuberosus* cultured in vitro. *Amer. J. Bot.* 1969;56(7):702-712.
18. Hassing BE. Meristematic activity during adventitious root primordial development. Influence of endogenous auxin and applied gibberellic acid. *Plant Physiol.* 1972;49:886-892.
19. Hiral, Patel R, Patel MJ, Singh smith. Effect Of Different Levels of Iba and naa on rooting of Hardwood and Semi Hardwood Cutting in Fig. *International Journal of Agricultural Science and Research (IJASR)*. 2017;7(4):2321-0087.
20. Ghosh A, Dey K, Mani A, Bauri FK, Mishra DK. Efficacy of different levels of IBA and NAA on rooting of Phalsa (*Grewia asiatica L.*) cuttings. *International Journal of Chemical Studies*. 2017;5(6):567-571
21. Khapare LS, Dahale MH, Bhusari RB. Propagational studies in fig as affected by plant growth regulator. *Asian Journal of Horticulture*. 2012; 7(1):118-120.

22. Singh KK. Effect of IBA concentrations on the rooting of pomegranate (*Punica granatum L.*) cv. Ganesh hardwood cuttings under mist house condition. Plant Archives. 2014;14(2):1111-1114.
23. Dahale M, Ningot EP, Deepa NM. Effect of Plant Growth Regulators on Rooting and Survival of Hard Wood Cuttings in Fig. International Journal of Current Microbiology and Applied Sciences. 2018; 6: 2386-2391.
24. Mehta SK, Singh KK, Harsana AS. Effect of IBA concentration and time of planting on rooting in pomegranate (*Punica granatum L.*) cuttings. J. Medi. Plants Studies. 2018;6(1): 250-253.
25. Kaur A. and Kaur A. Effect of IBA concentrations on success of cuttings of fig cv. Brown Turkey. International Journal of Recent Scientific Research. 2017; 8(11): 21576-21579.
26. Patel KD, Butani AM, Thummar BV, Purohit HP, Trambadiya RD. Response of different media and IBA on rooting and survival percentage of hardwood cutting in pomegranate (*Punica granatum L.*) cv. Bhagwa. Journal of Pharmacognosy and Phytochemistry. 2020;9(5):322-329.
27. Galston AW, Davies PJ. Hormonal regulation in higher plants. Science. 1969; 163:1288-1297
28. Kumari R. Studies on the effect of IBA and rooting media on rhizogenesis of cuttings of pomegranate (*Punica granatum L.*) cv. Bhagwa under shade net conditions. M. Sc. (Horti.) Thesis, Dr Y S R Horticultural University. AP; 2014.
29. Hakim A, Jaganath S, Honnabyraiah MK, Kumar M, Singh AK. and Dayamani KJ. Effect of biofertilizers and auxin on total chlorophyll content of leaf and leaf area in pomegranate (*Punica granatum L.*) cuttings. International Journal of Pure and Applied Bioscience. 2018; 6(1): 987-991.
30. Ghani MK, Habibi HK. Effect of growing media on rhizogenesis and growth of rooted stem cuttings of pomegranate (*Punica granatum*) cv. Phule Bhagwa Super under open field condition. International Journal of Current Microbiology and Applied Science. 2019;8(7):915-923.
31. Cheerna GS. Bhat SS, Naik FC. Commercial fruits of India with special reference to Western India, McMillan 3rd Co. Ltd; 1954.
32. Mewar D, Nautiyal MC, Singh KK. Response of various IBA concentrations to wild fig (*Ficus almate* Forsk.) cuttings under controlled condition. International Journal of Advanced Research and Management, Special Issue (I). 2018;2455-6378.
33. Patel HR, Patel MJ. Role of auxins on rooting of different types of cuttings in fig. International Journal of Current Microbiology and Applied Sciences. 2018;7(3):1317-1322.
34. Patel HR, Patel MJ and Singh S. Effect of different levels of IBA and NAA on rooting of hardwood and semi hardwood cutting in fig. International Journal of Agricultural Science and Research. 2017;7(4):519-524.
35. Shahab M, Ayub G, Rahman A, Rashid A, Jamal A, Ali J. Assessment of IBA (Indole Butyric Acid) levels and planting time for vii rooting and growth of alstonia cuttings. Journal of Natural Sciences Research. 2013;3(14):59-67.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/115234>