



Evaluation of the Phytochemical and Elemental Composition of *Andrographis paniculata* Leaves

Ime R. Ekanem ^{a*}, Nsikak B. Essien ^a, Ima I. Ekanem ^a,
R.S.Zayyan ^b and Iheanyichukwu E. Chukwuemeka ^c

^a Department of Chemical Sciences, Akwa Ibom State Polytechnic, Ikot Osurua, Nigeria.

^b Department of Basic and Applied Sciences, Hassan Usman Katsina Polytechnic, Katsina State, Nigeria.

^c Department of Science Laboratory Technology, Ibom Metropolitan Polytechnic, Uyo, Akwa Ibom State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajocs/2024/v14i4316>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119359>

Short Research Article

Received: 06/05/2024
Accepted: 08/07/2024
Published: 05/08/2024

ABSTRACT

This research work was designed to determine the phytochemical and elemental composition of *Andrographis paniculata* leaves grown in Ikot Ekpene Local Government Area of Akwa Ibom State, Nigeria. The analysis was carried out using standard procedures. The results of the phytochemical constituents were as follows: alkaloids ($2.48 \pm 0.36\%$), saponins ($9.56 \pm 0.77\%$), Flavonoids ($4.57 \pm 0.07\%$), Tannins ($12.34 \pm 0.36\%$) and cardiac glycosides ($0.39 \pm 0.01\%$). The results of the elemental composition were as follows: Magnesium ($3.02 \pm 0.05\text{mg/kg}$), calcium ($5.57 \pm 0.07\text{mg/kg}$), potassium ($20.70 \pm 0.11\text{mg/kg}$), sodium ($1.58 \pm 0.61\text{mg/kg}$), Zinc ($2.17 \pm 0.08\text{mg/kg}$),

*Corresponding author: E-mail: imeekanem55@gmail.com;

Cite as: Ekanem, Ime R., Nsikak B. Essien, Ima I. Ekanem, R.S.Zayyan, and Iheanyichukwu E. Chukwuemeka. 2024. "Evaluation of the Phytochemical and Elemental Composition of *Andrographis Paniculata* Leaves". Asian Journal of Chemical Sciences 14 (4):42-46. <https://doi.org/10.9734/ajocs/2024/v14i4316>.

Copper (10.45 ± 0.14 mg/kg) and Iron (12.50 ± 0.21 mg/kg). The phytochemicals present in this plant may be responsible for its medicinal properties. The elemental composition shows that the plant may be rich in minerals.

Keywords: *Andrographis paniculata*; phytochemicals; mineral elemental.

1. INTRODUCTION

Andrographis paniculata is an annual herbaceous plant in the family Acanthacea, native to India and Srilanka. It is widely cultivated in South Eastern Asia, where it has been believed to be a treatment for bacterial infections and some diseases [1].

Andrographis paniculata is a plant that grows in Shady places. The slender stem is dark green in colour. The leaves are lance-shaped measuring up to 8cm long by 2.5cm [2]. The fruit is a capsule around 2cm long and a few millimeters wide. The flowering time is September to December, *Andrographis paniculata* is a plant which grows in plane lands, hills, farms, moist habitats, seashores and roadsides. It can also be cultivated in gardens. This plant is commonly known as the king of bitters because of its intensely bitter taste.

Andrographis paniculata is being used in traditional medicine, as a remedy for the cold, fever and detoxification of the body [3]. *Andrographis paniculata* leaves and stems work by stimulating the immune system. It also prevents viruses from binding cells in the body [4]. People commonly use *Andrographis paniculata* for the infection of the throat and a type of bowel disease [3]. *Andrographis paniculata* is a medicinal plant traditionally used as an anti-inflammation and anti-bacterial herb. It exhibits diverse pharmacological activities including anti-cancer, anti-obesity, anti-diabetic and other activities [1]. These medicinal activities of this plant are due to the presence of certain phytochemicals in the plant. Phytochemicals, a bio-active chemical components of plants, are considered as secondary metabolites, synthesized naturally in all parts of the plant body and play a major role in the medicinal activity of the plant [5,6].

2. MATERIALS AND METHODS

2.1 Sample Collection and Treatment

The leaves of *Andrographis paniculata* were collected from a farm in Ikot Osurua village in

Ikot Ekpene Local Government Area of Akwa Ibom State, Nigeria. The leaves were air-dried for 7 days and later ground into powder form using an electric grinder. The powder samples were stored in air-tight containers for the analysis.

2.2 Chemical Analysis

Magnesium, calcium, potassium, sodium and trace elements (Zinc, Copper and Iron) were determined according to a method by Shahidi et al. [7].

Alkaloid was determined according to the method of Harbone [8]. Flavonoid was determined using the method of Boham and Kocipal [9]. Saponin was determined according to the method of Obadomi and Ochuko, 2001. Tannin was determined using the method of Van Burden and Roberson [10]. Cardiac glycosides were determined using the alkaline titration method of AOAC.

2.3 Statistical Analysis

All measurements were replicated three times and standard deviations were determined.

3. RESULTS AND DISCUSSION

The result of the phytochemical and elemental composition of *Andrographis paniculata* is shown in Tables 1 and 2.

Table 1. Result of the phytochemical composition of *Andrographis paniculata* leaves

Phytochemicals	Percentage composition (%)
Alkaloid	2.48 ± 0.36
Saponin	9.59 ± 0.72
Flavonoid	4.51 ± 0.02
Tannins	12.34 ± 0.04
Cardiac glycosides	0.39 ± 0.01

3.1 Phytochemical Screening

The bioactive substances present in the *Andrographis paniculata* leaves are presented in

Table 1, these include; saponin, flavonoid, cardiac glycosides and tannins which were present in high concentration. Alkaloid was moderately present. The presence of alkaloids commonly used in drug preparation may account for the medicinal use of the plant [5,11]. For example, alkaloids have many pharmacological activities including antihypertensive effects (many indole alkaloids), antimalarial activity (quinine), and anticancer actions (dimeric indoles, vincristine, vinblastine) [12,6]. Some alkaloids have stimulant properties such as caffeine and nicotine, morphine is used as an analgesic. Phytochemicals have roles in the protection of human health, when their dietary intake is significant [6].

Table 2. Result of the elemental composition of *Andrographis paniculata* leaves

Minerals	Concentration (mg/kg)
Magnesium (Mg)	3.02 ± 0.05
Calcium (Ca)	5.57 ± 0.07
Potassium (K)	20.70 ± 0.11
Sodium (Na)	1.58 ± 0.61
Zinc (Zn)	2.17 ± 0.08
Copper (Cu)	10.45 ± 0.14
Iron (Fe)	12.50 ± 0.21

Data are mean ± standard deviation of triplicate determinations on dry weight basis

As presented in Table 2, the concentration of potassium is 20.70 ± 0.11 mg/kg. Potassium plays a major role in maintaining the physiological balance within the human body [13]. It is an electrolyte that works in tandem with sodium to regulate fluid balance, nerve impulses, and muscle contractions. Potassium is particularly important for the proper functioning of the heart. It helps to maintain the electrical conductivity of Cardiac cells and also contributes to regular heartbeat [14]. The concentration of sodium is 1.58 ± 0.61 mg/kg in the leaves of the plant. Sodium is a vital element in the human body, playing a crucial role in maintaining fluid balance, nerve function and muscle contraction (Gupba et al., 2003). It is an essential electrolyte that works in conjunction with potassium to regulate the distribution of water in and out of the body cells. Sodium also plays a pivotal role in generating electrical impulses within nerve cells, by facilitating the transmission of signals through the nervous system [14].

The concentration of Calcium in the leaves of *Andrographis paniculata* is (5.57 ± 0.07mg/kg). Calcium is an essential mineral that plays a

crucial role in numerous physiological functions within the human body [15]. Primarily, calcium is known for its role in bone health providing strength and rigidity.

Approximately 99 % of the body's calcium is stored in bones and teeth, contributing to their development, maintenance, and overall structural integrity. Beyond its skeletal function, Calcium is also vital for various cellular processes, including muscle contraction, nerve transmission and blood Clotting [15].

The concentration of magnesium is 30.22 ± 0.05mg/kg in the leaves of the plant. Magnesium is a vital mineral that plays a multifaceted role in maintaining various physiological functions, within the human body [16]. As an essential cofactor for more than 300 enzymatic reactions, magnesium is involved in energy metabolism, DNA and RNA Synthesis and the regulation of ion channels. It contributes significantly to muscle and nerve function, supporting muscle contractions and the transmission of nerve impulses [14].

The concentration of Zinc is 21.66 ± 0/08mg/kg in the leaves of the plant. Zinc is an essential trace element that plays a vital role in various physiological processes within the human body. As a cofactor for numerous enzymes, zinc is involved in the regulation of gene expression, immune function, wound healing and DNA synthesis [14].

It is particularly crucial for the proper functioning of the immune system, as zinc supports the development and activity of immune cells, helping the body defend against infections and illnesses [17]. Additionally, zinc plays a role in cell division, growth and the maintenance of healthy skin.

The concentration of copper is 10.45 ± 0.14mg/kg in the leaves of the plants. Copper is an essential trace mineral that plays a crucial role in various physiological functions within the human body. As a cofactor for numerous enzymes, copper participates in fundamental processes such as energy metabolism, the formation of connective tissues and the synthesis of neurotransmitters [17]. One of the notable roles of copper is its involvement in the formation of hemoglobin, the protein responsible for transporting oxygen in the blood. Additionally, copper contributes to the maintenance of healthy bones and connective tissues by promoting the cross-linking of collagens [14].

The concentration of Iron is (12.50 ± 0.71 mg/kg) in the leaves of the plant. Iron is a vital trace element that plays a crucial role in several essential functions within the human body [18]. Its primary function involves oxygen transport, as iron is a key component of hemoglobin, the protein in red blood cells responsible for binding and carrying oxygen from the lungs to tissues throughout the body [14]. Additionally, iron is a critical component of hemoglobin, a protein that facilitates oxygen storage and release in muscle cells, supporting muscle function. The toxicity of any metal is a function of its concentration in the sample, exposure and the chemical nature of the metal ions [19]. A man requires between 2 and 3 mg/l of iron per day and 60% to 70% of ingested amounts are metabolized [20]. However, the high concentration of iron does not appear to present a health hazard but it may affect the taste of beverages or medicine made from the plant [21-25].

4. CONCLUSION

The elemental content of *Andrographis paniculata* reveals a significant concentration of key elements essential for human health, which include potassium, calcium, magnesium and sodium, alongside trace amounts of zinc, copper and iron. These findings underscore the potential nutritional value of *Andrographis paniculata* as a rich dietary supplement, as well as its potential in the treatment of various inflammatory conditions, including diabetes and inflammatory bowel disease. The study also revealed that *Andrographis paniculata* possesses medicinal properties and therefore should be used in herbal medicine for the treatment of various illnesses.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

Special gratitude goes to Tertiary Education Trust Fund (TETFUND), Abuja for a grant given to Ime R. Ekanem in support of this work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ekanem IR. Characterization of the ethanolic extract of *Andrographis paniculata* using fourier transform infrared spectrophotometer. International Journal of Sciences and Energy Research. 2018;4(2):2-5.
2. Akbar S. *Andrographis paniculata*: A review of phytochemistry, ethnopharmacology, and pharmacology. Journal of Ethnopharmacology. 2018;218:44-60.
3. Pramono AA. Anti-inflammatory effect of andrographolide in the brain of wister rats exposed to force swimming stress. Inflammation. 2016;39(6):109-112.
4. Sharma A. Andrographolide: A novel antiviral compound against hepatitis B virus infection. Frontiers in Microbiology. 2017;8:188-190.
5. Akpabio UD, Wilson LA, Akpakpan AE, Obot IB. Phytochemical screening and proximate composition of *Cassia hirsute* seeds. Elixir Journal of Applied Chemistry. 2012;8704-8707.
6. Akpakpan AE, Ukpong EJ, Odiongenyi AO, Willie IE. Phytochemical screening of the ethanol and aqueous extracts of *Dicliptera verticillata* Leaves. Archives of Current Research International. 2017;11(2):1-5.
7. Shahidi FU, Chavan AK, McKenzie OB. Chemical composition of beach Pea, *Lathyrus maritimum* plant parts. Food Chemistry. 2002;64:39-44.
8. Harborne JB. Phytochemical methods, Chapman and Hall, London. 1973;113-114.
9. Boham BA, Kocipai AC. Flavonoids and condensed tannins from leaves of *Hawaina vaccinium* and *Vaticulum calycinium*, Pacific Science. 1994;48:458-463.
10. Van-Burden TP, Robinson WC. Formation of complexes between protein and tannic acid. Journal of Agriculture and Food Chemistry. 1981;1:77-82.
11. Uwanta EJ, Udo IE, Willie IN, Akpakpan AE. Phytochemical screening, proximate and vitamin composition of Jack fruit (*Artocarpus heterophyllus*) pulp and seed. World Journal of Advanced Research and Reviews. 2024;22(01):1500-1506.
12. Wink M, Schmeller T, Latz-Briining B. Modes of action of allelochemical alkaloids: Interaction with neuroreceptors,

- DNA and other molecular targets. *Journal of Chemical Ecology*. 1998;24:1888-1937.
13. Gupta KK, Bhattacharjee S, Kar S, Chakrabarty S, Thakur P, Bhattacharyya G, Srivastava SC. Mineral composition of eight common spices. *Communication Oil Science and Plant Analogy*. 2003;34(6):6810-691.
 14. Abai EJ, Ekanem IR. Evaluation of phytochemical and elemental composition of *Hura crepitans* seeds. *Asian Journal of advances in Research*. 2020;4(4):12-16.
 15. Hefsi C, Debez PK. Calcium: A central regulator of plant growth and development, *The Plant Cell*2015;17(8):2142-2155.
 16. Carmak I, Kirkby EA. Role of magnesium in carbon partitioning and alleviating photo-oxidative damage. *Physiologia plantarum*. 2008;133(4):692-704
 17. Ansari TM, Ikram NM, Fayyaz Q, Ghafoor I, Khalid N. Essential trace metal (Zinc, manganese, copper and iron) levels in plants of medicinal importance. *Journal of Biological Chemistry*. 2004;4(2):95-100.
 18. Briat JF, Dubos C, Gaymard F. Iron nutrition, biomass production, and plant product quality. *Trends in plant Science*. 2007;12(1):40-47.
 19. Etuk, Bassey A, Udiog, Daniel S, Akpakpan, Aniekan E. Human health risk assessment of trace metals in water from cross river estuary, Niger Delta, Nigeria. *Asian Journal of Chemical Sciences*. 2020;7(3):1-11.
 20. World Health Organization (WHO), Guidelines for drinkingwater quality; 2011.
 21. Nsi EW, Uwanta EJ, Akpakpan AE, Ekwere IO. Analytical assessment of borehole water in some local government Areas of Akwa Ibom State, South-South Nigeria. *European Scientific Journal*. 2020;6(13):122-136.
 22. AOAC. Official methods of analysis. 15 Edition. Association of Official analytical chemists, Benjamin Franklin Station, Washington D.C. 1990;1160-1167.
 23. Chang RC, Lee HC, Ou AM. Investigation of the Physiochemical properties of concentration fruit vineger. *J. Food Drug Anal*. 2015;3:356-384.
 24. Kumar VS. A comprehensive Review on the medical plants from the Genus *Andrographis*. *Frontier in Microbiology*. 2016;7:743-745.
 25. Akpabio UD. and Akpakpan A E. Evaluation of Nutritive and Anti-nutritive compositions of the seeds of *Monodora myristica* (African Nutmeg). *World Journal of Applied Science and Technology*. 2012; 4(1): 49-55.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119359>