



# Assessing the Seroprevalence and Risk Factors of Bovine Brucellosis in Meghalaya, North-east India

**Hosterson Kylla <sup>a\*</sup>, Laureata Dkhar <sup>a</sup>, Lahun R. Dkhar <sup>a</sup>, Ernestine Basaiawmoit <sup>a</sup>, Iadarilin Warjri <sup>a</sup> and Evando Sumer <sup>a</sup>**

<sup>a</sup> *Disease Investigation and Surveillance, Department of A. H. & Veterinary, Government of Meghalaya. Shillong-793002, India.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/mrji/2024/v34i101491>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/121689>

**Original Research Article**

**Received: 22/07/2024**

**Accepted: 24/09/2024**

**Published: 03/10/2024**

## **ABSTRACT**

**Aim:** A cross-sectional prevalence study was conducted from 2015 to 2018 in Meghalaya state (India) to determine prevalence of bovine brucellosis by serological tests such as Rose Bengal Plate Test (RBPT), Indirect ELISA (iELISA) and Milk Ring Test (MRT).

**Materials and Methods:** Serum samples were collected from 3193 cattle aged 1 year to 12 years and screened for Brucella antibodies using Rose Bengal Plate Test (RBPT) and Indirect ELISA (iELISA). The samples were collected from both male and female cattle reared in organized farms and smaller private holdings. Individual milk samples (n=896) from representative dairy cattle were screened for brucellosis using the Milk Ring Test (MRT).

\*Corresponding author: E-mail: [hosterkylla123@yahoo.com](mailto:hosterkylla123@yahoo.com);

**Cite as:** Kylla, Hosterson, Laureata Dkhar, Lahun R. Dkhar, Ernestine Basaiawmoit, Iadarilin Warjri, and Evando Sumer. 2024. "Assessing the Seroprevalence and Risk Factors of Bovine Brucellosis in Meghalaya, North-East India". *Microbiology Research Journal International* 34 (10):59-67. <https://doi.org/10.9734/mrji/2024/v34i101491>.

**Results:** The iELISA revealed a serum positivity rate of 9.89% while the RBPT detected a positive rate of 9.33%. The prevalence rate was higher in females (12.32%) compared to males (2.21%). Cattle aged 1-4 showed higher susceptibility (11.66%) than other age groups. Higher prevalence were detected from organized farms maintaining high number of animals (13.09%) compared to smaller private holdings (9.02%). Cattle with a history of abortion and related clinical symptoms also had a higher and significant association with sero-positivity. The MRT indicated a high positivity of 17.29% among sampled dairy cattle.

**Conclusion:** Brucellosis is prevalent in the hilly state of Meghalaya, India. This study insight into the prevalence of bovine brucellosis in the state highlighting potential risk factors for transmission to human. The high prevalence of brucellosis highlights the urgent need for enhanced surveillance and control measures in Meghalaya such as implementing targeted vaccination programs of all eligible female calves and intense information education campaign (IEC) to all the farmers and general public which could significantly reduce the disease's impact.

**Keywords:** Brucellosis; seroprevalence; cattle; RBPT; iELISA.

## 1. INTRODUCTION

“Meghalaya state located in the North East Region of India relies on agriculture and livestock farming for its livelihood and economy. Amongst the various diseases affecting bovine species, brucellosis is a significant health problem in many countries. While the disease is prevalent globally, but there are countries such as Canada, Japan, Australia and New Zealand, cases of *Brucella abortus* and *Brucella melitensis* have never been reported” [1]. “The disease is endemic in many states and regions of India. In cattle, the disease is mainly caused by *Brucella abortus* and is associated with widespread contagious reproductive disease of dairy animals. In India it is highly prevalent among the bovine population thus causing huge economic losses to the farmers and other stake holders. Bovine brucellosis has also been attributed to *B. melitensis* and infrequently to *B. suis*” [2,3]. *Brucella* organism is coccobacillary shaped bacteria, Gram-negative, facultative, intracellular bacteria, comprising of many different species based on biochemical reactions and host affinity.

“Currently, there are ten spp. described in the genus *Brucella*. Each species may infect various animal hosts, but has preferences for particular host animals, such as *B. abortus* for cattle, *B. suis* for pigs, *B. melitensis* for sheep and goats, *B. ovis* for rams, *B. canis* for dogs, *B. microti* infect rodents - (*Microtus arvalis*), *B. neotomae* for rodents - (*Neotoma lepida*), *B. Pinnipedialis* for pinnipeds, *B. ceti* for cetaceans, *Brucella inopinata* which is first isolated from humans, does not yet have a known preferential host. The underlying mechanisms of intracellular survival, including the host determinants remain relatively unknown” [4,5]. “In young animals and non-

pregnant females, symptoms of brucellosis are usually not noticeable. However, in pregnant adult females, symptoms of *B. abortus* and *B. melitensis* includes placentitis often resulting in abortion between the fifth and ninth month of pregnancy. Adult male cattle may develop orchitis and/or epididymitis. In both males and females, may experiences infertility due to brucellosis. In some tropical countries, hygromas particularly of leg joints are a common manifestation of disease” [1,6]. “Additionally, the disease can cause chronic arthritis, reducing the productive lifespan of infected cattle” [7].

“To control the disease, many countries adopt a test and slaughter policy for infected animals, with proper disposal following confirmatory diagnostic tests” [8]. “Regulation of the disease often relies on vaccination and culling of infected animals to minimize the risk of spreading the disease to consumers and people involved in regular animal farming activities” [9]. In India, the main control programme for bovine brucellosis is based on timely vaccination of young female cattle (4-8 months old) and isolation of positive reactors.

The Milk ring test (MRT) first described by Fleischhauer in 1937 [7], is an important serological test to detect brucellosis particularly at the herd level. The test can detect IgM and IgA antibodies bound to fat globules. It is widely accepted due to its cost effectiveness, ease of performance and ability to cover a large population in a short time [10]. However, MRT can sometimes produce false positives commonly in colostrum or milk at the end of the lactation period and milk from cows suffering from hormonal disorder or mastitis, which therefore needs confirmatory testing through

more specific assays like RBPT and iELISA [11, 12].

Understanding the prevalence and risk factors of brucellosis in a region is critical to develop targeted intervention strategies for controlling the disease [13]. Information on the status of bovine brucellosis is limited in North East region of India particularly in a hilly and tribal state of Meghalaya. Therefore, this study aimed to determine seroprevalence of bovine brucellosis using RBPT and iELISA tests, as well as to detect infection through the MRT in individual milked cattle.

## 2. MATERIALS AND METHODS

### 2.1 Collection of Samples

A total of 3193 serum samples were collected from cattle of various organized farms (n=28) and private holdings (n=96) across different districts of Meghalaya, India between January 2015 and December 2018. The serum samples were collected by the Disease Investigation team, A.H & Veterinary Department, Meghalaya, Shillong. The samples were based on representative sampling with reference to age, sex, production and also through a purposive sample approach based on farmer's requests. The samples were collected from Holstein Friesian, Jersey and various cross breeds. Approximately 7 ml of blood was collected from the jugular vein of each animal using individual sterile syringes and commercially available serum separator vacutainers (Becton Dickson, USA). Samples were properly labelled, maintaining a cold chain at 4°C during transportation till serum separation process in the laboratory and all the clear serum samples were stored at -20°C until tested. Herd and animal level data were recorded including age group, sex, farm size, and history of abortion or repeat breeding. Of the 3193 animals sampled, 2425 were females (75.94%) and 768 (24.05%) were males. A total of 1937 sera were from cattle aged 1 to 4 years, 789 sera were from cattle aged 5 to 8 years and 467 sera were from cattle older than 8 years of age. Additionally, 1420 (44.47%) of the samples were from organized farms (including government run farms) with more than 20 lactating dairy cattle, while 1773 (55.52%) were from private holdings with fewer than 20 cattle.

### 2.2 Serological Tests

All the serum samples were used to evaluate disease prevalence using serological tests,

namely the Rose Bengal Plate test (RBPT) and the indirect Enzyme Linked Immunosorbent Assay (iELISA). According to [14], sensitivity and specificity were 96.8% and 96.3% respectively for iELISA, whereas estimates of RBPT were 89.6% and 84.5% respectively. The 3193 serum samples were analysed by RBPT according to standard protocol [15]. A known positive and negative serum was included in each day's tests. Briefly, 30µl of serum sample was mixed thoroughly using a clean tip for each sample with 30µl of Rose Bengal antigen on the clean slide. The mixture was mixed and agitated softly for 4 min. The test was read by examining for agglutination in a good light, best observed as the mixture flows on the slide. Results were recorded as positive (any degree of agglutination) or negative (no agglutination). The *B. abortus* S99 colored antigen used in the study was procured from the Institute of Animal Health and Veterinary Biologicals (IAH&VB), Kolkata, India.

The iELISA was performed on the serum samples using a commercial kit (IDEXX Brucellosis Serum X2 Ab Test, USA). The test followed the manufacturer's recommendation. Colour development in the form of the optical density (OD) was read at 450 nm using an ELISA microplate reader (Infinite F50, Tecan, Austria). Results were expressed as the ratio of the sample (OD) minus the mean kit negative control OD to the mean kit positive control OD minus the mean kit negative control OD (S/P ratio). According to manufacturer, a positive result was defined as an S/P ratio of ≥80%, and negative results was for an S/P ratio of <80%.

A total of 82 sampling units (cattle herds) from different farms in the state were sampled, totaling 896 individual lactating cows of various age groups. Approximately 10ml milk pooled from 4 quarters was collected and properly labelled. Milk samples were kept refrigerated at 4°C overnight prior to examination by Milk Ring Test (MRT) [15]. The milk samples were mixed well to ensure an even distribution of the milk cream.

"The MRT is based on agglutination of antibodies secreted into the milk allowing screening of large number of cattle by using milk samples from tanks or pools from several cows. This test is useful for monitoring cattle herds or areas free of brucellosis and is classified as surveillance or monitoring test" [16]. The MRT is also the most practical method for locating infected dairy animals. In this study, 896 milk samples were

analysed by MRT according to manufacturer's instructions. The *B. abortus* Bang Ring Antigen was procured from Institute of Animal Health and Veterinary Biologicals (IAH&VB), Kolkata, West Bengal, India. The test was performed by adding 30µl of *B. abortus* Bang Ring Antigen (hematoxylin-stained antigen) to 5ml milk in a test tube. The milk-antigen mixtures were incubated at 37°C for 1 hour, along with positive and negative control samples. A strongly positive reaction was indicated by formation of a dark blue ring above a white milk column. The test was considered negative if the color of the underlying milk exceeded that of the cream layer and the cream layer appeared normal.

### 3. RESULTS AND DISCUSSION

“Out of 3193 serum samples, 298 (9.33%) were detected positive by RBPT test, whereas 316 (9.89%) samples were positive by iELISA. The year-wise prevalence rate is depicted in Fig. 1. Indirect ELISA is a confirmatory diagnostic test for bovine brucellosis; hence, the overall seroprevalence of bovine brucellosis was recorded at 9.89% based on serum samples that tested positive by iELISA. In comparison, several researchers have reported lower prevalence rate. For instance, in Southern Ethiopia, a prevalence of 2.46% was reported” [17]. “Another study from Peninsular Malaysia reported Brucella antibodies in 2.5% of sampled cattle and in Central Ethiopia

0.7% prevalence was reported during 2013-14” [6,18]. “In India, antibodies to Brucella were reported to have a positivity rate of 5.22% by RBPT and 6.03% by iELISA. A study in Meghalaya reported a seroprevalence in cattle at 5.91% by RBPT and 11.29% by ELISA, particularly in neighbouring border areas with Assam State” [19]. “The seroprevalence of 9.89% in the present study is lower compared to findings of other countries such as Egypt (11%) and Nigeria (19.7%)” [20,21]. The disparity in disease prevalence report by different studies indicate the variability of brucellosis infection which may be attributed to various extrinsic factors such as the type of surveillance activities, farm management and production systems including cattle-rearing practices, and the level of stringency regarding disease-control measures adapted in different countries; infection status of the sampled animals and number of samples collected during the study period.

“By comparing the two commonly used serological tests, i.e., RBPT and iELISA, it shows that RBPT can also detect accountable number of sero-positive samples similar to iELISA. Even though its limitation and lack of specificity in detecting Brucella antibodies, RBPT is still widely used for preliminary screening for brucellosis in many countries” [22]. The high sensitivity of iELISA is detected in recovered or vaccinated




Fig. 1. Year-wise RBPT and iELISA Results for Sampled Cattle Population

**Table 1. Risk Factors Associated with Bovine Brucellosis in Meghalaya State, India**

Risk factors		No. of animals	RBPT positive	ELISA positive
<b>Sex</b>	Female	2425	286 (11.79%)	299 (12.32%)
	Male	768	12 (1.56%)	17 (2.21%)
<b>Size of herd</b>	Organized (>20)	1420	172 (12.11%)	186 (13.09%)
	Small holdings (<20)	1773	150 (8.46%)	160 (9.02%)
<b>Age (Yrs)</b>	1-4 years	1937	218 (11.25%)	226 (11.66%)
	5-8 year	789	55 (6.97%)	60 (7.60%)
	>8 years	467	25 (5.35%)	30 (6.42%)
<b>History of abortion and repeat breeding</b>	Yes	16	10 (62.5%)	10 (62.50%)
	No	3177	288 (9.06%)	306 (9.63%)


animals due to the persistence of IgG antibody for longer period. Hence, seroprevalence detected by iELISA may reflect either past or present exposure to *Brucella* organisms. However, since no brucella vaccination was administered to the sampled animals on different farms in Meghalaya during the study period, the presence of vaccinal antibody can be ruled out. Molecular detection by polymerase chain reaction (PCR) using serum and blood DNA could be to further validate the results.

“In this study, important risk factors associated to bovine brucellosis were also analysed and results were depicted in Table 1. The prevalence by sex showed that brucellosis is higher in females with a rate of 12.32% (299/2425) compared to males which had a prevalence of 2.21%; 17/768. This low prevalence of brucellosis in male cattle aligns with previous finding by other investigators” [18,23]. The spread of the disease in the herds is mainly due to lack of periodical screenings in large female bovine population and undiagnosed infected females. Researchers [18,24] reported that “1% and 6.63% female cattle were found to be seropositive”. Additionally, it was extensively reported by [25] that “the sex of the susceptible animal species is one of the many risk factors affecting the susceptibility of cattle to *Brucella abortus* infection”.


“A higher prevalence was detected from organized farms maintaining a larger number of lactating animals (13.09%; 186/1420) compared to smaller private holdings (9.02%; 160/1773). In the case of bovine brucellosis, the greater chances of spreading of infection is found especially in organized herds than in marginal herds” [26]. “In small farms various factors like sufficient unit floor space for each animal; stall feeding that minimizes contact with other infected animals and more personnel attention to the

animals from the farmer himself contribute to the reduced spread of infection. As per age category, the prevalence of brucellosis was higher in those cattle whose age ranged from 1 to 4 years compared to other age groups (Fig. 2)” [27]. According to [25] “susceptibility of animals to disease is in fact more commonly associated with sexual maturity. The higher prevalence in the age group of less than 4 years old may be attributed to new exposure to brucellosis infected animals in the farms. The present study also revealed that brucellosis is mostly prevalent (62.50%) in areas and farms where abortion, repeat breeding problems and other reproductive complications are reported. Hence, the prevalence of brucellosis was closely associated with their history of abortion in the examined animals, which infected animals may act as carriers of the infection. Representative individual dairy cattle milk sample collected between 2015-2018 and tested by MRT revealed a high positivity rate of 17.29% (155/896) regardless of the animal’ age groups, as depicted in Fig. 3” [35]. This finding correlates with the research of [28] who detected 18.35% positivity for *Brucella* and [29] in Iran, who detected a 14% positive reaction for *B. abortus* using MRT across various age groups into consideration. Another researcher [30] observed that 12.5% of animals were positive for brucellosis by MRT. While the MRT is preferred for its simplicity in detecting *B. abortus* infection, however, it has also been reported to produce a number of false positive results [31].

“Despite of various preventive and control measures being followed in India, there remains a high potential for the transmission and spread of *Brucella abortus* due to its widespread prevalence” [32]. “Timely confirmatory laboratory testing of animals along with emergency attentive animal health care, should be utilized to diagnose any related abortions cases, premature



**Fig. 2. RBPT and iELISA Results by Age Group in Sampled Cattle Population**



**Fig. 3. Year-wise Results of the Milk Ring Test with Total Prevalence**

births and other clinical signs. This should be followed by total disinfection of the farms using recommended disinfectants. Careful selection of animals before purchase particularly from brucella-free farm, along with pre-purchase tests and quarantine measures, should be

judiciously followed to keep the animals free of brucellosis” [33]. Current studies highlighted the importance of surveillance and biosecurity measures in controlling brucellosis in cattle populations. For example, [34] emphasized the need for improved vaccination strategies and

diagnostic capabilities in endemic regions to reduce the spread of the disease. Furthermore, the utilization of molecular diagnostics, such as conventional or real-time PCR, has more accuracy in identifying *Brucella* spp. in various samples, providing a more precise method for disease management [35]. These findings suggest that integrating traditional serological methods with molecular techniques, along with biosecurity protocols, could significantly decrease the burden of brucellosis in cattle populations.

#### 4. CONCLUSION

The study reveals that seroprevalence of bovine brucellosis (9.89%) is comparatively higher in the hilly state of Meghalaya with various potential risk factors that need proper attention to reduce the disease and prevent production loss. Cases of bovine brucellosis have been increased in certain areas of Meghalaya State, possibly due to increased cattle trade and demand, undetected movement of cattle from other states, and possibly from neighbouring border countries. Milk Ring test detected a high positivity of 17.29%, however, it is suggested that other confirmatory tests are to be used in conjunction. The presence of sero-positive reactors for brucellosis indicates foci of infection that leads to the spread of the disease. Hence, greater attention by concerned stakeholders in the State is urgently required, particularly focusing on mass vaccination programme of eligible animals, information educational campaign of general public and farmers on preventive measures to safeguard and prevent transmission risk of the infection to human population.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### ACKNOWLEDGEMENTS

The authors are thankful to the DAHD, Government of India for funding and the Director, A.H & Veterinary Department, Government of Meghalaya for providing facilities to conduct the study.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. OIE Terrestrial Manual. Manual of diagnostic tests and vaccines for terrestrial animals: Bovine brucellosis; 2016.
2. Patel MD, Patel PR, Prajapati MG, Kanani AN, Tyagi KK, Fulsoundar AB. Prevalence and risk factor's analysis of bovine brucellosis in peri-urban areas under intensive system of production in Gujarat, India. *Vet World*. 2014;7(7):509-16.
3. Sanogo M, Abatih E, Thys E, Fretin D, Berkvens D, et al. Risk factors associated with brucellosis seropositivity among cattle in the central savannah forest area of Ivory Coast. *Prev Vet Med*. 2012;107:51-56.
4. de Jong MF, Tsolis RM. Brucellosis and type IV secretion. *Future Microbiol*. 2012;7:47-58.
5. Hadush A, Pal M. Brucellosis: An infectious re-emerging bacterial zoonosis of global importance. *Int J Livestock Health*. 2013;3: 28-34.
6. Anka MS, Hassan L, Adzhar A, Bejo SK, Mohamad RB, Zainal MA. Bovine brucellosis trends in Malaysia between 2000 and 2008. *BMC Vet Res*. 2013;9: 230.
7. Fleischhauer G. Die Abortus-Bang-Ringprobe (ABR) zur Feststellung von bangverdächtigen Vollmilchproben. *Berl Tierarztl Wochenschr*. 1937; 53: 527-528.
8. Taleski V, Zerva L, Kantardjiev T, Cvetnic Z, Erski-Biljic M, et al. An overview of the epidemiology and epizootiology of brucellosis in selected countries of Central and Southeast Europe. *Vet Microbiol*. 2002; 90:147-156.
9. Mdegela RH, Kusiluk LMJ, Kapaga AM, Karimuuribo ED, Turuka FM, et al. Prevalence and determination of mastitis and milk born zoonosis in small holder farming sector in Kibaha and Orogoro districts in eastern Tanzania. *J of Vet Med Series*. 2004;51:123-128.
10. Cadmus SIB, Adesokan HK, Stack J. The use of the milk ring test and Rose Bengal test in brucellosis control and eradication in Nigeria. *J. of the South African. Vet. Asso*. 2008;79: 113-115.
11. OIE Terrestrial Manual. Bovine Brucellosis. 2018;2.4.3
12. Delegan Y, Shorokhova A, Zvonarev A, Abashina T, Suzina N. Draft Genome Sequence of Microbacterium sp. Strain KKR3/1, an Antimicrobial-Substance-

- Producing Strain Isolated from Banana Shrimp (*Atyopsis moluccensis*). Microbiology Resource Announcements. 2022;11(2):e01242-21.
13. Regasa F, Mekonnen D, Abunna F. Seroprevalence and associated risk factors of bovine brucellosis in Sendafa, Oromia Special Zone surrounding Addis Ababa, Ethiopia. PLOS ONE. 2021;16(2): e0246303.
  14. Getachew T, Getachew G, Sintayehu G, Getenet M, Fasil A. Bayesian estimation of sensitivity and specificity of rose bengal, complement fixation, and indirect ELISA tests for the diagnosis of bovine brucellosis in Ethiopia. Veterinary Medicine International. 2016;(1): 8032753.
  15. Alton GG, Jones LM, Angus RD, Verger JM. Techniques for the brucellosis laboratory. 1<sup>st</sup> edition. Institute Nationale de le Rech. France, Paris. 1988; 174.
  16. Office International Epizootic. Bovine brucellosis in terrestrial manual. Office International Epizootic Manual; 2009.
  17. Kassahun A, Shiv P, Yilkal A, Esayas G, Gelagaye A, Aschalew Z. Seroprevalence of brucellosis in cattle and high-risk professionals in Sidama Zone, Southern Ethiopia. Ethiopia Vet J.. 2007;11: 69-84.
  18. Bashitu L, Afera B, Tuli G, Aklilu F. Seroprevalence Study of Bovine Brucellosis and its Associated Risk Factors in Debrebirhan and Ambo Towns. J Adv Dairy Res.. 2015;3: 131.
  19. Shakuntala I, Ghatak S, Sanjukta R, Sen A, Das A, Puro AK, Dutta A, Kakoty K. Incidence of brucellosis in Livestock in North-Eastern India. Int J of Infect Dis.. 2016; 45S:1-477.
  20. Holt HR, Eltholth MM, Hegazy YM, El-Tras WF, Tayel AA, Guitian, J. *Brucella spp.* Infection in large ruminants in an endemic area of Egypt: cross-sectional study investigating seroprevalence, risk factors and livestock owner's knowledge, attitudes and practices (KAPs). BMC Public Health. 2011;11:341.
  21. Mai H, Irons P, Thompson P. A large seroprevalence survey of brucellosis in cattle herds under diverse production systems in northern Nigeria. BMC Vet Res.. 2012; 8:144.
  22. Poester FP, Nielsen K, Samartino LE, Yu WL. Diagnosis of brucellosis. The Open Vet Sci J. 2010; 4: 46-60.
  23. Kubuafor DK, Awumbila B, Akanmori BD. Seroprevalence of brucellosis in cattle and humans in the Akwapim-South district of Ghana: Public health implications. Acta Tropica. 2000;76(1):45-48.
  24. Shome R, Padmashree BS, Krithiga N, Triveni K, Sahay S, Shome BR, Singh P, Rahman H. Bovine Brucellosis in organized farms of India – An assessment of diagnostic assays and risk factors. Adv Anim Vet Sci.. 2014; 2 (10): 557-564.
  25. Radostits OM, Gay CC, Blood DC, Hinchcliff KW. Veterinary Medicine: A textbook of the diseases of cattle, sheep, pigs, goats and horses. 9<sup>th</sup> edition. New York: Saunders, W.B. 2000; 867-882.
  26. Jain U, Bist B, Sahzad P, Dwivedi K. Outbreak of brucellosis in buffaloes aborted in village Mahuan, district Mainpuri, U.P., India- A case report. Vet World. 2013; 6(1): 51-52.
  27. Kylla H, Passah P, Kharchandy M, Dkhar L, Warjri I, Dkhar L. Sero-prevalence of brucellosis in cattle and its associated risk factors in North East India (Meghalaya). International Journal of Tropical Disease & Health. 2017;27(3):1-6.
  28. Mohamand N, Gunaseelan L, Sukumar B, Porteen K. Milk Ring Test for spot identification of *Brucella abortus* infection in single cow herds. J Adv Vet Anim Res.. 2014;1(2):70-72.
  29. Rezaei M., Mohebbali SH, Abadi YK, Suri E, Zare A, Malamir SH, Rasuli ES, Maadi H. Investigation on the seroprevalence and pollution severity to *Brucella abortus* and *Brucella melitensis* bacteria in cows and sheeps living in the villager region of Toyserkan city, Hamedan, Iran. J. of Ani. and Vet. Adv.. 2010; 9: 2870-2872.
  30. Sukumar K, Tamilselvan P, Dorairajan N. Studies on infertility in cattle and buffaloes caused by *Brucella abortus*. Tamil Nadu. J. Vet. and Ani. Sci.. 2012;8:235-237.
  31. MacMillan AP. Conventional serological tests. Animal brucellosis. 1990;206: 153-97.
  32. Renukaradhya GJ, Isloor S, Rajasekhar M. Epidemiology, zoonotic aspects, vaccination and control/eradication of brucellosis in India. Vet Microbiol. 2002; 90: 183-195.
  33. Dadar M, Tiwari R, Sharun K, Dhama K. Importance of brucellosis control programs of livestock on improvement of one health. Veterinary Quarterly. 2021;41(1): 137-151.



34. Rahman A, et al. Vaccination strategies for brucellosis control in endemic regions. *Journal of Veterinary Science and Technology*. 2022; 14(2): 145-159.
35. Karthik K, et al. Application of real-time PCR for the diagnosis of brucellosis in Cattle. *Veterinary Microbiology*. 2023; 265:109357.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/121689>