



Mineralogical Composition and Bioactive Molecules in the Pulp and Seed of *Patauá* (*Oenocarpus bataua* Mart.): A Palm from the Amazon

**S. A. M. Saravia^{1*}, I. F. Montero², B. M. Linhares³, R. A. Santos⁴
and J. A. F. Marcia⁵**

¹*Faculty of Earth Sciences and Conservation, National University of Agriculture, Highway to Dulce Nombre de Culmi, Km 215, Neighborhood El Espino, Catacamas-Olancho, Honduras.*

²*Department of Organic and Inorganic Chemistry, Polytechnic School, University of Extremadura, University Avenue s/n, Cáceres, Spain.*

³*Federal Institute of Roraima, Campus Boa Vista, Av. Glaycon de Paiva, 2496 - Pricumã, Boa Vista - RR, 69303-340, Brazil.*

⁴*Department of Food Science, Louisiana State University, USA.*

⁵*Faculty of Technological Sciences, National University of Agriculture, Highway to Dulce Nombre de Culmi, Km 215, Neighborhood El Espino, Catacamas-Olancho, Honduras.*

Authors' contributions

This work was carried out in collaboration among all authors. Author SAMS contributed to the analysis, literature review and formatting of the article. Author IFM contributed to the supervision of the analysis, writing of the article and review of language and style. Author BML contributed to the collection of the material, sample preparation and analysis. Authors RAS and JAFM contributed to the review, editing of the article, the verification, validation of the methods and analysis being used in the experiments. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2019/v31i630228

Editor(s):

(1) Dr. Olanrewaju Folusho Olotuah, Professor, Department of Plant Science and Biotechnology, Adekunle Ajasin University, Akungba-Akoko, Ondo State, Nigeria.

Reviewers:

(1) Mohamed Gamal Shehata, City of Scientific Research and Technological Applications, Egypt.

(2) Jayath P. Kirthisinghe, University of Peradeniya Peradeniya, Sri Lanka.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/54389>

Original Research Article

Received 01 December 2019

Accepted 04 February 2020

Published 11 February 2020

ABSTRACT

The *Patauá* (*Oenocarpus bataua* Mart.) also known as Bataua or Patuá, is a palm native from the Amazon, consumed among the local populations as well as the wine obtained from its pulp with high energy value. It is a monocle palm tree reaching between 4-26 meters tall, distributed in the Amazon rainforest both in the wet forest of floodplains. The objective of this work was to study the proximal, mineralogical composition, as well as the total phenolic compounds and antioxidant activity of pulp and seed of *Patauá*. As for mineral composition, the high concentrations of sodium for the seed ($84.21 \text{ mg } 100 \text{ g}^{-1}$) and pulp ($71.21 \text{ mg } 100 \text{ g}^{-1}$), as well as magnesium values of $48.31 \text{ mg } 100 \text{ g}^{-1}$ for the seed and $41.23 \text{ mg } 100 \text{ g}^{-1}$ for the pulp. Among the micronutrients, the high concentration of iron in the pulp is $1.84 \text{ mg } 100 \text{ g}^{-1}$ for the pulps, and the manganese was $1.10 \text{ mg } 100 \text{ g}^{-1}$ for the seeds. The total phenolic compounds found in the seeds were relatively higher than for the pulps with values of $356.12 \pm 0.12 \text{ mg GAEq g}^{-1}$ and $321.03 \pm 0.43 \text{ mg GAEq g}^{-1}$, as well as the more significant antioxidant activity for the seeds than for the pulp. Carotenoids concentration in the seeds found of $2.52 \pm 0.04 \text{ mg mL}^{-1}$ and vitamin C concentrations were also quantified in trace concentrations, presenting the fruits of *Patauá* high biotechnological interest in the food and cosmetic industry.

Keywords: *Arecaceae*; *bioresidues valorization*; *carotenoids*; *antioxidant activity*; *minerals*.

1. INTRODUCTION

Arecaceae or Palmae families, constituted by about 2400 species distributed by various tropic and subtropical areas of the planet [1] as only the Neotropic areas in Africa and Asia [2] as well as the Neotropic regions of the Amazon Region and Central America [3] where more than 150 species of palm trees are found [4].

Among the palms of this family, it is worth mentioning the *Patauá* (*Oenocarpus bataua* Mart.), which has a concrescence with presence in the Amazon region, with famous traditional people, presenting great economic potential, serving as a source of food, as well as preparing medicines [5,6]. This species considered an oligarchic species, exceeding its natural production at $11.1 \text{ t ha}^{-1} \text{ year}$ [7]. The *Patauá* fruit has a purple color and weighs between 6-8 grams [8]. *Patauá*, presents numerous food applications, highlighting or *Patauá* wine, being an abundant source of rent, as well as a production of oil, which shows flavor and properties similar to olive oil, as well as manufacturing of soda and ice cream [9]. The oil extracted from *Patauá* presents massive quantities of unsaturated fatty acids [10]. The most important characteristic of this type of oil, and a high concentration of oleic acid and low level of palmitic acid, characteristic of other palm oils, as well as presenting applications to the cosmetics industry and not treating certainties [11]. Since there is work related to the fruit of *Patauá* a palm, this study aimed to study the chemical, nutritional composition and bioactive molecules in pulp and *Patauá* seed with

occurring in the state of Roraima, in the Northern Amazon of Brazil.

2. MATERIALS AND METHODS**2.1 Collection and Processing of Samples**

The *Patauá* samples were collected in Rorainópolis city in the state of Roraima (Brazil) and were taken to the Chemistry laboratory of the Federal Institute of Roraima, where previously selected were those that had an acceptable conservation state for consumption, and the samples were separated. Seed pulps and freezing at -80°C and lyophilized. The lyophilized material was sieved between 30-40 Mesh and stored in the dark until it was time for analysis.

2.2 Mineral Analysis

The extraction of minerals in the different parts of the fruit was done according to the methodology described by EMBRAPA [12], which uses the perchloric nitric digestion (3:1) in a TECNAL model TE 0079 digester block, washed with distilled water until 25 mL for subsequent analysis. Calcium (Ca), magnesium (Mg), iron (Fe), copper (Cu), zinc (Zn), manganese (Mn), and aluminum (Al) were determined by flame atomic absorption spectrophotometry (FAAS). Shimadzu AA-7000, coupled with ASC-7000 auto sample. Calibration was performed with standard solutions prepared from 1000 mgL^{-1} Qhemis High Purity PACU 1000-0125 commercial standards according to the specific conditions of each element (Table 1).

In the case of sodium (Na), it was determined in the same equipment, but in atomic emission mode. Potassium (K) was evaluated by flame photometry using the Digimed DH-62 Flame Photometer, calibrated using a Digimed standard solution with a concentration range of 2 - 100 mg L⁻¹. For the determination of phosphorus (P) and boron (B), the Ultraviolet-visible spectrophotometry technique (SHIMADZU model UV-1800 equipment) was applied using the methodology of EMBRAPA [12], which the colorimetric reaction was done by ammonium molybdate ((NH₄)₂MoO₄). In the case of P, a blue complex was formed, where the readings were taken at $\lambda = 660$ nm; In the case of B, a complex was formed with yellow azomethine-H and absorbed light at $\lambda = 460$ nm.

2.3 Characterization of Bioactive Molecules

The determination of total phenolic compounds (CFT) was being used gallic acid as a reference standard and absorbance readings in UV-VIS Photonics spectrophotometer at 765 nm [13]. The results were reported as GAE in 100 g⁻¹ of the sample. The antioxidant activity was

performed using two methodologies: on the one hand, the 1,1-diphenyl-2-picrylhydrazine (DDPH) absorption method was used absorbance readings at 515 nm [14]. The second method to evaluate total antioxidant activity was by the process of reducing Fe³⁺ → Fe²⁺ [15].

The absorbance readings made the concentration of total carotenoids, which were taken on UV-VIS molecular absorption spectrophotometer at 661 nm, 644 nm, and 470 nm, respectively [16]. Finally, vitamin C quantification was performed by UV-visible molecular spectrophotometry and recorded at 545 nm [17,18].

3. RESULTS AND DISCUSSION

3.1 Mineral Analysis

Minerals are essential nutrients for the body that cannot be synthesized and are incorporated into the diet [19], consisting of inorganic substances that help regulate body functions [20]. Table 2 shows the results of macro and micro minerals in the *Pataú* pulp and seed, performed by different spectrophotometric techniques.

Table 1. Analytical parameters

Element	Technique	(λ) nm	Calibration Line
Ca	FAAS	422.70	y= 0.0092 x - 0.0005 r ² = 0.999
Mg	FAAS	285.21	y= 0.2353 x - 0.0658 r ² = 0.997
P	UV-Vis	660.00	y= 0.2181 x - 0.0005 r ² = 0.999
K	EAE	766.50	y= 0.1231 x - 0.0013 r ² = 0.993
S	UV-Vis	420.00	y= 0.0213 x - 0.0012 r ² = 0.998
Fe	FAAS	248.33	y= 0.0399 x + 0.0067 r ² = 0.996
Zn	FAAS	213.80	y= 0.0600 x - 0.0171 r ² = 0.991
Mn	FAAS	279.48	y= 0.0282 x + 0.0041 r ² = 0.999
Cu	FAAS	324.75	y= 0.0512 x - 0.0099 r ² = 0.997
Na	EAS	589.00	y= 1.0000 x + 0.0005 r ² = 0.999
Al	FAAS	309.30	y= 0.0088 x + 0.0005 r ² = 0.998
B	UV-Vis	460.00	y= 0.0537 x + 0.0002 r ² = 0.999
Co	FAAS	240.73	y= 0.0286 x - 0.0066 r ² = 0.997

FAAS = Flame Atomic Absorption Spectroscopy. EAS = Flame Atomic Emission Spectroscopy. UV-Vis= UV-visible molecular spectrophotometry

Table 2. Mineralogical composition of *Pataú* pulp and seed

Macrominerals mg 100 g ⁻¹	Ca	K	Mg	Na	P
Pulp	2.35 ± 0.11	2.17 ± 0.07	41.23 ± 0.12	71.21 ± 0.02	41.23 ± 0.12
Seeds	0.71 ± 0.04	3.52 ± 0.04	48.31 ± 0.07	84.21 ± 0.02	1.17 ± 0.02
Microminerals mg 100 g ⁻¹	B	Cu	Fe	Mn	Zn
Pulp	0.37 ± 0.04	0.11 ± 0.01	1.84 ± 0.02	0.61 ± 0.09	0.97 ± 0.12
Seeds	0.22 ± 0.02	0.49 ± 0.07	0.91 ± 0.07	1.10 ± 0.03	0.21 ± 0.07

Average value of three repetitions and standard deviation at 95%

Among the macro-minerals Table 2, sodium and magnesium concentrations with better concentrations for seeds than for pulps stand out, with a value of 84.21 mg, 100 g⁻¹ sodium, 48.31 mg 100 g⁻¹ of magnesium for the seed. Another of the macro-minerals whose concentration is significant in *Patauá* is the phosphorus, being highly superior in the pulp (41.23 ± 0.12 mg 100 g⁻¹) compared to the seed. Phosphorus is an element that is synthesized in the body and is being found in the form of inorganic or organic phosphorus when it is covalently linked to other biomolecules such as proteins, sugars, and other cellular compounds [21].

On the other hand, the other two macro-minerals found in *Patauá* pulp and seed were, firstly, Ca, where the concentration was higher in the pulp with a value of 2.35 mg 100 g⁻¹ and, secondly, potassium with higher value in seed 3.52 mg. 100 g⁻¹. Calcium consumption is of vital importance to humankind since it is the main constituent of bones, as well as involved in intracellular regulation of different tissues [22]. On the other hand, potassium is another of the vitally important elements for the organism since it is responsible, among other functions, for maintaining the hydroelectric balance in cells along with sodium [23]. Mineral requirements vary between 4-15% of body weight, with 50% corresponding to calcium, 25% to phosphorus, and another 25% corresponding to other minerals such as magnesium, sodium, potassium, and copper [24].

The microminerals determined in this work (Table 2), were B, Cu, Fe, Mn, and Cu, they are necessary for health since their deficiencies are related to different diseases [25]. These include iron concentrations in the pulp (1.84 mg 100 g⁻¹) and zinc in the pulp with 0.97 mg 100 g⁻¹. Iron is one of the micronutrients of great importance for the organism. The iron content of 3-5 grams in the body is divided into two categories: the one belonging to functional compounds and the other corresponding to the iron stored in the hepatocytes and cells of the reticulum system-endothelial form of ferritin and hemosiderin [26] being daily iron recommendations for the organism of 11 mg day⁻¹ (men), and 15 mg day⁻¹ for women [27]. Zinc is an element distributed in the body in small proportions in concentrations ranging from 1.5 to 2.5 grams [28], and is an element implicated in the proper functioning of all cells in the body, being essential especially for the development of the immune system [29].

Manganese had higher concentrations for the seed than for the *Patauá* pulp being an essential trace element that participates in different metabolic reactions, involved in immunological responses and regulation of ATP synthesis, as well as crucial enzymatic reactions [30]. Boron is another of the elements identified in *Patauá* with a higher concentration in the pulp with 0.37 mg 100 g⁻¹, being of great importance for the organism in bone growth, prevention of arthritis as well as implicated in hormonal regulation processes with a daily intake of 3 mg day⁻¹ [31].

3.2 Bioactive Molecules

Table 3 shows the results of total phenolic compounds, antioxidant activity, total carotenoids and ascorbic acid for *Patauá* pulp and seeds.

Total phenolic compounds measured in gallic acid equivalents per gram of sample, using standard gallic acid reference standard, were higher in seed with a concentration of 356.12 ± 0.12 mg GAEg g⁻¹ than for the pulp with a level of 321.03 ± 0.43 mg GAEg. g⁻¹. According to the classification for phenolic compounds [32], they present a high concentration of phenolic compounds whose samples are over 500 mg GAEg g⁻¹, and according to this classification, the pulp and *Patauá* seed, have a high level of phenolic compounds. Antioxidant activity, measured by the DPPH technique and iron reduction method, showed higher antioxidant activity for the seed than for the pulp with both evaluated methods. The results obtained with this work are close to those obtained by Rezaire et al. [33], who studies extracts from *Euterpe oleracea* with GAE concentrations of 306.6 ± 7.4 mg GAEg g⁻¹ and antioxidant activity values determined by different methods within those surveyed in this work. As with other widely studied palm trees in this family, *Euterpe oleracea* has an interesting antioxidant potential, especially in vitro, and its antioxidant potential, which is related to the polyphenolic compounds such as anthocyanins (water-soluble pigments, responsible for the color purple) as well as other flavonoids [34].

The concentration of carotenoids in *Patauá* stands out, whose concentration is higher for seed than for pulp (Table 3), being a group of precursor bioactive molecules of vitamin E. The values obtained for carotenoids were 0.26 mg mL⁻¹ for the pulp and these values were higher

Table 3. Bioactive molecules in pulp and seeds of *Patauá*

Sample	Phenolic Compounds mg GAEq g ⁻¹	Antioxidant activity μmol g ⁻¹		Carotenoids μg mL ⁻¹	Vitamin C mg mL ⁻¹
		DPPH	Reduction Fe		
Pulps	321.03 ± 0.43	2147.12 ± 22.34	1856.21 ± 11.23	0.26 ± 0.02	0.008± 0.001
Seeds	356.12 ± 0.12	2632.21 ± 21.25	2123.12 ± 21.23	2.52 ± 0.04	0.005± 0.001

Average value of three repetitions and standard deviation at 95%

for the seed with a concentration of 2.52 mg mL⁻¹. Other fruits of the same family, such as the *Astrocaryum aculeatum* pulp case, presented carotenoid levels between 3.5-4.3 mg 100 g⁻¹ [35] and 0.4 mg 100 g⁻¹ for *Attalea maripa* [36]. Another type of compound with antioxidant activity is vitamin C, which was detected at low concentrations in *Patauá* pulp and seed Table 3, which its deficiency can cause scurvy [37].

4. CONCLUSION

In this work, it was placed the biotechnological importance of this Amazon palm in terms of its mineralogical composition, as well as the characterization of bioactive molecules, especially the phenolic compounds in the disposable parts such as the seed, a disposable part of the fruit that could be of interest to the recovery bioactive molecules containing these bio-residues at industrial level. Also, noteworthy are carotenoids, which can be used in the pharmaceutical and cosmetic industry. It is a species still little explored and can be used as a raw material in the food industry, since until now its use is limited to the local populations of the amazon.

ACKNOWLEDGEMENTS

The authors express their acknowledgements to the Federal Institute of Roraima and the FINEP to Brazil.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Trias-Blasi A, Baker WJ, Haigh AL, Simpson DA, Weber O, Wilkin P. A genus-level phylogenetic linear sequence of monocots. *Taxon*. 2015;64(1):552–581.

2. Govaerts R, Dransfields J. World checklist of Palms. Royal Botanic Gardens, Kew, Richmond, UK; 2005.
3. Eiserhardt WL, Svenning JC, Kissling WD, Balslev H. Geographical ecology of the palms (*Aracaceae*): Determinants of diversity and distributions across spatial scales. *Ann. Bot.* 2011;108:1391-1346.
4. Montúfar R, Laffargue A, Pintaud J-C, Avallone SHS, Dussert S. *Oenocarpus bataua* Mart. (*Arecaceae*): Rediscovering a source of high oleic vegetable oil from Amazonia. *Journal of the American Oil Chemists' Society*. 2010;87:67-172.
5. Gomes-Silva DAP, Wadt LHO, Ehringhaus C. Ecologia e manejo de *Patauá* (*Oenocarpus bataua* Mart.) para produção de frutos e óleo. *Embrapa Acre. Documentos*, 88. Rio Branco-AC; 2004.
6. Lorenzi. 2004. *Arvores brasileiras: Manual de identificação e cultivo de plantas arbóreas nativas do Brasil*, Plantarum: Nova Edesa. 2002;2.
7. Ruiz RR, Alencar JC. Comportamento fenológico da palmeira de *Patauá* (*Oenocarpus bataua*) na reserva florestal Adolpho Ducke, Manaus, Amazonas, Brasil. *Acta Amazônica*. 2004;34:553-558.
8. Rezaire A, Robinson JC, Bereau D, Verbaere A, Sommerer N, Khan MK, Durand E, Fils-Lycaon BF. Amazonian palm *Oenocarpus bataua* (Patawa): Chemical and biological antioxidant activity-Phytochemical composition. *Food Chemistry*. 2014;149:62-70.
9. Miranda IPA, Rabelo A, Bueno CR, Barbosa EM, Ribeiro MNS. *Frutos de Palmeiras da Amazônia*. MCT-INPA; 2001.
10. Shanley P, Medina G. *Frutíferas e Plantas Úteis na vida amazônica*. Imazon; 2005.
11. Teixeira D, Da Rocha, GN. Extração e caracterização do óleo de patuá (*Oenocarpus batua* Mart) e comparação de suas propriedades com o óleo de oliva. *Doctoral Thesis*. Federal University do Pará, Belém; 2001.

12. Empresa Brasileira de Pesquisa Agropecuária-EMBRAPA- Manuel o chemical analyzed of soils, plants and fertilizers. 2 Edition, Brasilia; 2009.
13. Wolfre K, Wu X, Liu RH. Antioxidant activity of apple peels. *J. Agric Food Chem.* 2003;51:609-614.
14. Miranda IPA, Rabelo A, Bueno CR, Barbosa EM, Ribeiro MNS. Frutos de Palmeiras da Amazônia. MCT-INPA; 2001.
15. Sánchez-Moreno C, Larrauri JA, Saura-Calixto F. A procedure to the antiradical efficiency of polyphenols. *Journal of the Science of Food and Agriculture.* 1998;76: 270-276.
16. Lichtenthaler HK, Buschmann C. Chlorophylls and carotenoids: Measurement and characterization by UV-VIS Spectroscopy. *Current protocols in Food Analytical Chemistry.* 2001;4:3-4.8.
17. Badolato MICB, Sabino M, Lamardo LCA, Antunes JLF. Comparative study of analytical methods for the determination of an ascorbic acid in the success of natural and industrialized fruits. *Ciênc. Tecnol. Aliment.* 1996;16:206-210.
18. Contreras-Gúzmán E, Strong IFC, Guernelli O. Determination of ascorbic acid (vitamin C), by reduction of copper ions. *Química Nova.* 1984;7:60-64.
19. Kessenich C. Alternative Choices for calcium supplementation. *The Journal for Nurse Practitioners.* 2008;4:36-39.
20. Roach S. Promovendo a saúde fisiológica. In: *Enfermagem na saúde do idoso.* 4 Ed. Rio de Janeiro: Guanabara Koogan; 2009.
21. Monteiro TH, Vannucchi H. Funções Plenamente reconhecidas de nutrientes Fósforo. São Paulo: Série de Publicações ILSI Brasil; 2010.
22. De França NA, Martini LA. Funções Plenamente reconhecidas de nutrientes Cálcio. São Paulo: Série de Publicações ILSI Brasil; 2014.
23. Cuppari L, Bazanelli AP. Funções Plenamente reconhecidas de nutrientes Potássio. São Paulo: Série de Publicações ILSI Brasil; 2010.
24. Caceres E, Garcia ML, Selgas ML. Design of a new cooked meat sausage enriched with calcium. *Meat Science.* 2006;73:368-377.
25. Smolin LA, Grosvenor MB. *Nutrition: Science and applications with bloklet package.* 1 Ed. Orlando: John Wiley & Sons Inc; 2007.
26. Fisberg M. Funções Plenamente reconhecidas de nutrientes Ferro. São Paulo: Série de Publicações ILSI Brasil; 2014.
27. Dietary Reference Intake for vitamin A, Vitamin K, Arsenic, boron, chromium, Copper, Iodine, Iron, Manganese, molybdenum, nickel, silicon, vanadium and Zinc. Capítulo 13. Washington: The National Academic Press; 2001. Available:<https://www.nap.edu/read/10026/chapter/15>. [Acesso em: 14 jan. 2019]
28. Hambidge MK. Dietary reference intakes for Zinc may require adjustment for Phytate intake based upon model Predictions. *Journal Nutrition.* 2008;138: 2363–2366.
29. Freitas EC, Silva ACM, da Silva MV. Análises de minerais zinco e manganês presentes na farinha do morango. *Revista Brasileira de Obesidade, Nutrição e Emagrecimento.* 2016;10:303-307.
30. Burton NC, Guilarte TR. Manganese neurotoxicity: Lessons learned from longitudinal studies in nonhuman primates. *Environ Health Perspect.* 2009;117:325-332.
31. Nielsen FH. Should bioactive trace elements not recognized as essential, but with beneficial health effects, have intake recommendations. *Journal Trace Element Medicine Biology.* 2014;28:406-408.
32. Vasco C, Ruales J, Kamal-Eldin A. Total phenolic compounds and antioxidant capacities of major fruits from Ecuador. *Food Chemistry.* 2008;111:816-823.
33. Rezaire A, Robinson JC, Bereau D, Verbaere A, Sommerer N, Khan MK, Durand E, Fils-Lycaon BF. Amazonian palm *Oenocarpus bataua* ("Patawa"): Chemical and biological antioxidant activity-Phytochemical composition. *Food Chemistry.* 2014;149:62-70.
34. Kang J, Li Z, Wu T, Jensen GS, Schauss AG, Wu X. Anti-oxidant capacities of flavonoid compounds isolated from açai pulp (*Euterpe oleracea* Mart.). *Food Chemistry.* 2010;122:610–617.
35. Aguiar JPL, Marinho HA, Rebêlo YS, Shrimpton R. Aspectos nutritivos de alguns frutos da Amazônia, *Acta Amazonica.* 1980;10:755–758.

36. Santos MDFG, Mamede RVS, Rufino MDSM, de Brito ES, Alves RE. Amazonian native palm fruits as sources of antioxidant bioactive compounds. *Antioxidants*. 2015; 4:591–602.
37. Chin KY, Ima-Nirwana S. Vitamin C and bone health: Evidence from cell, animal and human studies (Review). *Current Drug Targets*. 2018;19:439-450.

© 2019 Saravia et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/54389>