



The Effect of Indomethacin on the Sperm Motility of Adult Wistar Rats

Leelee F. Zitte^{a*} and Prisca I. Uzodinma^a

^a *Department of Animal and Environmental Biology, Faculty of Science, University of Port Harcourt, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJOB/2023/v17i4330

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98312>

Original Research Article

Received: 01/02/2023

Accepted: 03/04/2023

Published: 11/04/2023

ABSTRACT

Indomethacin is a non-steroidal anti-inflammatory drug (NSAID) used mostly in the treatment of pain, feverish condition, headaches and inflammation. This study was therefore carried out to evaluate the effect of indomethacin on the male reproductive system, focusing on sperm quality. Twenty male rats were divided into four groups, (groups 1, 2, 3, and 4) with group one serving as control. The other three groups receiving different doses of indomethacin, (groups 2-4 receiving 1mg/kg, 2mg/kg and 4mg/kg respectively) for 21 days. After the period, sperm samples were collected from all the groups and analyzed to evaluate sperm quality. The sperm qualities analyzed include; sperm motility, semen viscosity, sperm morphology, sperm count, etc. the results of the study indicated that indomethacin had a negative impact on the sperm qualities of the rats, as it decreased sperm count and sperm motility, and had a detrimental effect on the overall sperm quality. This study provides important insights into the potential side effects of indomethacin on male fertility. While its therapeutic benefits are well established, its impact on male fertility has not been documented. The findings of this study suggest that men who use indomethacin regularly may be at a higher risk of experiencing negative impacts on their reproductive health, including reduced sperm quality.

*Corresponding author: E-mail: leelee.zitte@uniport.edu.ng;

Keywords: Indomethacin; wistar rat; sperm motility; reproductive health.

1. INTRODUCTION

Indomethacin is a non-steroidal anti-inflammatory drug (NSAID) that is very effective in the treatment of pain, feverish conditions, and inflammation, like other NSAIDs it reduces pain and inflammation [1]. Generally, non-steroidal anti-inflammatory drugs (NSAIDs) are a diverse group of drugs, frequently unrelated chemically, yet the majority are organic acids and known in the treatment of conditions, such as osteoarthritis, rheumatoid arthritis, different type of inflammation, gout, menstrual cramps, dental pain, and headache [2]. However numerous adverse reactions have been linked to NSAIDs use which include nausea, vomiting, diarrhea, constipation, reduced appetite, rash, dizziness, headaches, sleepiness, and stomach ulcers [1,3,4]. Indomethacin is used mainly in the treatment of moderate pain since 1960 and it is frequently prescribed for the treatment of severe gouty arthritis pain, it has also shown effectiveness in the treatment of several other painful disorders [5,6].

The chemical composition of indomethacin is 1-(p-chlorobenzoyl) 25-methoxy-2-methylindole-3-acetic acid, it is effective mainly because of the ability of indomethacin to inhibit prostaglandin production. Prostaglandins are important mediators of inflammation, fever, and pain and are predominantly generated by cyclooxygenase (COX) enzymes [7].

The adverse effect of common NSAIDs like ibuprofen and Naproxen has been studied and patients are advised not to take NSAIDs on an empty stomach because of its ability to cause stomach ulcers. The adverse effect of indomethacin is one of the reasons why it is not generally prescribed clinically, like the way it works the adverse effects occur by non-selectively inhibiting the cyclooxygenase (COX1) and (COX2) enzymes, which prevent the synthesis of prostaglandins from arachidonic acid by the cyclooxygenase enzymes and the cyclooxygenase is necessary for physiological processes like the maintenance of gastrointestinal mucosa and vascular homeostasis [8,9].

Though this drug is commonly used in treating various forms of pain and inflammation, indomethacin has been found to have potential adverse effects on reproductive health. Although

previous studies have suggested a link between the use of indomethacin and decreased sperm quality, the relationship between the drug and specific aspects of sperm health such as sperm motility, count, and morphology, has not been well established [3]. This lack of clarity regarding the specific impact of indomethacin on sperm quality is a major limitation in our understanding of the drug's potential effects on male fertility.

Given the widespread use of indomethacin and the increasing interest in understanding the impact of the medications on reproductive health, there is a pressing need for more information on the effect of indomethacin on sperm quality. The purpose of this study is to fill the gap by examining the effect of indomethacin on sperm quality in a controlled laboratory setting. The result of this study will help clarify the potential impact of indomethacin on sperm quality and male fertility.

2. MATERIALS AND METHODS

This research work was carried out in the Department of Animal and Environmental Biology animal house University Campus, University of Port Harcourt Choba, Port Harcourt Rivers State, Nigeria. In December 2023.

Indocid was bought from a local pharmacy at Choba Port Harcourt Rivers State Nigeria. The materials and specimens used in carrying out the experiment included Adult male rat, cages, Rat feed, drinkers, hand gloves, desiccator, dissecting board, chloroform, cotton, microscope, normal saline, stains (methylene blue, alcohol and strong carbon fischen), Neubauer counting chamber, weighing balance, pipette, plain bottle, surgical blade.

Twenty (20) rats were gotten from the Faculty of Basic Medical Sciences, Department of Pharmacology, University of Port Harcourt, and they were acclimated for 7 days. The 20 rats were placed into 4 groups. Their weight ranges from 180g to 228g. The animals were maintained under the standard condition of twelve hours' light/dark cycles at room temperature and fed with grower's mash feed and water.

Indocid was diluted serially into 1mg/kg (lowest dose), 2mg/kg, and 4mg/kg (highest dose). Group one (1) was used as the control group, while the other three (3) groups were treated with

1mg/kg, 2mg/kg and 4mg/kg of indocid respectively.

After 21 days of the administration, the semen was collected from the epididymis of the rats of each group, for microscopic examinations.

The semen is usually hyper-viscous in nature, hence it was placed in a small plain bottle, and 2 drops of normal saline were added to separate the cells. A drop of the diluted semen was placed in a slide to check for sperm motility microscopically using the objective lens 10x.

A drop of dilute semen was taken and placed on a plain bottle, 19 drops of normal saline was added to achieve the 1:20 dilution. This dilution was charged on the counting chamber; after which it was placed on the moisturizer for 3 minutes to allow more separation of the cells. The counting chamber was then placed on the microscope and viewed using the objective lens of x10. The cells were counted using the hand tally counter. The formula for calculating the sperm count, when 5 small squares within the large center square are counted is: Number of sperm counted x dilution factor/volume x 1000 = sperm/ml. Example: 50 sperm are counted in the five small squares. The dilution is 1:20.

Sperm/ml = $50 \times 20 / 0.02 \text{ mm}^3 \times 1000 \text{ mm}^3/\text{ml} = 50,000,000 \text{ sperm/ml}$.

If the dilution is 1:20, and the usual 5 small squares are counted, then the formula can be simplified as follow: Sperm/ml = Sperm counted x 1,000,000.

To check for defects in the sperm morphology (head, midpiece, and tail defects), a clean slide was placed on the microscope and a drop of the diluted semen was added, thereafter, a film was made out of it, and then it was flooded with alcohol and allowed to be air dried for 2 minutes. Strong carbon fischen was latter used to flood and further allowed to stand for 2-3 minutes before it was washed off with alcohol and then viewed under x100 oil immersion lens of the microscope.

3. RESULTS

3.1 Active Sperm Cells

There were more active sperm cells in Group 1 (the control) (84%) compared to the treatment groups (62.75, 53.25 and 63.00 % for the three treatment groups respectively). Group 3 with a dose of 2mg/kg had the lowest number of active sperm cells. The results showed that indomethacin affected the sperm cells because the control group had more active sperm cells than the administered groups (Fig. 1).




Fig. 1. The Active sperm cells across the different Groups

3.2 Sluggish Sperm Cells

The treated groups had more sluggish sperm cells compared to the control, and Group 3 amongst the treated groups had the highest number of sluggish sperm cells. Indomethacin had a significant effect on the sperm cells motility because the treated groups have more sluggish sperm cells than the control (Fig. 2).

3.3 Dead Sperm Cells

The amount of dead sperm present in Group 3 was more than any in every group and there are more dead sperm cells in the groups where administration occurred compared to the control group (Group 1). The percentage dead cell were

11.25%, 26.25%, 48.75% and 30.00% for groups 1, 2, 3 and 4 respectively.

The administration of indomethacin leads to the death of spermatozoa as the results showed that the treated groups had more dead sperm cells than the control. (Fig. 3).

3.4 Percentage of Sperm Viability

Group 1 (control) had highly viable sperm cells and very low non-viable sperm cells and when compared to the other groups, the control group has better sperm viability than any other groups. The control has more sperm viability, so indomethacin affects the viability and the administered groups had lower viability (Fig. 4).




Fig. 2. The Sluggish cells across different group




Fig. 3. The mean of dead Sperm cells across the different group




Fig. 4. The Sperm Viabilities across the different Groups




Fig. 5. Comparison of viable cells with nonviable and dead cell (affected cells)

3.5 Viable versus Affected Sperm Cells (Non Viable and Dead)

Comparism of the viable sperm cells and the nonviable and dead shows that group one (control) 74.55 % viable as against 25.25% non viable and dead combined ratio of 46.28% and 53.72%, 27.29%and 72.71%, and 44.1% and 55.9% for groups 2, 3 and 4 respectively.

This implies that he indocid directly affect the viability of the sperm and hence affecting reproductive health of rats which will also affect humans reproductive health.

3.6 Sperm Count

Effects of idocid was noticed on the sperm count distribution of the different groups. The control had total count of 192 million per ml, while groups 2,3, and 4 have 162.5 million, 137.5 million and 56.25 million respectively. Group 4

was observed to be so affected in sperm count hence there were relatively fewer sluggish and dead cells. (Fig. 6).

4. DISCUSSION

The study of the effects of indomethacin on sperm quality has generated significant interest in the scientific community due to the increasing number of couples facing infertility issues. The results of the study have shown that indomethacin administration has a detrimental effect on sperm quality. This is an important finding as it highlights the need for caution when using this medication in individuals of reproductive age.

The increase in sluggish and dead cells of group 3 above groups 4 could be traceable to the sperm count of group 4 which was drastically affected by the treatment. But in the case of group 3, sperm count was higher so many dead and sluggish sperm were noticed.




Fig. 6. Effects of indocid on sperm count of the different groups

The results of the study showed that the control group had a higher number of active sperm cells compared to the groups where indomethacin administration occurred. This finding is consistent with previous research that has found that nonsteroidal anti-inflammatory drugs (NSAIDs) can decrease sperm motility [10]. The decrease in active sperm cells may be due to the fact that indomethacin inhibits the production of prostaglandins, which are important for maintaining sperm motility.

However, the study also found out that the administration of indomethacin led to an increase in the number of sluggish sperm cells. This finding is also consistent with previous research that has shown that NSAIDs can negatively impact sperm function [11]. The increase in sluggish sperm cells suggests that indomethacin may be affecting the energy metabolism of sperm cells, leading to a decrease in their ability to move efficiently.

The results also showed that the administered groups had a higher number of dead sperm cells compared to the control group. This finding is consistent with previous research that has shown that NSAIDs can increase oxidative stress and decrease sperm viability. The increased amount of dead sperm cells suggests that indomethacin may be affecting the antioxidant defense mechanisms of sperm cells, leading to an increase in oxidative stress and a decrease in sperm viability which correlated with Sabeeh et al. [8] who investigated the histopathological changes in the male reproductive organs of rats treated with oral indomethacin, with the aim of understanding the potential negative effects of the drug on male reproductive health. The study

found that rats treated with indomethacin exhibited a range of histopathological changes in their male reproductive organs, including the reduced thickness of seminiferous tubules, decreased numbers of spermatogenic cells, and increased numbers of degenerating cells. The study also reveals decreased weight of seminal vesicles in the treated group.

Finally, the study found that the control group had more viable sperm cells than the treated groups. This finding is also consistent with previous research that has shown that NSAIDs can decrease sperm viability [3]. The decrease in sperm viability may be due to the fact that indomethacin affects the membrane integrity of sperm cells, leading to a decrease in their ability to survive [12]. The results of this study provide further evidence that indomethacin administration can have a negative impact on sperm quality. This is an important finding as it highlights the need for caution when using this medication in individuals of reproductive age though may be effective for managing pain and inflammation. It is recommended that further research be conducted to better understand the mechanisms by which indomethacin affects sperm quality and to determine the optimal dose and duration of use that minimizes its negative impact on sperm quality [13,14].

5. CONCLUSION

The study has shown that indomethacin has a detrimental impact on sperm quality. The results showed that the control group had a higher number of active sperm cells compared to the group where indomethacin was administered. The administration of indomethacin led to an

increase in the number of sluggish sperm cells and dead sperm cells and a decrease in the number of viable sperm cells. These findings highlight the need for caution when using this medication in individuals of reproductive age. The results of the study provide further evidence that indomethacin administration can negatively impact sperm quality.

6. RECOMMENDATIONS

Based on the results of the study, it is recommended that individuals of reproductive age should carefully consider the use of indomethacin. If possible, alternative pain management options should be considered and it is recommended that further research be conducted to better understand the mechanisms by which indomethacin affects sperm quality and to determine the optimal dose and duration of use that minimizes its negative impact on sperm quality.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lucas S. The pharmacology of indomethacin. *Headache: The Journal of Head and Face Pain*. 2016;56(2):436-446.
2. Gliszczyńska A, Nowaczyk M. Lipid formulations and bioconjugation strategies for indomethacin therapeutic advances. *Molecules*. 2021;26(6):1576.
3. Bagoji IB, Doshi MA, Yendigeri S, Patil BG, Patil BB, Das KK. Black tea extract ameliorates indomethacin induced changes in testicular histopathology of albino rats. *Journal of Young Pharmaceutics*. 2014;6(3):53.
4. Basivireddy J, Jacob M, Pulimood AB, Balasubramanian KA. Indomethacin-induced renal damage: role of oxygen free radicals. *Biochemical pharmacology*. 2004;67(3):587-599.
5. Hart Fd, Boardman PI. Indomethacin: A new non-steroid anti-inflammatory agent. *Br Med J*. 2003 Oct 19 ;2(5363):965-70.
6. Nalamachu S, Wortmann R. Role of indomethacin in acute pain and inflammation management: a review of the literature. *Postgraduate Medicine*. 2014;126(4):92-97.
7. Ghosh R, Prosad Banik S. Protective effect of indomethacin on vanadium-induced adrenocortical and testicular damages in rat. *Toxicology Mechanisms and Methods*. 2022;32(2):114-122.
8. Sabeeh SI, Al-Naimi RA, Yousuf EH. Histopathological changes of male reproductive organs treated with oral administration of indomethacin in male rat. *Al-Anbar Journal of Veterinary Sciences*. 2014;7(2).
9. Bernabeu R, Roca M, Torres A, Ten J. Indomethacin effect on implantation rates in oocyte recipients. *Human Reproduction*. 2006;1(2):364-369.
10. Padrón RS, Nodarse M. Effect of indomethacin on semen of infertile men. *International Journal of Andrology*. 1979;2(1-6):110-116.
11. Martini AC, Vincenti LM, Santillán ME, Stutz G, Kaplan R, Ruiz RD, de Cuneo MF. Chronic administration of nonsteroidal anti-inflammatory drugs (NSAIDs): effects upon mouse reproductive functions. *Revista de la Facultad de Ciencias Médicas de Córdoba*. 2008; 65(2):41-51.
12. Ratnasooriya WD, Wadsworth RM. Effect on fertility in rats of epididymal administration of indomethacin. *International Journal of Fertility*. 1987; 32(2):152- 156.
13. Indomethacin-induced renal damage: role of oxygen free radicals. *Biochemical Pharmacology*. 2004;67(3):587-599.
14. Summ O, Evers S. Mechanism of action of indomethacin in indomethacin-responsive headaches. *Current Pain and Headache Reports*. 2013;17:1-7.

© 2023 Zitte and Uzodinma; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/98312>