



# **Prescription Audit and Dispensing Errors in the Outpatient Pharmacy of Tertiary Care Multispecialty Teaching Hospital**

**T. Irsath Ahamed<sup>a</sup>, K. Vaishnavi<sup>a</sup>, R. R. Harini<sup>a</sup>, Likhita Yalavarthi<sup>a</sup>  
and Jagadeesan M.<sup>a\*</sup>**

<sup>a</sup> Department of Pharmacy Practice, SRM College of Pharmacy, SRM Institute of Science and Technology, SRM Nagar, Kattankulathur, Chengalpattu-603203, Tamil Nadu, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author TIA designed the data collection form, also identified and documented dispensing errors. Author KV contributed to obtaining Institutional Ethical Committee (IEC) approval and wrote the manuscript. Author RRH edited the article and reviewed the literature. Author LY performed the area of citations, references and final editing of the manuscript. Author JM selected the topic, guided and supervised throughout the progress of the research work. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JPRI/2021/v33i54B33790

Editor(s):

(1) Dr. Juan Carlos Troiano, University of Buenos Aires, Buenos Aires, Argentina.

Reviewers:

(1) Shoaeb Mohammad Syed, Dr. Vedprakash Patil Pharmacy College, India.

(2) Kapil Kalra, Alpine College of Management & Technology, India.

(3) Ilker Ilhanlı, Ondokuz Mayıs University, Turkey.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/76861>

**Original Research Article**

**Received 01 October 2021**

**Accepted 03 December 2021**

**Published 13 December 2021**

## **ABSTRACT**

**Background:** Although errors occurring during the process of dispensing may affect the goal of the treatment, they also can be the significant cause of morbidity and mortality. There are only few published evidences which focuses on the errors of dispensing that occurs in the pharmacy. This study focuses on identifying the dispensing errors, impact of brand substitution on cost and DDIs.

**Objective:** To identify the dispensing errors, impact of brand substitution on cost and DDIs.

**Methodology:** Prospective observational study conducted over a period of eight weeks in outpatient hospital pharmacy of tertiary care multispecialty teaching hospital, Tamil Nadu. Prescriptions and medication bills and dispensed medicines are collected from the hospital

pharmacy to obtain data required for the study. Prescription containing only medical devices were excluded from the study.

**Results:** Out of 1010 prescriptions, dispensing errors were present in 419 (41.48%) prescriptions which consist of 557 errors. The errors include dispensing multi-pills to make the required dose is 3.77%, tablet splitting is 0.8%, incorrect strength is 1.07%, omission error due to unavailability is 31.4%, dispensing drugs with brands other than prescribed brands is 63.7%. Prescriptions which had more than three drugs were analyzed for drug-drug interactions (n=389). DDIs were present in 156(40.1%) prescriptions which had a total of 281 interactions.

**Conclusion:** The brand substitution and omission errors are the major causes of dispensing errors. Brand substitution is not always recommended as it may have some adverse effects because of salt and excipients variation. Pharmacists are in the position to identify and reduce DDIs by discussing with the physicians.

**Keywords:** Brand substitution; dispensing errors; cost effectiveness; drug-drug interactions; omission error.

## 1. INTRODUCTION

Medication errors are common, preventable errors but they are the important source of errors that may lead to morbidity and mortality in the health care settings [1,2]. They can occur at any time during drug therapy such as prescribing, dispensing, administering, or monitoring [3]. Dispensing errors may occur during dispensing drugs by pharmacists, which may affect the goal of the therapy and may increase the chance of morbidity and mortality [4,5]. The pharmacy staffs are primarily responsible for this error and they can occur at any stage during the process of medication dispensing. A dispensing error is defined as the discrepancy between the written order in a prescription and the medications delivered to the patients by the pharmacists [5-10].

The different types of dispensing errors reported worldwide which include dispensing of wrong drugs, dosage, formulation, quantity, failure to supply the drug, labeling error (includes an error in the drug name, strength, directions and warnings, quantity, patient name, and completely wrong label) [10,11-21]. The hospital pharmacy is the place of origin of medication errors and potential adverse drug reactions [2,22].

Drug-Drug Interactions (DDIs) are defined as the influence of the drug on one another in pharmacokinetics or pharmacodynamics, which may result in unwanted effects, decreased efficacy and/or increased toxicity [23]. They are the significant cause of preventable adverse clinical outcomes [24]. Drug-drug interactions are one of the easily preventable errors among medication errors. Usually, more than one drug is used for treating a disease that carries an increased risk of DDIs with serious health

consequences. There are some factors that could raise the potential negative effect of drug interactions. They are patient's age, number of underlying disease conditions, or drug administration with a low therapeutic index [25]. As many of the physicians may not aware of potential interactions of drugs, it is the responsibility of the pharmacist to avoid dispensing of drugs together that may cause serious drug-drug interactions [26].

There are several patients harmed by medication errors and the studies are mainly concentrating on administration and prescribing errors while dispensing errors are also an important cause of producing harmful effects [27]. As there is only a limited number of studies that focus on the dispensing errors occurs in pharmacy, this study focuses on identifying the rate of dispensing errors, impact of brand substitution on cost, and DDIs [28].

## 2. METHODOLOGY

A prospective observational study was conducted for the period of eight weeks at the hospital pharmacy department of tertiary care multi-specialty teaching hospital, Tamil Nadu. The patient's prescriptions, dispensed medications and medication bills are collected for auditing. Inclusion of all the prescriptions irrespective of the department received from the hospital pharmacy and prescriptions that had only medical devices were excluded from the study.

### 2.1 Statistical Analysis

The data obtained were analyzed using Microsoft Excel® 2016 software and the results have been expressed in percentages and shown in tables and charts.

### 3. RESULTS AND DISCUSSION

Of the total 1060 prescriptions collected, 50 were excluded as it had only medical devices. Finally, 1010 prescriptions were included, in which 41.4% of prescriptions had dispensing errors which consists of about 557 errors. The dispensing errors were identified and classified as in Fig. 1. The total dispensing error rate was 16.45 % which was low when compared to the Beso Adnan et al. study [27]. However, most of these errors are due to brand substitution which may cause adverse effects because of the varying salt forms and it may also cause cost variation [29,30]. In content errors, omission errors were high (31.40%) which was associated with the unavailability of drugs at the time of dispensing. This result is similar to the study of Azevedo Tania Anacleto et al study which was done in Brazil [31].

The total number of drugs prescribed in 1010 prescriptions was 3384. Out of which, only 25% of drugs were prescribed in generic name whereas the remaining were prescribed in the brand name. Due to the unavailability of the

prescribed brand, the pharmacist had to dispense the alternative brand available in the pharmacy. Though the active ingredient did not vary, this alteration may increase or reduce the total cost of medications [29]. Hence, we have analysed the cost variation of prescribed brands and dispensed brands. The difference in cost and the percentage cost variation was calculated using the formula,

$$\text{Cost variation} = \text{Dispensed brand cost} - \text{Prescribed brand cost}$$

$$\text{Percentage of cost variation} = (\text{Cost variation} / \text{Prescribed cost}) * 100$$

It was found that 353 drugs dispensed in other brands, the cost difference of 23 (6%) drugs was found to be zero. The percentage cost variation of 137 (39%) drugs was 22.1% which indicates the dispensed brand cost is higher than the prescribed brand and the percentage cost variation of other 193 (55%) drugs was -21.6% (the negative sign indicates that the cost of dispensed brands is less than the prescribed brands) as given in Table 1.





Fig. 1. Dispensing errors identified and classified

Table 1. Cost variation analysis of prescribed and dispensed brands

No. of Drugs n=353 (%)	Sum of all Dispensed Brand Medications Cost (INR)	Sum of all Prescribed Brand Medication Cost (INR)	Difference in cost (Indian rupee)	Percentage of cost variation
23 (6)	450.52	450.52	0	-
137(39)	3045.78	2494.53	551.25	22.1
193(55)	7119.98	9080.48	-1960.5	-21.6

**Table 2. Prescriptions containing drug-drug interactions**

S. No.	Department	Total No. of Prescriptions (n= 389)	Prescriptions with drug-drug interactions (%)	Prescriptions without drug-drug interactions (%)
1	General Medicine	113	53 (46.9)	60 (53.1)
2	Cardiology & Cardiothoracic	50	36 (72)	14 (28)
3	Dermatology	27	6 (22.23)	21 (77.78)
4	ENT	27	4 (14.81)	23 (85.19)
5	Septic Ward	5	1 (20)	4 (80)
6	Surgery & ICU	47	16 (34.04)	31 (65.96)
7	Nephrology	9	6 (66.67)	3 (33.33)
8	Neurology	17	5 (29.41)	12 (70.59)
9	Orthopedics & Rheumatology	43	16 (37.21)	27 (62.79)
10	Respiratory	9	2 (22.22)	7 (77.78)
11	OBG	9	2 (22.22)	7 (77.78)
12	Pediatrics	7	0(0)	7 (100)
13	Urology	14	4 (28.57)	10 (71.43)
14	Psychiatry	12	5 (41.67)	7 (58.33)

**Fig. 2. Numbers of DDI in the prescription analyzed**

The prescriptions which contain more than three drugs were analyzed for DDIs. Medscape drug interaction checker (v1033.0), Stockley's drug interaction book (12th edition), Micromedex drug interaction software (v2130) were used to identify drug-drug interactions in the prescriptions. 389 prescriptions were examined for DDIs. A total number of 281 interactions were identified in 156 prescriptions (40.1 % from 389) as shown in (Table 2 & Fig. 2), which was less compared to the study of Ajay Chandra et al. [32]. More number of drug interactions was found in Cardiology (72%) and Nephrology (66.6%), there were no drug interaction found in the prescriptions collected in the pediatrics department.

The DDIs found were classified into mild, moderate and severe and it was found that 55

interactions (19.57%) present were mild, 155 (55.16%) were moderate and 71(25.26%) were severe. The moderate drug interactions were found to be high in our study, which was similar to the Ajay Chandra et al study [32] where the rate of moderate DDI is 74.37%.

#### 4. CONCLUSION

In this study, the brand substitution and omission errors are the major causes of dispensing errors which was mainly due to the non-availability of prescribed medications. Brand substitution is not always recommended as it may have some adverse effects because of salt and excipients variation. Pharmacists should make sure that the drugs given in hospital formulary are available at all the time. Dispensing alternative brands are only recommended when the brand has similar

active ingredient, same pharmaceutical form, cost effective, only after getting approval from the physician. Pharmacists are in the position to identify and reduce the drug-drug interactions by discussing with the physicians and providing counseling to the patients.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

Institutional ethical committee [IEC] approval was obtained 1871/IEC/2019 for the study on 13 December 2019.

## ACKNOWLEDGEMENT

We would like to thank, Dean, Head of the Department of Pharmacy Practice, SRM College of Pharmacy and Chief Pharmacist and all the pharmacists, SRM Medical College Hospital and Research Centre.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Brennan TA, et al. Incidence of adverse events and negligence in hospitalized patients. Results from the Harvard Medical Practice Study *N Engl J Med.* 1991;324: 370–376.
2. Bates DW, et al.: Incidence of adverse drug events and potential adverse drug events: Implications for prevention. ADE Prevention Study Group. *JAMA.* 1995;274: 29–34.
3. Rheinskin PH, McGinnis TJ. Medication errors. *Am Fam Physician;* 1992.
4. US Food and Drug Administration. Medication Errors. Available:<http://www.fda.gov/drugs/drugsafety/medicationerrors>.
5. van den Bemt PM, Egberts AC. Drug related problems: Definitions and classification. *Eur J Hosp Pharm Pract.* 2007;13:62-4.
6. Beso A, Franklin BD, Barber N. The frequency and potential causes of dispensing errors in a hospital pharmacy. *Pharm World Sci.* 2005;27:182–90.
7. Teagarden JR, Nagle B, Aubert RE, Wasdyke C, Courtney P, Epstein RS. Dispensing error rate in a highly automated mail-service pharmacy practice. *Pharmacotherapy.* 2005;25:1629–35.
8. Cina JL, Gandhi TK, Churchill W, Fanikos J, McCrea M, Mitton P, Rothschild JM, Featherstone E, Keohane C, Bates DW, Poon EG. How many hospital pharmacy medication dispensing errors go undetected? *JtComm J Qual Patient Saf.* 2006;32:73–80.
9. Maviglia SM, Yoo JY, Franz C, Featherstone E, Churchill W, Bates DW, Gandhi TK, Poon EG. Cost–benefit analysis of a hospital pharmacy bar code solution. *Arch Intern Med.* 2007;167: 788–94.
10. Ashcroft DM, Quinlan P, Blenkinsopp A. Prospective study of the incidence, nature and causes of dispensing errors in community pharmacies. *Pharmacoepidemiol Drug Saf.* 2005;14: 327–32.
11. Flynn EA, Barker KN, Carnahan BJ. National observational study of prescription dispensing accuracy and safety in 50 pharmacies. *J Am Pharm Assoc (Wash).* 2003;43:191-200.
12. Hoxsie DM, Keller AE, Armstrong EP. Analysis of community pharmacy workflow processes in preventing dispensing errors. *J Pharm Pract.* 2006;19:124-30.
13. Chua SS, Wong IC, Edmondson H, Allen C, Chow J, Peacham J, et al. A feasibility study for recording of dispensing errors and near misses' in four UK primary care pharmacies. *Drug Saf.* 2003;26:803-13.
14. Franklin BD, O'Grady K. Dispensing errors in community pharmacy: Frequency, clinical significance and potential impact of authentication at the point of dispensing. *Int J Pharm Pract* 2007;15:273-81.
15. Allan EL, Barker KN, Malloy MJ, Heller WM. Dispensing errors and counseling in community practice. *Am Pharm.* 1995;NS35:25-33.
16. Flynn EA, Dorris NT, Holman, GT Camahan BJ, Barker KN. Medication dispensing errors in community pharmacies: A nationwide study. *Proc Hum Factor Ergon Soc.* 2002;46:48-51.
17. James KL, Barlow D, McArtney R, Hiom S, Roberts D, Whittlesea C, et al. Incidence, type and causes of dispensing errors: A review of the literature. *Int J Pharm Pract.* 2009;17:9-30.

18. Szeinbach S, Seoane-Vazquez E, Parekh A, Herderick M. Dispensing errors in community pharmacy: Perceived influence of sociotechnical factors. *Int J Qual Health Care*. 2007;19:203-9.
19. Varadarajan R, Barker KN, Flynn EA, Thomas RE. Comparison of two error-detection methods in a mail service pharmacy serving health facilities. *J Am Pharm Assoc*. 2003;2008;48:371-8.
20. Peterson GM, Wu MS, Bergin JK. Pharmacist's attitudes towards dispensing errors: Their causes and prevention. *J Clin Pharm Ther*. 1999;24:57-71.
21. Knudsen P, Herborg H, Mortensen AR, Knudsen M, Hellebek A. Preventing medication errors in community pharmacy: Frequency and seriousness of medication errors. *QualSaf Health Care*. 2007;16:291-6.
22. Leape LL, et al. Systems analysis of adverse drug events. ADE Prevention Study Group. *JAMA*. 1995;74:35-43.
23. Edwards IR, Aronson JK. Adverse drug reactions: definitions, diagnosis, and management. *Lancet*. 2000;356(9237):1255-9.
24. Lazarou J, Pomeranz BH, Corey PN. Incidence of adverse drug reactions in hospitalized patients: A meta-analysis of prospective studies. *JAMA*. 1998;279:1200-5.
25. Juurlink DN, Mamdani M, Kopp A, Laupacis A, Redelmeier DA. Drug-drug interactions among elderly patients hospitalized for drug toxicity. *JAMA*. 2003;289:1652-8.
26. Dirin MM, Mousavi S, Afshari AR, Tabrizian K, Ashrafi MH. Potential drug-drug interactions in prescriptions dispensed in community and hospital pharmacies in East of Iran. *J Res Pharm Pract*. 2014;3(3):104-107.
27. Beso Adnan, Franklin Bryony Dean, Barber Nick. *Pharmacy World And Science* 2005;27:182-90
28. Matos, Vanessa. Evaluation of Drug-Dispensing Errors at the Internal Medicine of an University Hospital. *Latin American Journal of Pharmacy*. 2013;32:26-30.
29. Gothe H, Schall I, Saverno K, et al. The Impact of Generic Substitution on Health and Economic Outcomes: A Systematic Review. *Appl Health Econ Health Policy*. 2015;13:21-33.
30. Paveliu MS, Bengea S, Paveliu FS. Generic Substitution Issues: Brand-generic Substitution, Generic-generic Substitution, and Generic Substitution of Narrow Therapeutic Index (NTI)/Critical Dose Drugs. *Maedica (Bucur)*. 2011;6(1):52-58.
31. AnacletoTâniaAzevedo, Perini Edson, Rosa Mário Borges, César CibeleComini. Drug-dispensing errors in the hospital pharmacy. *Clinics [Internet]*. 2007;62(3):243-250.
32. Ajay Chandra et al. Prescription Audit with Special Emphasis on Drug-Drug Interactions Study in a Private Tertiary Care Teaching Hospital. *Mod ApplBioequiv Availab*. 2017;2(3).

© 2021 Ahamed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/76861>